Neutrosophic Diagram and Classes of Neutrosophic Paradoxes

Florentin Smarandache University of New Mexico, Gallup Campus, USA

Abstract.

These paradoxes are called "neutrosophic" since they are based on indeterminacy (or neutrality, i.e. neither true nor false), which is the third component in neutrosophic logic. We generalize the Venn Diagram to a Neutrosophic Diagram, which deals with vague, inexact, ambiguous, ill-defined ideas, statements, notions, entities with unclear borders. We define the neutrosophic truth table and introduce two neutrosophic operators (*neuterization* and *antonymization* operators) give many classes of neutrosophic paradoxes.

1. Introduction to the Neutrosophics.

Let <A> be an idea, or proposition, statement, attribute, theory, event, concept, entity, and <*nonA*> what is not <A>.

Let $\langle antiA \rangle$ be the opposite of $\langle A \rangle$. We have introduced a new notation [1998], $\langle neutA \rangle$, which is neither $\langle A \rangle$ nor $\langle antiA \rangle$ but in between. $\langle neutA \rangle$ is related with $\langle A \rangle$ and $\langle antiA \rangle$.

Let's see an example for vague (not exact) concepts: if $\langle A \rangle$ is "tall" (an attribute), then $\langle antiA \rangle$ is "short", and $\langle neutA \rangle$ is "medium", while $\langle nonA \rangle$ is "not tall" (which can be "medium or short"). Similarly for other $\langle A \rangle$, $\langle neutA \rangle$, $\langle antiA \rangle$ such as: $\langle good \rangle$, $\langle so so \rangle$, $\langle bad \rangle$, or $\langle perfect \rangle$, $\langle average \rangle$, $\langle imperfect \rangle$, or $\langle high \rangle$, $\langle medium \rangle$, $\langle small \rangle$, or respectively $\langle possible \rangle$, $\langle sometimes possible and other times impossible \rangle$, $\langle impossible \rangle$, etc.

Now, let's take an exact concept / statement: if $\langle A \rangle$ is the statement "1+1=2 in base 10", then $\langle antiA \rangle$ is " $1+1\neq 2$ in base 10", while $\langle neutA \rangle$ is undefined (doesn't exist) since it is not possible to have a statement in between "1+1=2 in base 10" and " $1+1\neq 2$ in base 10" because in base 10 we have 1+1 is either equal to 2 or 1+1 is different from 2. $\langle nonA \rangle$ coincides with $\langle antiA \rangle$ in this case, $\langle nonA \rangle$ is " $1+1\neq 2$ in base 10".

Neutrosophy is a theory the author developed since 1995 as a generalization of dialectics. This theory considers every notion or idea $\langle A \rangle$ together with its opposite or negation $\langle antiA \rangle$, and the spectrum of "neutralities" in between them and related to them, noted by $\langle neutA \rangle$.

<u>The Neutrosophy</u> is a new branch of philosophy which studies the origin, nature, and scope of neutralities, as well as their interactions with different ideational spectra.

Its Fundamental Thesis:

Any idea $\langle A \rangle$ is T% true, I% indeterminate (i.e. neither true nor false, but neutral, unknown), and F% false.

Its Fundamental Theory:

Every idea $\langle A \rangle$ tends to be neutralized, diminished, balanced by $\langle nonA \rangle$ ideas (not only by $\langle antiA \rangle$ as Hegel asserted) - as a state of equilibrium.

In between $\langle A \rangle$ and $\langle antiA \rangle$ there may be a continuous spectrum of particular $\langle neutA \rangle$ ideas, which can balance $\langle A \rangle$ and $\langle antiA \rangle$.

To neuter an idea one must discover all its three sides: of sense (truth), of nonsense (falsity), and of undecidability (indeterminacy) - then reverse/combine them. Afterwards, the idea will be classified as neutrality.

There exists a <u>Principle of Attraction</u> not only between the opposites $\langle A \rangle$ and $\langle antiA \rangle$ (as in dialectics), but also between them and their neutralities $\langle neutA \rangle$ related to them, since $\langle neutA \rangle$ contributes to the <u>Completeness of Knowledge</u>.

Hence, neutrosophy is based not only on analysis of oppositional propositions as dialectic does, but on analysis of these contradictions together with the neutralities related to them.

Neutrosophy was extended to Neutrosophic Logic, Neutrosophic Set, Neutrosophic Probability and Neutrosophic Statistics, which are used in technical applications.

In the <u>Neutrosophic Logic</u> (which is a generalization of fuzzy logic, especially of intuitionistic fuzzy logic) every logical variable *x* is described by an ordered triple x = (T, I, F), where *T* is the degree of truth, *F* is the degree of falsehood, and *I* the degree of indeterminacy (or neutrality, i.e. neither true nor false, but vague, unknown, imprecise), with *T*, *I*, *F* standard or non-standard subsets of the non-standard unit interval $\int 0$, $I^+[$. In addition, these values may vary over time, space, hidden parameters, etc.

<u>Neutrosophic Probability</u> (as a generalization of the classical probability and imprecise probability) studies the chance that a particular event $\langle A \rangle$ will occur, where that chance is represented by three coordinates (variables): T% chance the event will occur, I% indeterminate (unknown) chance, and F% chance the event will not occur.

<u>Neutrosophic Statistics</u> is the analysis of neutrosophic probabilistic events.

<u>Neutrosophic Set</u> (as a generalization of the fuzzy set, and especially of intuitionistic fuzzy set) is a set such that an element belongs to the set with a neutrosophic probability, i.e. T degree of appurtenance (membership) to the set, I degree of indeterminacy (unknown if it is appurtenance or non-appurtenance to the set), and F degree of non-appurtenance (non-membership) to the set.

There exist, for each particular idea: PRO parameters, CONTRA parameters, and NEUTER parameters which influence the above values.

Indeterminacy results from any hazard which may occur, from unknown parameters, or from new arising conditions. This resulted from practice.

2. Applications of Neutrosophics:

Neutrosophic logic/set/probability/statistics are useful in artificial intelligence, neural networks, evolutionary programming, neutrosophic dynamic systems, and quantum mechanics.

3. Examples of Neutrosophy used in Arabic philosophy (F. Smarandache & S. Osman):

- While Avicenna promotes the idea that the world is contingent if it is necessitated by its causes, Averroes rejects it, and both of them are right from their point of view.

Hence *<A>* and *<antiA>* have common parts.

- Islamic dialectical theology (*kalam*) promoting creationism was connected by Avicenna in an extraordinary way with the opposite Aristotelian-Neoplatonic tradition.

Actually a lot of work by Avicenna falls into the frame of neutrosophy.

- Averroes's religious judges (qadis) can be connected with atheists' believes.

- al-Farabi's metaphysics and general theory of emanation vs. al-Ghazali's Sufi writings and mystical treatises [we may think about a coherence of al-Ghazali's "Incoherence of the Incoherence" book].

- al-Kindi's combination of Koranic doctrines with Greek philosophy.

- Islamic Neoplatonism + Western Neoplatonism.

- Ibn – Khaldun's statements in his theory on the cyclic sequence of civilizations, says that:

Luxury leads to the raising of civilization (because the people seek for comforts of life)

but also Luxury leads to the decay of civilization (because its correlation with the corruption of ethics).


- On the other hand, there's the method of absent–by–present syllogism in jurisprudence, in which we find the same principles and laws of neutrosophy.

- In fact, we can also function a lot of Arabic aphorisms, maxims, Koranic miracles (Ayat Al-Qur'ãn) and Sunna of the prophet, to support the theory of neutrosophy.

Take the colloquial proverb that "The continuance of state is impossible" too, or "Everything, if it's increased over its extreme, it will turn over to its opposite"!

4. The Venn Diagram.

In a Venn Diagram we have with respect to a universal set U the following:



Therefore, there are no common parts amongst $\langle A \rangle$, $\langle neutA \rangle$, and $\langle antiA \rangle$, and all three of them are (completely) contained by the universal set U. Also, all borders of these sets $\langle A \rangle$, $\langle neutA \rangle$, $\langle antiA \rangle$, and U are clear, exact. All these four sets are well-defined.

While *<neutA>* means neutralities related to *<A>* and *<antiA>*, what is outside of

 $<A> \cup < neutA> \cup < antiA>$ but inside of U are other neutralities, not related to <A> or to < antiA>.


Given $\langle A \rangle$, there are two types of neutralities: those related to $\langle A \rangle$ (and implicitly related to $\langle antiA \rangle$), and those not related to $\langle A \rangle$ (and implicitly not related to $\langle antiA \rangle$).

5. The Neutrosophic Diagram, as extension of the Venn Diagram.

Yet, for ambiguous, vague, not-well-known (or even unknown) imprecise ideas / notions / statements / entities with unclear frontiers amongst them the below relationships may occur because between an approximate idea noted by $\langle A \rangle$ and its opposite $\langle antiA \rangle$ and their neutralities $\langle neutA \rangle$ there are not clear delimitations, not clear borders to distinguish amongst what is $\langle A \rangle$ and what is not $\langle A \rangle$. There are buffer zones in between $\langle A \rangle$ and $\langle antiA \rangle$ and $\langle neutA \rangle$, and an element x from a buffer zone between $\langle A \rangle$ and $\langle antiA \rangle$ may or may not belong to both $\langle A \rangle$ and $\langle antiA \rangle$ simultaneously. And similarly for an element y in a buffer zone between $\langle A \rangle$ and $\langle antiA \rangle$ and $\langle antiA \rangle$, or to $\langle antiA \rangle$. We may have a buffer zone where the confusion of appurtenance to $\langle A \rangle$, or to $\langle neutA \rangle$, or to $\langle antiA \rangle$ is so high, that we can consider that an element w belongs to all of them simultaneously (or to none of them simultaneously).

We say that all four sets $\langle A \rangle$, $\langle neutA \rangle$, $\langle antiA \rangle$, and the neutrosophic universal set U are illdefined, inexact, unknown (especially if we deal with predictions; for example if $\langle A \rangle$ is a statement with some degree of chance of occurring, with another degree of change of not occurring, plus an unknown part). In the general case, none of the sets $\langle A \rangle$, $\langle neutA \rangle$, $\langle antiA \rangle$, $\langle nonA \rangle$ are completely included in U, and neither U is completely known; for example, if U is the neutrosophic universal set of some specific given events, what about an unexpected event that might belong to U? That's why an approximate U (with vague borders) leaves room for expecting the unexpected.

The <u>Neutrosophic Diagram</u> in the general case is the following:




The borders of *<A>*, *<antiA>*, and *<neutA>* are dotted since they are unclear.

Similarly, the border of the neutrosophic universal set U is dotted, meaning also unclear, so U may not completely contain $\langle A \rangle$, nor $\langle neutA \rangle$ or $\langle antiA \rangle$, but U "approximately" contains each of them. Therefore, there are elements in $\langle A \rangle$ that may not belong to U, and the same thing for $\langle neutA \rangle$ and $\langle antiA \rangle$. Or elements, in the most ambiguous case, there may be elements in $\langle A \rangle$ and in $\langle neutA \rangle$ and in $\langle antiA \rangle$ which are not contained in the universal set U. Even the neutrosophic universal set is ambiguous, vague, and with unclear borders.

Of course, the intersections amongst $\langle A \rangle$, $\langle neutA \rangle$, $\langle antiA \rangle$, and U may be smaller or bigger or even empty depending on each particular case.

See below an example of a particular neutrosophic diagram, when some intersections are contained by the neutrosophic universal set:



A neutrosophic diagram is different from a Venn diagram since the borders in a neutrosophic diagram are vague. When all borders are exact and all intersections among $\langle A \rangle$, $\langle neutA \rangle$, and $\langle antiA \rangle$ are empty, and all $\langle A \rangle$, $\langle neutA \rangle$, and $\langle antiA \rangle$ are included in the neutrosophic universal set *U*, then the neutrosophic diagram becomes a Venn diagram.

The neutrosophic diagram, which complies with the neutrosophic logic and neutrosophic set, is an extension of the Venn diagram.

6. Classes of Neutrosophic Paradoxes.

The below classes of neutrosophic paradoxes are not simply word puzzles. They may look absurd or unreal from the classical logic and classical set theory perspective.

If $\langle A \rangle$ is a precise / exact idea, with well-defined borders that delimit it from others, then of course the below relationships do not occur.

But let $\langle A \rangle$ be a vague, imprecise, ambiguous, not-well-known, not-clear-boundary entity, $\langle nonA \rangle$ means what is not $\langle A \rangle$, and $\langle antiA \rangle$ means the opposite of $\langle A \rangle$.

<*neutA*> means the neutralities related to <*A*> and <*antiA*>, neutralities which are in between them.

When $\langle A \rangle$, $\langle neutA \rangle$, $\langle antiA \rangle$, $\langle nonA \rangle$, U are uncertain, imprecise, they may be selfcontradictory. Also, there are cases when the distinction between a set and its elements is not clear.

Although these neutrosophic paradoxes are based on 'pathological sets' (those whose properties are considered atypically counterintuitive), they are not referring to the theory of Meinongian objects (*Gegenstandstheorie*) such as round squares, unicorns, etc. Neutrosophic paradoxes are not reported to objects, but to vague, imprecise, unclear ideas or predictions or approximate notions or attributes from our everyday life.

- 7. Let's introduce for the first time two new Neutrosophic Operators.
- An operator that "neuterizes" an idea. To neuterize [neuter+ize, transitive verb; from the Latin word neuter = neutral, in neither side], n(.), means to map an entity to its neutral part. {We use the Segoe Print for "n(.)".}

"To neuterize" is different from "to neutralize" [from the French word *neutraliser*] which means to declare a territory neutral in war, or to make ineffective an enemy, or to destroy an enemy.

 $n(\langle A \rangle) = \langle neutA \rangle$. By definition $n(\langle neutA \rangle) = \langle neutA \rangle$.

For example, if $\langle A \rangle$ is "tall", then n(tall) = medium, also n(short) = medium, n(medium) = medium.

But if $\langle A \rangle$ is "l+l=2 in base 10" then $n(\langle l+l=2 \text{ in base } 10 \rangle)$ is undefined (does not

exist), and similarly $n(<1+1\neq 2 \text{ in base } 10>)$ is undefined.

And an operator that "antonymizes" an idea. To antonymize [antonym+ize, transitive verb; from the Greek work antōnymia = instead of, opposite], a(.), means to map an entity to its opposite. {We use the Segoe Print for "a(.)".}

 $a(\langle A \rangle) = \langle antiA \rangle.$

For example, if $\langle A \rangle$ is "tall", then a(tall) = short, also a(short) = tall, and a(medium) = tall or short.

But if <A> is "1+1=2 in base 10" then $a(<1+1=2 \text{ in base } 10>) = <1+1\neq 2$ in base 10> and reciprocally $a(<1+1\neq 2 \text{ in base } 10>) = <1+1=2$ in base 10>.

The classical operator for <u>negation / complement</u> in logics respectively in set theory, "to negate" (\neg) , which is equivalent in neutrosophy with the operator "to nonize" (i.e. to non+ize) or nonization (i.e. non+ization), means to map an idea to its neutral or to its opposite (a union of the previous two neutrosophic operators: neuterization and antonymization):

 $\neg \langle A \rangle = \langle nonA \rangle = \langle neutA \rangle \cup \langle antiA \rangle = n(\langle A \rangle) \cup a(\langle A \rangle).$

Neutrosophic Paradoxes result from the following <u>neutrosophic logic / set connectives</u> following all apparently impossibilities or semi-impossibilities of neutrosophically connecting $\langle A \rangle$, $\langle antiA \rangle$, $\langle neutA \rangle$, $\langle nonA \rangle$, and the neutrosophic universal set U.

8. Neutrosophic Truth Tables.

For <*A*> = "*tall*":

<a>	a (<a>)	n(<a>)	¬ <a>
tall medium short	short or tall	medium	short or medium short or tall tall or medium

To remark that $n(< medium >) \triangleq medium$.

If $\langle A \rangle = tall$, then $\langle neutA \rangle = medium$, and $\langle neut(neutA) \rangle = \langle neutA \rangle$, or $n(\langle n(\langle A \rangle) \rangle) = n(\langle A \rangle)$.

For $\langle A \rangle = (l+1=2 \text{ in base } 10)$ we have $\langle antiA \rangle = \langle nonA \rangle = (l+1\neq 2 \text{ in base } 10)$, while $\langle neutA \rangle$ is undefined (N/A) - whence the neutrosophic truth table becomes:

<a>	a (<a>)	n(<a>)	$\neg $
True	False	N/A	False
False	True	N/A	True

In the case when a statement is given by its neutrosophic logic components $\langle A \rangle = (T, I, F)$, i.e. $\langle A \rangle$ is T% true, I% indeterminate, and F% false, then the neutrosophic truth table depends on the defined neutrosophic operators for each application.

9. Neutrosophic Operators and Classes of Neutrosophic Paradoxes.

a) Complement/Negation

 $\neg <A \ge \neq < nonA \ge$ and reciprocally $\neg < nonA \ge \neq <A \ge$.

 $\neg (\neg \langle A \rangle) \neq \langle A \rangle$ $\neg (\neg \langle antiA \rangle) \neq \langle antiA \rangle$ $\neg (\neg \langle nonA \rangle) \neq \langle nonA \rangle$ $\neg (\neg \langle neutA \rangle) \neq \langle neutA \rangle$ $\neg (\neg U) \neq U$, where U is the neutrosophic universal set. $\neg (\neg \langle \emptyset \rangle) \neq \langle \emptyset \rangle$, where $\langle \emptyset \rangle$ is the neutrosophic empty set.

b) Neuterization.

 $< n(<A>) \neq < neutA>$ $< n(<antiA>) \neq < neutA>$ $< n(< nonA>) \neq < neutA>$ $< n(n(<A>)) \neq <A>$

c) Antonymization.

 $< a(<A>) \neq < antiA>$ $< a(<antiA>) \neq <A>$ $< a(<nonA>) \neq <A>$

 $< a(a(<A>)) \neq <A>$

d) Intersection/Conjunction

 $<A> \cap < nonA> \neq \emptyset$ (neutrosophic empty set) {symbolically $(\exists x)(x \in A \land x \in \neg A)$ }, or even more $<A> \cap < antiA> \neq \emptyset$ {symbolically $(\exists x)(x \in A \land x \in a(A))$ }, similarly $<A> \cap < neutA> \neq \emptyset$ and $< antiA> \cap < neutA> \neq \emptyset$, up to $<A> \cap < neutA> \cap < antiA> \neq \emptyset$. The symbolic notations will be in a similar way. This is <u>Neutrosophic Transdisciplinarity</u>, which means to find common features to uncommon entities

For examples:

There are things which are good and bad in the same time.

There are things which are good and bad and medium in the same time (because from one point of view they may be god, from other point of view they may be bad, and from a third point of view they may be medium).

e) Union / Weak Disjunction

 $<A> \cup < neutA> \cup < antiA> \neq U.$ $< antiA> \cup < neutA> \neq < nonA>.$ Etc.

f) Inclusion/Conditional

 $<A>\subset <antiA>$ $(\forall x)(x \in A \rightarrow x \in a(A))$ All is <antiA>, the <A> too. All *good* things are also *bad*. All is *imperfect*, the *perfect* too.

 $< antiA > \subset <A >$ $(\forall x)(x \in a(A) \rightarrow x \in A)$ All is $\langle A \rangle$, the $\langle antiA \rangle$ too.

All *bad* things have something *good* in them {this is rather a fuzzy paradox}. All is *perfect* things are *imperfect* in some degree.

 $< nonA > \subset <A >$ $(\forall x)(x \in \neg A \rightarrow x \in A)$ All is <A >, the < nonA > too. All bad things have something good

All *bad* things have something *good* and something *medium* in them {this is a neutrosophic paradox, since it is based on good, bad, and medium}.

All is *perfect* things have some *imperfectness and mediocrity* in them at some degree.

 $<A> \subset <neutA>$ $(\forall x)(x \in A \rightarrow x \in n(A))$ All is <neutA>, the <A> too.

 $< nonA > \subset < neutA >$ {partial neutrosophic paradox of inclusion} $(\forall x)(x \in \neg A \rightarrow x \in n(A))$ All is < neutA >, the < nonA > too.

<*nonA* $> \subset <$ *antiA*>{partial neutrosophic paradox of inclusion} ($\forall x$)($x \in \neg A \rightarrow x \in a(A)$) All is <*antiA*>, the <*nonA*> too.

 $< antiA > \subset < neutA >$ $(\forall x)(x \in a(A) \rightarrow x \in n(A))$ All is < neutA >, the < antiA > too.

 $<A> \cup <antiA> \subset <neutA>$ $(\forall x)((x \in A \lor x \in a(A)) \rightarrow x \in n(A))$ All is <neutA>, the <A> and <antiA> too.

Paradoxes of some Neutrosophic Arguments

 $<\!\!A \!\!> \rightarrow <\!\!B \!\!> \\<\!\!B \!\!> \rightarrow <\!\!antiA \!\!> \\\cdots \\ \cdots \\ <\!\!A \!\!> \rightarrow <\!\!antiA \!\!>$

Example: too much work produces sickness; sickness produces less work (absences from work, low efficiency); therefore, too much work implies less work (this is a Law of Self-Equilibrium).

g) Equality/Biconditional

<u>Unequal Equalities</u> $\langle A \rangle \neq \langle A \rangle$ which symbolically becomes $(\exists x)(x \in A \leftrightarrow x \notin A)$ or even stronger inequality $(\forall x)(x \in A \leftrightarrow x \notin A)$. Nothing is $\langle A \rangle$, nor even $\langle A \rangle$.

 $\langle antiA \rangle \neq \langle antiA \rangle$ which symbolically becomes $(\exists x)(x \in \neg A \leftrightarrow x \notin \neg A)$ or even stronger inequality $(\forall x)(x \in \neg A \leftrightarrow x \notin \neg A)$

<*neutA* $> \neq <$ *neutA*>which symbolically becomes $(\exists x)(x \in vA \leftrightarrow x \notin vA)$ or even stronger inequality $(\forall x)(x \in vA \leftrightarrow x \notin vA)$

<*nonA* $> \neq <$ *nonA*>which symbolically becomes $(\exists x)(x \in \neg A \leftrightarrow x \notin \neg A)$ or even stronger inequality $(\forall x)(x \in \neg A \leftrightarrow x \notin \neg A)$

Equal Inequalities $\langle A \rangle = \langle antiA \rangle$ $(\forall x)(x \in A \leftrightarrow x \in a(A))$ All is $\langle A \rangle$, the $\langle antiA \rangle$ too; and reciprocally, all is $\langle antiA \rangle$, the $\langle A \rangle$ too. Or, both combined implications give: All is $\langle A \rangle$ is equivalent to all is $\langle antiA \rangle$.

And so on: <*A*> = <*neutA*> <*antiA*> = <*neutA*> <*nonA*> = <*A*>

 $\begin{array}{l} \hline Dilations and Absorptions \\ <antiA> = <nonA>, \\ \mbox{which means that } <antiA> \mbox{ is dilated to its neutrosophic superset } <nonA>, \mbox{ or } <nonA> \mbox{ is absorbed to its neutrosophic subset } <antiA>. \\ \mbox{Similarly for:} \\ <neutA> = <nonA> \\ <A> = U \\ <neutA> = U \\ <antiA> = U \\ <antiA> = U \end{array}$

< nonA > = U

h) Combinations of the previous single neutrosophic operator equalities and/or inequalities, resulting in more neutrosophic operators involved in the same expression.

For examples:

```
<neutA> \cap (<A> \cup <antiA>) \neq Ø {two neutrosophic operators}.
<A> \cup <antiA> \neq \neg <neutA> and reciprocally \neg (<A> \cup <antiA>) \neq <neutA>.
<A> \cup <neutA> \neq \neg <antiA> and reciprocally.
\neg (<A> \cup <neutA> \cup <antiA>) \neq Ø and reciprocally.
```

Etc.

- i) We can also take into consideration other logical connectors, such as *strong disjunction* (we previously used the *weak disjunction*), *Shaffer's connector*, *Peirce's connector*, and extend them to the neutrosophic form.
- j) We may substitute $\langle A \rangle$ by some entities, attributes, statements, ideas and get nice neutrosophic paradoxes, but not all substitutions will work properly.

10. Some particular paradoxes:

A Quantum Semi-Paradox.

Let's go back to 1931 Schrödinger's paper. Saul Youssef writes (flipping a *quantum coin*) in arXiv.org at *quant-ph/9509004*:

"The situation before the observation could be described by the distribution (1/2, 1/2) and after observing heads our description would be adjusted to (1,0). The problem is, what would you say to a student who then asks: "Yes, but what causes (1/2, 1/2) to evolve into (1,0)? How does it happen?" {http://god-does-not-play-dice.net/Adler.html}.

It is interesting.

Actually we can say the same for any probability different from 1: If at the beginning, the probability of a quantum event, P(quantum event) = p, with 0 , and if later the event occurs, we get to <math>P(quantum event) = 1;

but if the event does not occur, then we get P(quantum event) = 0, so still a kind of contradiction.

Torture's paradox.

An innocent person P, who is tortured, would say to the torturer T whatever the torturer wants to hear, even if P doesn't know anything.

So, *T* would receive incorrect information that will work against him/her.

Thus, the torture returns against the torturer.

Paradoxist psychological behavior.

Instead of being afraid of something, say <A>, try to be afraid of its opposite <antiA>, and thus – because of your fear – you'll end up with the <anti<antiA>>, which is <A>.

Paradoxically, <u>negative publicity</u> attracts better than positive one (enemies of those who do negative publicity against you will sympathize with you and become your friends).

Paradoxistically [word coming etymologically from *paradoxism, paradoxist*], to be <u>in opposition</u> is more poetical and interesting than being opportunistic.

At a sportive, literary, or scientific competition, or in a war, to be on the side of the weaker is more challenging but on the edge of chaos and, as in Complex Adoptive System, more potential to higher creation.

Law of Self-Equilibrium.

{Already cited above at the Neutrosophic Inclusion / Conditional Paradoxes.}

 $<A> \rightarrow $ and $ \rightarrow <antiA>$, therefore $<A> \rightarrow <antiA>$!

Example: too much work produces sickness; sickness produces less work (absences from work, low efficiency); therefore, too much work implies less work.

References:

- 1. Ashbacher, Charles, "The Most Paradoxist Mathematician of the World', by Charles T. Le", review in Journal of Recreational Mathematics, USA, Vol. 28(2), 130, 1996-7.
- Begay, Anthony, "The Smarandache Semantic Paradox", Humanistic Mathematics Network Journal, Harvey Mudd College, Claremont, CA, USA, Issue #17, 48, May 1998; Begay, A., *Paradoja Semántica de Smarandache*: "Todo es posible, ¡lo imposible también!", traducido por Sebastian Martin Ruiz, <u>http://www.scribd.com/doc/27275122/Paradoja-Semantica-de-Smarandache-%E2%80%9CTodo-es-posible-%C2%A110-imposible-tambien-%E2%80%9D</u>.
- 3. Burton, Emil, On Smarandache Class of Paradoxes, mss., 1995.
- 4. Carol Horn Greenstein, Dictionary of logical terms and symbols, Van Nostrand Reinhold Co., 1978.
- 5. Crăciun, Gheorghe, *Florentin Smarandache / Creatorul Mişcării Literare Paradoxiste / The Smarandache Semantic Paradox: All is possible, the impossible too!*, in Timpul, Melbourne, Australia, Anul III, Nr. 32, p. 2, 1994.
- 6. Dr. Devaraj, Ramasamy, *Florentin Smarandache set up the paradoxist literary movement*, in <Parnassus of World Poets 1994>, Madras, India, September 1994.
- 7. Hintikka, Jaako, The Logic of Epistemology and the Epistemology of Logic.
- 8. Holden editor, The Smarandache Semantic Paradox: All is possible, the impossible too!, in <Short Fuse>, Santa Barbara, CA, USA, issue 60, p. 1, 1995.
- 9. Bernd Hutschenreuther, *Die Smarandache'sche Klasse von Paradoxien* [German], <u>http://portal.acm.org/citation.cfm?id=1006537</u>.
- 10. Jacquette, Dale, *Logic: The Semantics of Existence and Nonexistence*, Berlin: de Gruyter, 1996.

- 11. Le, Charles T., "The Smarandache Class of Paradoxes", Bulletin of the Transylvania University of Brasov, Vol. 1 (36), New Series, Series B, 7-8, 1994.
- 12. Le, Charles T., "The Smarandache Class of Paradoxes", Bulletin of Pure and Applied Sciences, Delhi, India, Vol. 14 E (No. 2), 109-110, 1995.
- Le, Charles T., "The Most Paradoxist Mathematician of the World: Florentin Smarandache", Bulletin of Pure and Applied Sciences, Delhi, India, Vol. 15E (Maths & Statistics), No. 1, 81-100, January-June 1996.
- 14. Le, Charles T., "The Smarandache Class of Paradoxes", Journal of Indian Academy of Mathematics, Indore, Vol. 18, No. 1, 53-55, 1996.
- 15. Le, Charles T., *The Smarandache Class of Paradoxes / (mathematical poem)*, Henry C. Bunner / An Anthology in Memoriam, Bristol Banner Books, Bristol, IN, USA, 94, 1996.
- 16. Le, C., *Clasa de paradoxuri Smarandache*, in Tempus, Bucharest, Anul III, Nr. 2(5), p. 4, 1994.
- 17. Mirescu, Marian, Totul e posibil ... chiar si imposibilul (Paradox de Florentin Smarandache), Abracadabra, Salinas, CA, USA, p. 19, March 1995.
- 18. Mitroiescu, I., *The Smarandache Class of Paradoxes Applied in Computer Sciences*, <Abstracts of Papers Presented to the American Mathematical Society>, New Jersey, USA, Vol. 16, No. 3, 651, Issue 101, 1995.
- 19. Mitroiescu, Ilie, *The Smarandache Class of Paradoxes*, The Mathematical Gazette, Vol. 79, No. 484, p. 125, 1995.
- 20. Rapauzu, Mircea, *Țara Paradoxurilor*, Scrisoare către Autor, București, 22 Nov. 1994.
- Mudge, Michael R., "A Paradoxist Mathematician: His Function, Paradoxist Geometry, and Class of Paradoxes", Smarandache Notions Journal, Vail, AZ, USA, Vol. 7, No. 1-2-3, 127-129, 1996.
- Popescu, Marian, "A Model of the Smarandache Paradoxist Geometry", Abstracts of Papers Presented to the American Mathematical Society, New Providence, RI, USA, Vol. 17, No. 1, Issue 103, 96T-99-15, 265, 1996.
- 23. Popescu, Titu, "Estetica paradoxismului", Editura Tempus, Bucarest, 26, 27-28, 1995.
- 24. Rotaru, Ion, *Din nou despre Florentin Smarandache*, Vatra, Tg. Mures, Romania, Nr. 2 (299), 93-94, 1996.
- 25. Seagull, Larry, *Clasa de Paradoxuri Semantice Smarandache* (translation), Abracadabra, Salinas, CA, USA, Anul 2, Nr. 20, 2, June 1994.

- 26. Smarandache, Florentin, Neutrosophy. / Neutrosophic Probability, Set, and Logic, American Research Press, Rehoboth, USA, 105 p., 1998; Republished in 2000, 2003, 2005, as A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability and Statistics (fourth edition), American Research Press, 156 p.; Chinese translation by F. Liu, "A Unifying Field in Logics: Neutrosophic Logic. / Neutrosophy, Neutrosophic Set, Neutrosophic Probability and statistics", Xiquan Chinese Branch, 121 p., 2003; Russian partial translation by D. Rabounski: Hexis, Сущность <u>нейтрософии</u>, 32 p., 2006.
- 27. Smarandache, F., "Mixed Non-Euclidean Geometries", Arhivele Statului, Filiala Vâlcea, Rm. Vâlcea, 1969.
- Smarandache F., Osman, S., *Neutrosophy in Arabic Philosophy* [English version], Renaissance High Press (Ann Arbor), 291 pp., 2007; translated into Arabic language by Dr. Osman Salah, 418 pp., *Munsha't al-Ma'arif* Publ. Hse., Jalal Huzie & Partners, Alexandria, Egypt, 2007.
- 29. Smarandache, Florentin, *Mathematical Fancies & Paradoxes*, The Eugene Strens Memorial on Intuitive and Recreational Mathematics and its History, University of Calgary, Alberta, Canada, 27 July 2 August, 1986.
- 30. Florentin Smarandache, *Paradoxul vieții noastre (poem matematic)*, in <Poesis>, Satu Mare, Romania, Nr. 9 (69), September 1995.
- Vasiliu, Florin, "Paradoxism's main roots", Translated from Romanian by Ștefan Benea, Xiquan Publishing House, Phoenix, USA, 64 p., 1994; review in Zentralblatt für Mathematik, Berlin, Germany, No. 5, 830 - 17, 03001, 1996.
- 32. Tilton, Homer B., "Smarandache's Paradoxes", Math Power, Tucson, AZ, USA, Vol. 2, No. 9, 1-2, September 1996.
- 33. Weisstein, Eric W., "Smarandache Paradox", *CRC Concise Enciclopedia of Mathematics*, CRC Press, Boca Raton, FL, 1661, 1998; <u>http://mathworld.wolfram.com/SmarandacheParadox.html</u>.
- 34. Zitarelli, David E., "Le, Charles T. / The Most Paradoxist Mathematician of the World", Historia Mathematica, PA, USA, Vol. 22, No. 4, # 22.4.110, 460, November 1995.
- 35. Zitarelli, David E., "Mudge, Michael R. / A Paradoxist Mathematician: His Function, Paradoxist Geometry, and Class of Paradoxes", Historia Mathematica, PA, USA, Vol. 24, No. 1, #24.1.119, 114, February 1997.