

## **Stellar Anchor Black Holes as the remnants of former Herbig Haro Objects.**

Leo Vuyk,  
Architect,  
Rotterdam, the Netherlands.  
[LeoVuyk@Gmail.com](mailto:LeoVuyk@Gmail.com)

### **Abstract,**

FUNCTION FOLLOWS FORM in Quantum FFF Theory.

The FORM and MICROSTRUCTURE of elementary particles, is supposed to be the origin of FUNCTIONAL differences between Higgs- Graviton- Photon- and Fermion particles. As a consequence, a NEW splitting, accelerating and pairing MASSLESS Black Hole, able to convert vacuum energy (ZPE) into real energy by entropy decrease, seems to be able to explain quick Galaxy- and Star formation, by so called Stellar Anchor Black Holes (SABHs), Herbig Haro Objects, down to Sunspots, Comets and even Ball Lightning.

### **Introduction.**

Quantum FFF theory states, that the vacuum is seeded with fast oscillating massless Higgs particles, oscillating along a complex chiral tetrahedral vacuum lattice, which has the ability to transfer Photon and Graviton FORM information in bunches of oscillations, through the vacuum lattice with the local speed of light. Ref [5,6,8,9]. As a consequence there are no attraction forces on propeller shaped Fermions. Gravity seems to be the result of two different impulse arrays: the massless Higgs impulse array and the (less effective) massless Graviton impulse array. As a consequence Black holes suffer only Higgs impulses and NO gravitons are escaping because the nucleus of the black hole is supposed to be a super dense knot of Higgs particles. (Ref. 2). Only the statistical sum of the different kinds of vacuum impulses from all directions on Fermions are responsible for all (energetic) phenomena in the universe. Ref.[1,2,3,4,9]

The accelerating plasma tail of NEW paradigm black holes (Ref.1,4), and the splitting black holes by smaller pairing black holes is supposed to be the origin of a smooth proliferation of black holes in the universe. The Eagle Spire Nebula, (fig. 5) the Carina and other nebulas, are examples of a smooth proliferation of black holes through Galaxies and Nebula. What we can observe in these nebulas is the pairing and splitting of black holes ending up into multiple Herbig Haro objects as the origin of new stars as the start of small open star clusters.

Our sun is supposed to be also a remnant of a former open star cluster located in between two Stellar Anchor black holes, representing the so called hotspots of a former Herbig Haro object.

The solar planetary system and planets are supposed to be expelled from the ever growing Sun by instable stellar rotation effects.

As a result we may discriminate two different kinds of stars:

Stars accompanied by two or even more Stellar Anchor Black holes ( SABHs) and stars that are not.

As a result we may observe clear differences in the way stars form planetary nebula at the end of their life.

There are Planetary Nebula's (PNs) with and without clear hotspots located on both sides of the symmetrical exploded gas distribution.

Next pages should give visual support for these statements.

## The origine of Double MWC922 interference rings





Figure 1, The Red Square Planetary nebula MWC922, equipped with double cone structure of expelled material including trailing starspots (Micro Black Holes). There are FOUR cones (figure 3) of interferences originated by the two Stellar Anchor black holes (SABHs). These four cones seem to interfere with the TWO cones of expelled material of the star, leading to FOUR rings of interference.

## MWC922 with additional electron jets like Cygnus-A and the resulting e-magnetic ribbon of protection.



If the sun is a remnant of an open star cluster, then only comparable systems are electromagnetic protected. (author: Leo Vuyk)

Figure 2, The same MWC922 image with additional representation of the so called magnetic ribbon of protection for Galaxy Cosmic Rays, as is found to be present around the Sun by NASA's Interstellar Boundary Explorer (IBEX).




Figure 3, Higgs Vacuum structure around a set of two Black holes, as is found to be present around Planetary nebula and early Galaxies and Herbig Haro objects.







Figure 4, Cygnus A, equipped with clear electron jets and Hotspots.




This Exploding Star /Planetary Nebula M2-9, is presenting only two Stellar Anchor black holes located at A and B. The star is located at C. The shape of this system is an example of double ring shaped dark matter/ sunspot gravity influence.



This Exploding Star /Planetary Nebula is presenting even a double pair of Stellar Anchor black holes located at A,B,C and D.. Our solar system is supposed to have also at least two Stellar Anchor black holes located on both sides of the planetary plane..

Figure 5, The Spire Eagle nebula, with two Planetary nebula, examples of the importance of black holes in the universe.

## The Fermion repelling horizon of new black holes



7x Higgs quantum curves (2) are depicted for Fermions (3) with different distance to the BH. At location (10) the spin of the fermions will FLIP due to the propeller shape of the Fermion and an opposing force away from the horizon is created by the Higgs vacuum itself down to location (7). The zone between (7) and (10) ...

Figure 6, Principle of Fermion repulsion around a black hole according to Quantum FFF theory..

## References:

- [1] [viXra:1103.0068](#)  
**Function Follows Form, at the Quantum Scale and Beyond.**  
Authors: Leo Vuyk.
- [2] [viXra:1103.0024](#)  
**Quantum Gravity and Electro Magnetic Forces in FFF-Theory.**  
Authors: Leo Vuyk.  
Category: [Quantum Gravity and String Theory](#)
- [3] [viXra:1103.0015](#)  
**Wavefunction Collapse and Human Choice-Making Inside an Entangled Mirror Symmetrical Multiverse.**  
Authors: Leo Vuyk.  
Category: [High Energy Particle Physics](#)
- [4] [viXra:1103.0011](#)  
**An Alternative Black Hole, Provided with Entropy Decrease and Plasma Creation.**  
Authors: Leo Vuyk  
Category: [High Energy Particle Physics](#)
- [5] [viXra:1103.0002](#)  
**3-Dimensional String Based Alternative Particle Model.**  
Authors: Leo Vuyk.  
Category: [High Energy Particle Physics](#)
- [6] [viXra:1102.0052](#)  
**Construction Principles for Chiral "atoms of Spacetime Geometry".**  
Authors: Leo Vuyk
- [7] [viXra:1102.0054](#)  
**Atomic Nuclear Geometry Based on Magic Number Logic.**  
Authors: Leo Vuyk.  
Category: [Nuclear and Atomic Physics](#)
- [8] [viXra:1102.0056](#)  
**Experiments to Determine the Mass Related Lightspeed Extinction Volume Around the Earth and Around Spinning Objects in the Lab.**  
Authors: Leo Vuyk.
- [9] [viXra:1103.0097](#) **ZPE Zero Point Energy Examples Around Black Holes.**  
Authors: [Leo Vuyk](#)
- [10] **The New God Particle and Free Will.**  
Author: Leo Vuyk, LuLu publishers, 2008.  
ISBN number 978-1-4092-1031-3