

Space Doesn't Expand. New Proof of Hubble's Law and Center of the Universe

Hyoyoung Choi

After the expansion of universe was observed in the 1920s, [1] [2] physicists and astronomers introduced the concept of "space expands" into physics and many observations and research results were used based on this. [3] [4] However, we can't explain why space expands and why it has a specific velocity and is no observations of expansion of space. This study proves that the expansion of the universe and Hubble's law doesn't result from the expansion of space, but is a dynamical result from the movement of galaxies in space. We could confirm that Hubble's law was always valid when the effect of acceleration was smaller than initial velocity. We can define the center of the universe and find it. There is a possibility that 2.7K background radiation is not radiation in the early days of the universe. In that case, we can't estimate that our universe is isotropic and uniform from CMBR. Also, this shows that cosmological redshift comes out from the Doppler effect of light. Expansion of space was explained that it was related to redshift and scale factor. Therefore, it is influencing many areas of astronomy and cosmology. Therefore, if this discovery is true, all matters related to redshift and scale factor should be reviewed.

Keyword: Hubble's law, expansion of space, redshift, Doppler effect, scale factor, Inflation, CMBR
Email : 7icarush7@gmail.com

I. Introduction

By *Lemaître* and Hubble in the 1920s, expansion of universe, redshift of the galaxy, and recession velocity based on Earth were observed, [1] [2] scientists introduced the concept of "space expands" into physics to explain this.

Observed cosmological redshift was similar to the Doppler shift which occurs when the light source becomes further away from the observer in space, but it was replaced with the concept that space itself expands. [3] [4]

From the two facts of observation of all distant galaxies receding with Earth in the center and that Earth isn't the center of universe, it is presumed that cosmological redshift isn't Doppler shift of the galaxy moving in space.

Moreover, because scale factor is separated and marked by the solution of field equation and this can be corresponded to expansion of space, it was thought that observed cosmological redshift results from the expansion of space. [3]

A recent study put some other interpretation on the expansion of space. [5] [6]

However, significant matters related to expansion of space haven't been proved or explained during the 80s until today and these results aren't being observed.

1. Expansion of space isn't an obvious matter.

Thinking of space expanding, there are 3 cases.

- A. Expansion
- B. Contraction

C. Maintenance - Condition without expansion and contraction

These three conditions can be possible and there is no thought that "expansion" among these is the most natural value. If force does not exist, it is natural that any physical quantity has the same value, so "maintenance" is the most natural value.

2.If space expands, the expansion speed of space can vary from - infinity to + infinity. There is no basis that a specific value among these should be chosen.

3.We have never observed the expansion of space.

The physical meaning of "space expands" is that all space expands.

A. Space between an atomic nucleus and electrons also expands.

B. Space between the Sun and Earth also expands.

C. Space between galaxies also expands.

Like the above content, it means that all space expands. Scientist who claim expansion of space, space all expands in A, B, C, but

For A, binding is consisted by electromagnetic force, space actually expands but position is compensated by electromagnetic force in time we don't feel, and therefore it is explained that is why we can't observe that effect. [4]

For B, space between the Sun and Earth expands every second, but position is immediately compensated because the Sun and Earth is strongly combined by gravity and explains that is why we can't observe that effect. [4]

On the other hand, for C, space between galaxies also expands, but it is explained that expansion of space appears in C because their gravitational binding is weak. [4]

It is a possible explanation. However, this is a possible explanation for Hubble's law, but it is clear that we didn't observe the "expansion of space."

Thus, we have never directly observed expansion of space between electrons and protons, and energy loss used in compensation of position was never measured, and was never measured between Earth and Sun either.

4. Expansion of the universe and expansion of space isn't the same concept.

The fact that the universe expands shows that distance between galaxies become further. This can be explained from the expansion of space between galaxies, but this can be explained even when galaxies have +r direction initial speed in condition where space doesn't expand.

5. The metaphor of a balloon is 4 dimensional or 2 dimensional, the observed Hubble's law is an observational matter in 3 dimensional space.

Balloon analogy [4] is just a pedantic metaphor, not a precise explanation.

This study proves that Hubble's law is a natural result from the dynamics of galaxies in 3 dimension and tries to prove the fact that all far away galaxies have recession speed with Earth in the center.

II. Proof of Hubble's law through dynamics

1. After accelerating expansion (inflation) of early universe has almost finished, particles started to have some velocity.

This velocity distribution naturally has higher velocity when it is further away from the center of the universe and has lower velocity when it is closer to the center.

A. Big bang simulation in the zero energy universe [Video for Big bang Simulation] [7] [8]

Even if the velocity of particles is zero in the early universe, there are particles with higher velocity in further areas from the center and particles close to the center have relatively low velocity by accelerating expansion of early universe. When positive mass gravitationally contracts to form a galaxy, momentum must be conserved, so higher initial velocity continues to exist as it becomes further away from the center of universe.




Figure 1: Velocity distribution of galaxies at early universe. Red arrows show the velocity vector of particles. [8] It can be known that the magnitude of velocity vector is bigger as it become further from the center.

B. Natural distribution of velocity in the 3D space Thinking in another way, 3 dimensional space can be divided into 3 areas (from the center) to far, middle, and close area. Even if the velocity of the far area is lower than the middle area, middle area particles exceed far area particles when time passes because the velocity of middle particles are higher. As a result, velocity distribution of particles shows that the velocity of far areas is highest, middle area is second, and the close area becomes third.

C. Velocity distribution when some kind of anti-gravitational source exists If some kind of anti-gravitational source or effect in 3 dimension exists, M exists with even density, the above velocity distribution can exist.

$$m\vec{a} = + \frac{G(\frac{4\pi}{3}r^3\rho)m}{r^2}\hat{r} \quad (1)$$

$$\vec{a} = + \frac{4\pi G}{3}\rho r\hat{r} \quad (2)$$

If anti-gravitational source is evenly distributed in accelerating expansion time like the inflation of early universe, a bigger acceleration a exists as r becomes larger and velocity distribution has a higher velocity as the radius of the universe becomes larger. As a result, higher velocity exists for particles of far area from the center of the universe after accelerating expansion ends.

The 3 explanations shown above mean that higher velocity for larger R (distance from the center of universe) after accelerating expansion in the early universe isn't a peculiar phenomenon. If speed in small area in the early universe distributes from 0 to c and if some time passes, velocity distribution will be in order as above.

2. Derivation of Hubble's law in space without expansion

A. Decelerating expansion time

First to look into the possibility of this model, let's look into the case in which the direction of V_{a1} and V_{b1} is the same.

$$V_{a1} = V_{a0} + (-a_1 t_1) \quad (3)$$

$$V_{b1} = V_{b0} + (-a_1 t_1) \quad (4)$$

V_{a0}, V_{b0} : It is the speed in which A and B galaxy has when an accelerating expansion (like inflation) ends.

$-a_1$: Acceleration by force (maybe gravity) occurred from some unknown energy source. It is the acceleration of decelerating expansion because decelerating expansion seems to have taken place in the early universe. It is actually a function of time. To make the problem simple, we plan to solve the problem making it as a constant.

t_1 : Total time of universe decelerating expansion.

$$R_{a1} = R_{a0} + V_{a0} t_1 - \frac{1}{2} a_1 t_1^2 \quad (5)$$

$$R_{b1} = R_{b0} + V_{b0} t_1 - \frac{1}{2} a_1 t_1^2 \quad (6)$$

The above equations are equations of speed and distance when acceleration is constant.

B. Accelerating expansion time

After decelerating expansion ends, there was a time of accelerating expansion. Acceleration is given as a_2 this time and the duration time is set as t_2 .

$$V_{an} = V_{a1} + a_2 t_2 = (V_{a0} - a_1 t_1) + a_2 t_2 \quad (7)$$

$$V_{bn} = V_{b1} + a_2 t_2 = (V_{b0} - a_1 t_1) + a_2 t_2 \quad (8)$$

V_{an}, V_{bn} is the now speed of galaxy a and galaxy b.

$$\begin{aligned} R_{an} &= R_{a1} + V_{a1} t_2 + \frac{1}{2} a_2 t_2^2 \\ &= R_{a0} + (V_{a0} t_1 - \frac{1}{2} a_1 t_1^2) + (V_{a0} - a_1 t_1) t_2 + \frac{1}{2} a_2 t_2^2 \\ &= R_{a0} + V_{a0} (t_1 + t_2) - a_1 t_1 t_2 - \frac{1}{2} a_1 t_1^2 + \frac{1}{2} a_2 t_2^2 \end{aligned} \quad (9)$$

$$\begin{aligned} R_{bn} &= R_{b0} + V_{b1} t_2 + \frac{1}{2} a_2 t_2^2 \\ &= R_{b0} + (V_{b0} t_1 - \frac{1}{2} a_1 t_1^2) + (V_{b0} - a_1 t_1) t_2 + \frac{1}{2} a_2 t_2^2 \\ &= R_{b0} + V_{b0} (t_1 + t_2) - a_1 t_1 t_2 - \frac{1}{2} a_1 t_1^2 + \frac{1}{2} a_2 t_2^2 \end{aligned} \quad (10)$$

C. Deriving Hubble's law (when direction is the same)

$V_{rel} = V_{bn} - V_{an}$ is the relative speed of galaxy a and galaxy b.

$$\begin{aligned} V_{rel} &= V_{bn} - V_{an} \\ &= (V_{b0} - a_1 t_1) + a_2 t_2 - (V_{a0} - a_1 t_1) - a_2 t_2 = V_{b0} - V_{a0} \end{aligned} \quad (11)$$

$$\begin{aligned} R_{rel} &= R_{bn} - R_{an} = (R_{b0} - R_{a0}) + (V_{b0} - V_{a0})(t_1 + t_2) \\ &\simeq (V_{b0} - V_{a0})t \end{aligned} \quad (12)$$

Because the galaxies or particles in the early universe were concentrated in a very close distance, it can be set as

$$(R_{b0} - R_{a0}) = 0 \quad (13)$$

$$t = t_1 + t_2 \quad (14)$$

This is the age of the universe.

Deriving the relation between V_{rel} and R_{rel} ,

$$V_{rel} = V_{bn} - V_{an} = V_{b0} - V_{a0} = \frac{1}{t} R_{rel} = H R_{rel} \quad (15)$$

It can be known that Hubble's law comes out.

Especially, the Hubble constant is $H=1/t$ and this is a result that the Hubble constant in Hubble's law corresponds to the reciprocal of the age of universe. Considering decelerating expansion and accelerating expansion and movement of relative particles, the actual age of the universe is $0.993t_H$. It is very close to 1. [3]

Therefore, the above model contains simple equation, but has possibility.

Thus, the recession velocity and Hubble's law between galaxies don't come from some vague concept(unknown concept without empirical experience) of "expansion of space" and shows possibility that it comes from a simple movement equation called $R = V_0 t + \int \int a(t) d^2 t$.

In $R = V_0 t + \int \int a(t) d^2 t$, if $a(t)$ (acceleration) is small, this is because a $V_{rel} = \frac{1}{t} R_{rel} = H R_{rel}$ shape Hubble's law comes out.

D. The observation of "all galaxies become further from us and all galaxies have recession velocity from Hubble's law" isn't from the expansion of space, it is result of dynamics that galaxies show.




Figure 2: Hubble's observation of all galaxies receding with Earth in the center

It is assumed that interpretation issues of observation results above applied most in physicists and astronomers

introducing expansion of space. When observed from Earth, it is observed that all galaxies recede from Earth and the recession velocity also follow all relations of $\vec{V} = H\vec{R}$.

To explain this, if position of the Earth is the center of expansion, namely if position of the Earth is the center of universe, this issue can be simply solved but it can be clearly known that Earth isn't the center of the universe from the observation of the universe until now.

It is because Earth isn't the center of the solar system, but is clear to be just a planet and that the solar system isn't the center of the galactic system either.

Therefore, physicists and astronomers had to find a way to explain this and as this couldn't be explained by dynamics, a new concept that "space expands" was introduced. To explain more specifically, it is assumed that the stereotype that Hubble's observation isn't valid in places where expansion isn't in the center had influence.

[Derivation of Hubble's law]




Figure 3: Hubble's law doesn't result from the expansion of space, but is a dynamical result from the movement of galaxies in space.

Set as

$$|\vec{a}_{E1}| = |\vec{a}_{\alpha 1}| = |\vec{a}_{\beta 1}| = |\vec{a}_{\gamma 1}| = |\vec{a}_{\delta 1}| = a_1,$$

$$|\vec{a}_{E2}| = |\vec{a}_{\alpha 2}| = |\vec{a}_{\beta 2}| = |\vec{a}_{\gamma 2}| = |\vec{a}_{\delta 2}| = a_2$$

$$\vec{V}_{E1} = \vec{V}_{E0} - \vec{a}_{E1}t_1 \quad (16)$$

$$\vec{V}_{\alpha 1} = \vec{V}_{\alpha 0} - \vec{a}_{\alpha 1}t_1 \quad (17)$$

$$\vec{V}_{En} = \vec{V}_{E1} + \vec{a}_{E2}t_2 = (\vec{V}_{E0} - \vec{a}_{E1}t_1) + \vec{a}_{E2}t_2 \quad (18)$$

$$= ((V_{E0} - a_1t_1) + a_2t_2)\hat{x}$$

Set as x-axis.

$$\vec{V}_{\alpha n} = \vec{V}_{\alpha 1} + \vec{a}_{\alpha 2}t_2 = (\vec{V}_{\alpha 0} - \vec{a}_{\alpha 1}t_1) + \vec{a}_{\alpha 2}t_2$$

$$= (V_{\alpha 0} - a_1t_1 + a_2t_2)\cos\theta\hat{x} + (V_{\alpha 0} - a_1t_1 + a_2t_2)\sin\theta\hat{y} \quad (19)$$

$$\vec{R}_{En} = \vec{R}_{E1} + \vec{V}_{E1}t_2 + \frac{1}{2}\vec{a}_{E2}t_2^2$$

$$= (\vec{V}_{E0}t_1 - \frac{1}{2}\vec{a}_{E1}t_1^2) + (\vec{V}_{E0} - \vec{a}_{E1}t_1)t_2 + \frac{1}{2}\vec{a}_{E2}t_2^2$$

$$= (t_1 + t_2)\vec{V}_{E0} - t_1t_2\vec{a}_{E1} - \frac{1}{2}t_1^2\vec{a}_{E1} + \frac{1}{2}t_2^2\vec{a}_{E2} \quad (20)$$

$$\vec{R}_{En} = [(t_1 + t_2)V_{E0} - t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2]\hat{x} \quad (21)$$

$$\vec{R}_{\alpha n} = \vec{R}_{\alpha 1} + \vec{V}_{\alpha 1}t_2 + \frac{1}{2}\vec{a}_{\alpha 2}t_2^2$$

$$= (\vec{V}_{\alpha 0}t_1 - \frac{1}{2}\vec{a}_{\alpha 1}t_1^2) + (\vec{V}_{\alpha 0} - \vec{a}_{\alpha 1}t_1)t_2 + \frac{1}{2}\vec{a}_{\alpha 2}t_2^2 \quad (22)$$

$$= (t_1 + t_2)\vec{V}_{\alpha 0} - t_1t_2\vec{a}_{\alpha 1} - \frac{1}{2}t_1^2\vec{a}_{\alpha 1} + \frac{1}{2}t_2^2\vec{a}_{\alpha 2}$$

$$\vec{R}_{\alpha n} = [(t_1 + t_2)V_{\alpha 0} - t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2]\cos\theta\hat{x}$$

$$+ [(t_1 + t_2)V_{\alpha 0} - t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2]\sin\theta\hat{y} \quad (23)$$

[Relative Velocity]

$$\vec{V}_{rel} = \vec{V}_{\alpha n} - \vec{V}_{En}$$

$$= [(V_{\alpha 0}\cos\theta - V_{E0}) + (-a_1t_1 + a_2t_2)(\cos\theta - 1)]\hat{x}$$

$$+ (V_{\alpha 0} - a_1t_1 + a_2t_2)\sin\theta\hat{y} \quad (24)$$

[Relative Distance]

$$\vec{R}_{rel} =$$

$$t\{[(V_{\alpha 0}\cos\theta - V_{E0}) + \frac{1}{t}(-t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2)(\cos\theta - 1)]\hat{x}$$

$$+ [V_{\alpha 0} + \frac{1}{t}(-t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2)]\sin\theta\hat{y}\} \quad (25)$$

$$(t = t_1 + t_2)$$

1) When θ is zero.

$$\vec{V}_{rel} = (V_{\alpha 0} - V_{E0})\hat{x} \quad (26)$$

$$\vec{R}_{rel} = t(V_{\alpha 0} - V_{E0})\hat{x} \quad (27)$$

$$\text{Therefore, } \vec{V}_{rel} = \frac{1}{t}\vec{R}_{rel} = H\vec{R}_{rel}$$

2) When θ is small.

$$\vec{V}_{rel} \simeq (V_{\alpha 0} - V_{E0})\hat{x} + (V_{\alpha 0} - a_1t_1 + a_2t_2)\theta\hat{y} \quad (28)$$

$$\vec{R}_{rel} \simeq t\{(V_{\alpha 0} - V_{E0})\hat{x} + [V_{\alpha 0} + \frac{1}{t}(-t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2)]\theta\hat{y}\} \quad (29)$$

Hubble's law is valid for the 2 following cases.

i) $V_{E0}, V_{\alpha 0} \gg -a_1t_1 + a_2t_2$

When initial speed is much larger than speed change by deceleration and acceleration :

* Because there is high possibility that there was a time of accelerating expansion of the early universe, particles gained high speed after inflation and the galaxy composed by these particles also had high speed, so the above supposition has validity.

* $-a_1t_1 + a_2t_2 \approx 0$: When the effects of deceleration and acceleration are offset by each other.

Considering decelerating expansion and accelerating expansion and movement of relative particles, the actual

age of the universe is $0.993t_H$. It is very close to 1 [3]. Namely, our universe has a state of $-a_1t_1 + a_2t_2 \approx 0$.

* Zero Energy Universe : In principle, the total energy is zero. So deceleration and acceleration terms are small. [9] [10]

$$\begin{aligned} \vec{V}_{rel} &\simeq (V_{\alpha 0} - V_{E0})\hat{x} + (V_{\alpha 0} - a_1t_1 + a_2t_2)\theta\hat{y} \\ &\simeq (V_{\alpha 0} - V_{E0})\hat{x} + V_{\alpha 0}\theta\hat{y} \end{aligned} \quad (30)$$

$$\vec{R}_{rel} \simeq t[(V_{\alpha 0} - V_{E0})\hat{x} + V_{\alpha 0}\theta\hat{y}] \quad (31)$$

Therefore, $\vec{V}_{rel} = \frac{1}{t}\vec{R}_{rel} = H\vec{R}_{rel}$ Hubble's law is valid.

ii) Hubble's law is valid in $t_1 = t_2, a_1 = 3a_2$ condition.

Because the term of decelerating expansion and accelerating expansion is almost similar from the current observation, it can be set as $t_1 = t_2$ [3] [11]. This condition is the result gained from the condition of assuming expansion of space. Therefore, if the result of this study is true, it can be revised.

$$\begin{aligned} \vec{V}_{rel} &\simeq [(V_{\alpha 0}(1 - \frac{\theta^2}{2!}) - V_{E0}) + (-a_1t_1 + a_2t_2)(-\frac{\theta^2}{2!})]\hat{x} \\ &+ (V_{\alpha 0} - a_1t_1 + a_2t_2)\theta\hat{y} \\ &= [(V_{\alpha 0}(1 - \frac{\theta^2}{2}) - V_{E0}) + a_2t_2\theta^2]\hat{x} + (V_{\alpha 0} - 2a_2t_2)\theta\hat{y} \end{aligned} \quad (32)$$

$$\begin{aligned} \vec{R}_{rel} &\simeq t\{[(V_{\alpha 0}(1 - \frac{\theta^2}{2!}) - V_{E0}) \\ &+ \frac{1}{2t_2}(-3a_2t_2^2 - \frac{1}{2}3a_2t_2^2 + \frac{1}{2}a_2t_2^2)(-\frac{\theta^2}{2!})]\hat{x} \\ &+ [V_{\alpha 0} + \frac{1}{2t_2}(-3a_2t_2^2 - \frac{1}{2}3a_2t_2^2 + \frac{1}{2}a_2t_2^2)]\theta\hat{y}\} \\ &= t\{[(V_{\alpha 0}(1 - \frac{\theta^2}{2}) - V_{E0}) + a_2t_2\theta^2]\hat{x} + (V_{\alpha 0} - 2a_2t_2)\theta\hat{y}\} \end{aligned} \quad (33)$$

Therefore, $\vec{V}_{rel} = \frac{1}{t}\vec{R}_{rel} = H\vec{R}_{rel}$ is valid.

3) When θ is big.

$$\begin{aligned} \vec{V}_{rel} &= [(V_{\alpha 0} \cos \theta - V_{E0}) + (-a_1t_1 + a_2t_2)(\cos \theta - 1)]\hat{x} \\ &+ (V_{\alpha 0} - a_1t_1 + a_2t_2) \sin \theta\hat{y} \end{aligned} \quad (34)$$

$$\begin{aligned} \vec{R}_{rel} &= \\ &t\{[(V_{\alpha 0} \cos \theta - V_{E0}) \\ &+ \frac{1}{t}(-t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2)(\cos \theta - 1)]\hat{x} \\ &+ [V_{\alpha 0} + \frac{1}{t}(-t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2)] \sin \theta\hat{y}\} \end{aligned} \quad (35)$$

i) $V_{E0}, V_{\alpha 0} \gg -a_1t_1 + a_2t_2$

$$\vec{V}_{rel} \simeq (V_{\alpha 0} \cos \theta - V_{E0})\hat{x} + V_{\alpha 0} \sin \theta\hat{y} \quad (36)$$

$$\vec{R}_{rel} \simeq t[(V_{\alpha 0} \cos \theta - V_{E0})\hat{x} + V_{\alpha 0} \sin \theta\hat{y}] \quad (37)$$

Therefore, $\vec{V}_{rel} = \frac{1}{t}\vec{R}_{rel} = H\vec{R}_{rel}$ is valid.

ex.1) $\theta = \frac{\pi}{2}$

$$\vec{V}_{rel} = -(V_{E0} - a_1t_1 + a_2t_2)\hat{x} + (V_{\alpha 0} - a_1t_1 + a_2t_2)\hat{y} \quad (38)$$

$$\begin{aligned} \vec{R}_{rel} &= t\{-(V_{E0} + \frac{1}{t}(-t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2))\hat{x} \\ &+ [V_{\alpha 0} + \frac{1}{t}(-t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2)]\hat{y}\} \end{aligned} \quad (39)$$

If, $V_{E0}, V_{\alpha 0} \gg -a_1t_1 + a_2t_2$

$$\vec{V}_{rel} \simeq -V_{E0}\hat{x} + V_{\alpha 0}\hat{y} \quad (40)$$

$$\vec{R}_{rel} \simeq t(-V_{E0}\hat{x} + V_{\alpha 0}\hat{y}) \quad (41)$$

Therefore, $\vec{V}_{rel} = \frac{1}{t}\vec{R}_{rel} = H\vec{R}_{rel}$. Hubble's law is valid.

ex.2) $\theta = \pi$

$$\vec{V}_{rel} = [(-V_{\alpha 0} - V_{E0}) - 2(-a_1t_1 + a_2t_2)]\hat{x} \quad (42)$$

$$\vec{R}_{rel} = t[(-V_{\alpha 0} - V_{E0}) - \frac{2}{t}(-t_1t_2a_1 - \frac{1}{2}t_1^2a_1 + \frac{1}{2}t_2^2a_2)]\hat{x} \quad (43)$$

If, $V_{E0}, V_{\alpha 0} \gg -a_1t_1 + a_2t_2$

$$\vec{V}_{rel} \simeq (-V_{\alpha 0} - V_{E0})\hat{x} \quad (44)$$

$$\vec{R}_{rel} \simeq t(-V_{\alpha 0} - V_{E0})\hat{x} \quad (45)$$

Therefore, $\vec{V}_{rel} = \frac{1}{t}\vec{R}_{rel} = H\vec{R}_{rel}$ is valid.

When initial speed is much larger than velocity change from deceleration and acceleration, Hubble's law is valid in a very wide area. Also this initial speed is the velocity gained from the accelerating expansion of the early universe.

ii) If $t_1 = t_2, a_1 = 3a_2$

$$\begin{aligned} \vec{V}_{rel} &= [(V_{\alpha 0} \cos \theta - V_{E0}) + (-3a_2t_2 + a_2t_2)(\cos \theta - 1)]\hat{x} \\ &+ (V_{\alpha 0} - 2a_2t_2 + a_2t_2) \sin \theta\hat{y} \\ &= [(V_{\alpha 0} \cos \theta - V_{E0}) - 2a_2t_2(\cos \theta - 1)]\hat{x} \\ &+ (V_{\alpha 0} - a_2t_2) \sin \theta\hat{y} \end{aligned} \quad (46)$$

$$\begin{aligned} \vec{R}_{rel} &= t\{[(V_{\alpha 0} \cos \theta - V_{E0}) - 2a_2t_2(\cos \theta - 1)]\hat{x} \\ &+ (V_{\alpha 0} - 2a_2t_2) \sin \theta\hat{y}\} \end{aligned} \quad (47)$$

Therefore, $\vec{V}_{rel} = \frac{1}{t}\vec{R}_{rel} = H\vec{R}_{rel}$ is valid.

E. Direct meaning of proof

1) Hubble's law is valid in a very wide area in 3 dimensional space when the initial speed of galaxies is much larger than the velocity change by deceleration and acceleration (in the same meaning, when velocity change by deceleration and acceleration is smaller compared to initial speed).

2) This means that even though initial velocity isn't much bigger than the effect by deceleration and acceleration, Hubble's law can be valid in some specific condition. For example, $t_1 = t_2, | -a_1 | = 3a_2$

3) Even though Earth isn't the center of the universe, the belief (something not experienced such as "expansion of space") isn't necessarily needed to explain the reason all galaxies recede from Earth.

III. Meaning including proof

Hubble's law isn't a matter only explained by special condition such as "center of the universe" or a new concept that we haven't experienced such as "expansion of space".

Hubble's law is a result of dynamics valid in almost all areas when change of acceleration is small in the universe.

1. Even if $-a_1(t)$ and $+a_2(t)$ is a function of time, Hubble's law is always valid when the effect of decelerating expansion and accelerating expansion is smaller than initial velocity.

To derive the Hubble's law, we presumed decelerating expansion in the early term and accelerating expansion of the later term. $-a_1$ and $+a_2$ was set as a constant in this process. However, more closely speaking, $-a_1$ and $+a_2$ is a function of time.

2. When the effect of decelerating expansion and accelerating expansion has some specific ratio, Hubble's law can be valid.

For example : $t_1 = t_2, |-a_1| = 3a_2$

3. Hubble's law doesn't come from the expansion of space, but results from dynamics from velocity of individual galaxies.




Figure 4: Hubble's law is a dynamical result from the movement of galaxies in 3D space. Two situations are same.

Maybe this would be a good way of looking at it,¹

¹I'd once introduced this thesis to Science Community(<http://www.scienceforums.net/>) and above is an explanation written by a user named Iggy. As the explanation is intuitive, I copy here.




Figure 5: So that all of the red lines are the same size, the ones where the tangent is the line of expansion too. It's important also that the lengths are proportional, so for example OB/OB' is equal to OC/OC' and OA/OA' since velocity from the center needs to be proportional to distance.

4. Therefore, redshift comes from the Doppler shift of light and implies that the existing equation of redshift should be revised.

Existing equation :

$$z = \frac{R_{obs}}{R_{emitted}} - 1 \quad (48)$$

R is scale factor.

Equation by this discovery :

$$z = \frac{\lambda_o}{\lambda_e} - 1 = \sqrt{\frac{1 + \frac{V}{c}}{1 - \frac{V}{c}}} - 1 \quad (49)$$

The two equations show similar results in close galaxies, but show difference in far galaxies. [3] [4]

5. redshift was the role of a ruler measuring the distance of the universe, but if this model is true, the inaccuracy of the existing ruler is implied and all data (including distance, scale factor) through redshift should be reviewed.

6. We can define the center of the universe and find it. (Revival of absolute coordinate system)

A. Direction of center of the universe

Considering homogeneous, isotropy, and dependence of r of gravity, Hubble's law will be well valid as the direction is closer to the perpendicular plane to the center direction. Draw several lines with Earth in the center

8. The change of redshift eq. influences the "discovery that the universe accelerating expands." [12] [13] Therefore, there is necessity to review accelerating expansion.

IV. Inflation and cosmic microwave background radiation

1. Inflation - expansion faster than light

To explain the flatness and horizon problem, expansion faster than light (inflation) was assumed. [14] However, positive energy and negative energy are cancelled in the zero energy universe. In the field equation, if stress-energy term is zero, curvature term is zero. So, zero energy universe is flat. Therefore to explain flatness, there is no need to assume expansion faster than light.

The horizon problem occurs from the wrong Hubble radius which is derived from the assumption that space expands. If particles don't have velocity faster than light, all areas in the early universe will be inside the area of light(radiation) and are all causally connected. Therefore, thermal equilibrium takes place.

Horizon problem doesn't occur and expansion faster than light isn't needed.

2. Cosmic microwave background radiation

A. When 2.7K background radiation is radiation in the early days of the universe

In the density equation of radiation,

$$R^{3(1+w_{rad})}u_{rad} = R^4u_{rad} = u_{rad,0} \quad (59)$$

It is estimated that R^3 was formed due to the volume of the universe increase, and that the rest R^1 was generated because of photon's cosmological redshift. Thus, we can derive a formula, $RT = T_0$, to explain CMBR's temperature.

Likewise, in this model, as the volume of the cosmos increases, the term R^3 is applied as it is. And, as we know, the difference between the redshift based on the Doppler shift model and that resulting from spatial expansion is not significant, as long as $z < 2$. [3]

Strictly speaking, a new relational expression between the redshift and the scale vector R should be derived from this model. Yet, out of R^4 terms, R^3 is equivalent, and R^1 term is to return a similar result, considering the estimation abovementioned. Hence, based on this model as well, CMBR's temperature will turn out to be a similar value.

B. Provided the 2.7K radiation comes from negative masses surrounding galactic structure

As mentioned in the previous article [9] [10], there was a process of pair production between negative and positive masses in the early days of the cosmos, which could explain dark energy and dark matter.

According to pair creation model of negative mass and positive mass, currently, individual galaxies and clusters of galaxies are surrounded by negative masses.

If the temperature of negative masses outside galactic structure is 2.7K, and if they give off radiation, what will happen?

As negative masses surround our galaxy and clusters of galaxies, we on the earth will observe a uniform 2.7K background radiation coming from all directions.

The radiation temperature decreases as $T \propto \frac{1}{a}$, whilst the temperature of matters drops as $T \propto \frac{1}{a^2}$, when it comes to adiabatic expansion of idea gases. [15]

1) Model 1 - temperature of negative mass

Temperature of radition :

$$T \propto \frac{1}{a}T_0 \approx \left(\frac{1}{1100}\right) \times 3000K = 2.72K \quad (60)$$

Temperature of (negative mass) matter :

$$T \propto \frac{1}{a^2}T_0 \approx \left(\frac{1}{1100}\right)^2 \times 3000K = 2.48 \times 10^{-3}K \quad (61)$$

That is, to think of matters as ideal gases so as to find current temperature, the resulting value is very close to 2.72K. It isn't that bad, considering Alpher and Herman yielded approximately 5K. [16]

However, the problem is baryon matters are not same as ideal gases and join to form stars and galaxies. Another problem is that heat sources exist following nuclear reactions. Therefore, baryonic matters do not follow the adiabatic expansion.

Nonetheless, negative masses interact with each other via repulsive interaction, which makes it difficult for them to join each other. This meets the same conditions as ideal gases. Also, up to now, it is estimated that negative masses are limited in attributes relevant to nuclear reactions. [10] In other words, negative masses may be regarded as similar to ideal gases. As a result, the current temperature of negative masses can be approximately 2.7K as aforementioned.

In the adiabatic expansion model of ideal gases, their temperature is lower than 2.7K. However, due to motions around galaxies and energy gained from a range of heat sources, it can rise a bit further, which can explain the 2.7K.

2) Model 2 - Temperature of separation time

$$T = \left(\frac{1}{1100}\right)^2 \times xK = 2.72K \quad (62)$$

$$x = 3, 291, 200K \quad (63)$$

The formula above implies that once negative masses form neutral particles approximately at 3.3million degrees, they will undergo a process different from radiation, and then adiabatic expansion could happen. That is, in that case, negative masses can explain the current temperature 2.72K to the full.

Negative masses are also affected by gravitational potential. So, although CMBR is a radiation coming from negative masses, the energy ratio inferred from BAO and WMAP can be explained. [9]

If this conjecture is right, there is no estimating that our universe is isotropic and uniform from CMBR.

C. The interpretation of CMB might be wrong or nonuniform characteristics are likely to be discovered in CMB in the future.

D. Verification methods

1) Intensity of CMBR

In the standard model, CMBR is a light released from the last scattering surface, and its intensity should be constant at all directions of the 3D sphere. However, if it is a radiation from certain particles surrounding galactic structure, the distribution of masses surrounding our galaxy becomes important. Thus, CMBR intensity may be different although its temperature is same. (intensity= (photon number)/(unit area or angle)

For example, as our galaxy and clusters of galaxies revolve, negative masses are likely to revolve. Accordingly, the distribution of negative masses may be axially symmetrical. In that case, it is possible that the density of negative masses is a bit higher when they are distributed on the plane perpendicular to the axis.

In that case, the intensity of radiation from them is likely to reach a maximal level on the plane perpendicular to the axis. But due to the repulsive gravitational effect between negative masses, negative masses tend to be distributed uniformly. Hence, such an effect is likely to be insignificant.

2) Implicit assumption of CMBR

It is not certain whether this kind of experiment has already been conducted. Still, it need be tested whether the wavelength distribution of 3000k radiation cooling down to 2.7K is completely identical to that of the radiation from a 2.7K black body.

For the experiment, 900K–1200K radiation needs cooling down to 300K or so, and the resulting values(especially wavelength distribution) need be compared with those of radiation from the black body has a 300K.

CMBR is a very important subject in cosmology. Therefore, even minor questions should be verified.

How can we tell the difference between “the radiation from unknown 2.7K particles surrounding our galactic structure” and “the light radiated from the last scattering surface in the early days of the universe” ?

V. Conclusion

From the observation of the universe, we found Hubble’s law and introduced the concept that “space expands” into physics and astronomy to explain this.

However, not only can’t we explain the reason of expansion, expansion velocity also couldn’t be reasonably explained so the expansion of space isn’t an observed fact.

According to these research results, Hubble’s law comes out from dynamics that galaxies from 3 dimensional space has and for effect of acceleration smaller than initial velocity, it can be confirmed that Hubble’s law is valid in a very wide area.

redshift and scale factor by expansion of space has much influence in much areas of astronomy and cosmology. Therefore, we must review all these related matters.

References

- [1] Lematre, Georges. Un univers homogène de masse constante et de rayon croissant rendant compte de la vitesse radiale des nébuleuses extra-galactiques. Annales de la Société Scientifique de Bruxelles A47: 4956. (1927) Bibcode [1927ASSB...47...49L]
- [2] Hubble, Edwin. A Relation between Distance and Radial Velocity among Extra-Galactic Nebulae. Proceedings of the National Academy of Sciences of the United States of America. Volume 15, March 15, 1929: Issue 3, pp. 168-173, communicated January 17, 1929.
[<http://www.pnas.org/content/15/3/168.full.pdf+html>].
- [3] Bradley W. Carroll, Dale A. Ostlie. Introduction to Modern Astrophysics. 2nd Edition. Pearson Education, Inc.. (2007)
- [4] Charles H. Lineweaver and Tamara M.Davis. Misconceptions about the Big Bang. Scientific american.(2005).[Misconceptions about the Big Bang.pdf]
- [5] J.A. Peacock. A diatribe on expanding space (2008). [arxiv:0809.4573v1]
- [6] Emory F. Bunn and David W. Hogg. The kinematic origin of the cosmological redshift. (2009). [arxiv:0808.1081v2]

- [7] Hyoyoung Choi. Birth of the Universe from Zero Energy State. (2010). [<http://www.youtube.com/watch?v=vYEPbCpkLa8>].
- [8] Hyoyoung Choi. Inflation, accelerating expansion with pair creation of negative and positive mass. (2012). [<http://www.youtube.com/watch?v=SRUqQM2FfNU>].
- [9] Hyoyoung Choi, The Change of Gravitational Potential Energy and Dark Energy in the Zero Energy Universe, (2011). [viXra:1110.0019]
- [10] Hyoyoung Choi, Hypothesis of Dark Matter and Dark Energy with Negative Mass, (2009). [viXra:0907.0015]
- [11] E. Komatsu et al. Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation, *Astrophys.J.Suppl.*192:18, (2011).
- [12] Riess, A. G. et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. *Astron. J.* 116, 10091038 (1998).
- [13] Perlmutter, S. et al. Measurements of omega and lambda from 42 high-redshift supernovae. *Astrophys. J.* 517, 565586 (1999).
- [14] A.H. Guth, The inflationary universe: A possible solution to the horizon and flatness problems, *Phys. Rev.*, D23, 347-356,(1980).
- [15] Hans C. Ohanian and Remo Ruffini. *Gravitation and Spacetime.* (W.W. Norton & Co Inc, New York, 1999)
- [16] Alpher, R. A and Herman, R. C. On the Relative Abundance of the Elements. *Physical Review* 74(12):1737-1742. (1948)