

What are the Mathematical and Physical Concepts of “Flat” Euclidean and “Curved” Non-Euclidean Gravitational Fields?

**By
Robert Louis Kemp**

***Super Principia Mathematica
The Rage to Master Conceptual & Mathematical Physics***

www.SuperPrincipia.com

www.Blog.Superprincipia.com

***Flying Car Publishing Company
P.O Box 91861
Long Beach, CA 90809
January 7, 2013***

Abstract

In this paper a general introduction to basic concepts for the geometric description of Euclidean “Flat-Space” Geometry and Non-Euclidean “Curved-Space” Geometry, and Spherically Symmetric Metric equations which are used for describing the causality and motion of the “Gravitational” interaction between mass with vacuum energy space, and the mass interaction with mass.

This paper gives a conceptual and mathematical description of the differential geometry, of flat and curved space, space-time, or gravitational fields, using the “metric theory” mathematics of Euclidean, Minkowski, Einstein, and Schwarzschild, Spherically Symmetric metrics, and geodesic line elements.

This paper postulates a “Vacuum Energy Perfect Fluid” model and a “Dark Matter Force and Pressure” associated with the Non-Euclidean Spherically Symmetric metric equations, and also gives a conceptual and mathematical description and rationale, for selecting the Schwarzschild Metric over the Einstein Metric, as a physical description of the gradient gravitational, field surrounding a localized net inertial mass/matter source.

This paper also gives a ***new generalized mathematical formalism*** for describing “Non-Euclidean” Spherically Symmetric Metrics, of space, space-time, or the gravitational field, using a generalized ***“Metric “Curvature” Coefficient”***.

Keywords: General Relativity, Special Relativity, Einstein Field Equation, Gravitational Field, Coordinate Singularity, Physical Singularity, Gravitation, Dark Energy Gravitation, Black Hole Event Horizon, Spherically Symmetric Metric, Euclidean Geometry, Non-Euclidean Geometry, Minkowski Metric, Einstein Metric, Schwarzschild Metric

Contents

- 1.0 [Introduction](#)
- 1.1 [Introduction to Basic Concepts of Euclidean and Non-Euclidean Geometry and Spherically Symmetric Metrics](#)
- 1.2 [General Spherically Symmetric Metric – Euclidean and Minkowski \(Pseudo-Euclidean\) Metrics](#)
- 1.3 [General Spherically Symmetric Metric – Schwarzschild Metric – Non-Euclidean Metric](#)
- 1.4 [General Spherically Symmetric Metric – Schwarzschild Metric – Non-Euclidean Metric – Dynamic Fluid Pressure Gradient Solution](#)
- 1.5 [General Spherically Symmetric Metric – Einstein Metric – Non-Euclidean Metric – Static Fluid Pressure Gradient Solution](#)
- 1.6 [New Mathematical Formalism – General Spherically Symmetric Metric – Einstein Metric – Non-Euclidean Metric](#)

1. Introduction

This work is written to physicists that are interested in understanding from a conceptual view, the rationale for selecting “Flat Geometry” Euclidean Space, or selecting a “Curved Geometry” Non-Euclidean Space; as causality for gravity, and motion of the “Gravitational” interaction between mass with vacuum energy space, and the mass interaction with mass.

In this paper, I do weave *some of my own theory, ideas, and mathematics* into these well established physics concepts and mathematics; therefore, this work is written for those that have a very good basis and understanding, of the concepts of differential geometry, and General Relativity; to be able to distinguish what is newly proposed, and what is being discussed in general throughout this paper.

Likewise this work is written to physicists that are interested in understanding why a Schwarzschild Spherically Symmetric Metric (ds^2), is preferred over the Einstein Spherically Symmetric Metric (ds^2); which is based on a particular choice of “pressure” and “density” in the universe, and on the surface of the Black Hole Event Horizon, source of a localized gradient gravitational field.

A “Spherically Symmetric Metric” (ds^2) is used for describing the “flat” or “curved” **Differential Geometry of Space, Time, & Surfaces**, of spherically symmetric space, space-time, or gravitational field, in the presence or absence of condensed matter.

In this work I have limited the discussion only to the: Euclidean, Minkowski, Einstein, and Schwarzschild Spherically Symmetric Metrics of space, space-time, or the gravitational field, however there are many other geometric “metric” equations, and theories of gravitation, that are accepted by the mainstream physics. And there are many “Spherically Symmetric Metrics” that are in use in physics today.

In a paper written by M.S.R. Delgaty and Kayll Lake (1998) **“Physical Acceptability of Isolated, Static, Spherically Symmetric, Perfect Fluid Solutions of Einstein’s Equations”**[\[1\]](#), they describe various “Spherically Symmetric Metrics” (ds^2) equations.

M.S.R. Delgaty and Kayll Lake [”\[1\]](#), state,

“It is fair to say then that most of the spherically symmetric perfect fluid “exact solutions” of Einstein’s field equations that are in the literature are of no physical interest.”

And likewise in a paper by, Petarpa Boonserm, Matt Visser, and Silke Weinfurter (2005) “**Generating perfect fluid spheres in general relativity**” [2], they describe that there are over 127 solutions to the “Spherically Symmetric Metrics”.

But only nine (9) of those “metric” equations satisfy the criteria for predicting actual physical measurable results.

The various “**Spherically Symmetric Metric**” (ds^2), equations which are either Euclidean or Non-Euclidean, describes physical and observable results of gravitational interaction between mass and space, and between mass and mass, predicts that the vacuum energy, and inertial matter in motion interact, through a **space, space-time, or gravitational field, that is either flat or curved, and surrounding a localized gravity source, that is either matter dependent, or matter independent**, is described in the following sections of this paper.

My goal is to bring into correlation, the concepts of “Gravitational Force” of the Newtonian style, and the concepts of Non-Euclidean geometric “curvature of space-time” as described by the mathematical differential geometry of the “**Spherically Symmetric Metric**” (ds^2) of space, space-time, or a gravitational field, and due to the presence or absence of mass or matter as the source of a gradient gravitational field, in a localized region of the universe.

Also I will introduce a **new generalized mathematical formalism** for describing the “**Spherically Symmetric Metric**” (ds^2), by expression of a term known as the “**Metric “Curvature” Coefficient**” ($\kappa_{Curvature}$). The “**Metric “Curvature” Coefficient**” ($\kappa_{Curvature}$) is term and quantity, used for describing the “amount of curvature” due to motion through a perfect fluid vacuum energy, and the presence or absence of mass or matter as the source of a gradient gravitational field, in a localized region of the universe.

The Non-Euclidean “**Spherically Symmetric Metric**” (ds^2) which describes the actual physical geometry of space, space-time or a gravitational field, is used to describe space where there is condensed matter, mass, and energy; as will be described in following section of this paper.

Furthermore, the “**Spherically Symmetric Metric**” (ds^2) can describe the space, space-time or a gravitational field, of or surrounding the: universe, stars, planets, galaxies, quasars, electrons, protons, neutrons, atoms, molecules, photons, etc...

1.1. Introduction to Basic Concepts of Euclidean and Non-Euclidean Geometry and Spherically Symmetric Metrics


The goal of this section is to describe an algorithm for describing a general “Spherically Symmetric Metric”, which can predict Euclidean and Non-Euclidean Metrics, in a “flat” or “curved” space, space-time, or gravitational field.

For any surface, embedded in a Euclidean space of three (3) dimensions or higher, it is possible to measure: the length of a curve (ds) on the surface, the angle ($d\Omega_{Map\ \theta\phi}$) between any two curves on the surface, and the area (ds^2) of a region on the surface. This concept is extended into greater than three dimensions with Bernhard Riemann Geometry (1826 - 1866).

This surface embedded (mapping one surface to another surface) structure to any space, space-time, or gravitational field, is encoded infinitesimally in a generalized spherically symmetric metric, on the surface, through geodesic line elements (ds) and area elements (ds^2).

Carl Friedrich Gauss (1777 - 1855) in his “Theorema Egregium” [3], which is Latin for "remarkable theorem", states that Gaussian curvature of a surface can be determined from the measurements of length (ds) on the surface itself.

This “extraordinary” result shows that the Gaussian curvature of a surface can be computed solely in terms of a “**Metric**” (ds^2), and is thus an intrinsic invariant of the surface. The Gaussian curvature is invariant under isometric deformations of the surface.




Gaussian curvature can be used to describe the position, location, or a change in position or location, of any object on a surface in space.

The “remarkable” result predicted by Gauss, is that the curvature does not depend on its surface embedding, and in spite of all bending and twisting deformations, undergone during rotations; a surface can be determined from the measurements of length (ds) on the surface itself, which is described by a Metric (ds^2).

The Gaussian curvature of a surface is independent of any embedding (mapping one surface to another surface) and is unchanged under coordinate transformations. In general the isometrics of all surfaces preserve Gaussian curvature, during bending, twisting, deformations, and rotations; which is described by a Metric (ds^2).

In Riemannian geometry, Gauss's lemma predicts that any small sphere centered at a point in any surface or Riemannian manifold, is perpendicular to every geodesic through the point. The geodesic is describes by a Metric (ds^2) with line element (ds) on the surface of a space, space-time, or gravitational field; that is either a “flat space” geometry or “curved space” geometry.



Getting a good understanding for what a metric describes is crucial for describing the space and time considerations of Special Relativity (SR) and General Relativity (GR). In (SR) and (GR) the concept of the “Metric” (ds^2), is used to describe a line element (ds) on the surface of a space, space-time, or gravitational field; that is either a “flat space” geometry or “curved space” geometry.


A “Euclidean” Metric (ds^2), in Newtonian Gravitation (NG), describes a gravitational field as a “flat space”, and may be thought of as a concept where mass attracts mass, from across the vastness of empty “flat space”.

A “Non-Euclidean” Metric (ds^2), in General Relativity (GR) may be thought of as a generalization of the gravitational field, in a “curved space or space-time”. The curvature is a form of “attraction” and is a result of the presence of the mass bodies in interaction; with a similar analogy to the Newtonian gravitation attraction concept.

The Metric (ds^2) captures all the geometric and causal structure of space, space-time or a gravitational field, which are used to define notions such as distance, volume, curvature, angle, future and past.


For any Gaussian Surface or Riemannian Manifold, the **sum of the angles of any triangle**, described on a surface is given by any Metric (ds^2):

- is equal to 180° if the geometry is **Euclidean**;
- is less than 180° if the geometry is **Non-Euclidean hyperbolic**;
- is greater than 180° if the geometry is **Non-Euclidean elliptic**



This introduces a conceptual difference between the straight lines of the Euclidean geometry, and the curves of Non-Euclidean geometry, which physically bend in space. This "bending" property of space is not a property of the “Euclidean” or “Pseudo-Euclidean” lines elements and mathematics; but is only described by the “Non-Euclidean” lines elements and mathematics.

Thus Euclidean geometry describes straight lines in flat space geometry, and Non-Euclidean describes hyperbolic geometry and elliptic geometry in curved space geometry.



Another way of considering the differences between the Euclidean and Non-Euclidean geometries, are to consider two straight lines, infinitely extended in a two-dimensional plane that are both perpendicular to a third line:

- In **Euclidean “flat” geometry** the lines in a space, space-time or gravitational field remain in a “flat space” at a constant distance from each other, even if extended into infinity, and are known as parallels.


The **Euclidean “flat” geometry** is **independent** of any condensed “mass” or “energy” in that “flat space”.

- In **Non-Euclidean “open” hyperbolic geometry** the lines in a space, space-time or gravitational field “curve away” from each other, increasing in distance as one moves further from the points of intersection with the common perpendicular.

The **Non-Euclidean “open” hyperbolic geometry** is “curvature” that is **dependent** of the condensed “mass” or “energy” in that “curved space”.


- In **Non-Euclidean “Closed” elliptical geometry** the lines in a space, space-time or gravitational field “curve toward” from each other, decreasing in distance as one moves towards the points of intersection.

The **Non-Euclidean “Closed” elliptical geometry** is “curvature” that is **dependent** of the condensed “mass” or “energy” in that “curved space”.



The **“Spherically Symmetric Metric”** (ds^2) which describes the actual physical geometry of space, space-time or a gravitational field, can be used to describe space where there is condensed matter, mass, and energy.

The **“Spherically Symmetric Metric”** (ds^2) can describe the space, space-time or a gravitational field, of or surrounding the: universe, stars, planets, galaxies, quasars, electrons, protons, neutrons, atoms, molecules, photons, etc...




1.2. General Spherically Symmetric Metric – Euclidean and Minkowski (Pseudo-Euclidean) Metrics

In the differential geometry of theoretical physics, the Minkowski space is often contrasted with Euclidean space because they are both considered “flat space” geometry for space, space-time, or the gravitational field.


The Metric (ds^2) which describes the geometry of space, space-time or a gravitational field, can be used to describe space where there is condensed matter, mass, and energy. And can describe the space, space-time or a gravitational field, of or surrounding the: universe, stars, planets, galaxies, quasars, electrons, protons, neutrons, atoms, molecules, photons, etc...

In the differential geometry of theoretical physics, the Euclidean space describes the "ordinary" triangle distance given by the Pythagorean calculation, on a flat space, and describes the physical space, between two points that one would measure with a ruler, and using the Pythagorean formula.



The Euclidean space has only space-like dimensions, and a Minkowski space has the space-like dimensions and one time-like dimension. On an orthonormal basis the **Euclidean Space** is a four-dimensional real vector space with signature (+, +, +), (x, y, z).

The **Euclidean Metric** ($ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2$) is a Spherically Symmetric Metric that describes the differential geometry of “Flat” space/space-time; and is defined as the “**net sum**” of a square differential radial ($dr^2 = c_{\text{Light}}^2 \cdot d\tau^2$) component, and the “invariant” or “co-variant” square differential surface component ($r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2$), of a “Flat” space/space-time.



The **Euclidean Metric** (ds^2) is **independent** of the mass or energy present in a localized space, space-time, or gravitational field.

Spherically Symmetric Metric – Euclidean Metric

1.1

$$ds^2 = \left[dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

$$c_{\text{Light}}^2 \cdot d\tau'^2 = \left[c_{\text{Light}}^2 \cdot d\tau^2 + r^2 \cdot (d\theta_{\text{Latitude}}^2 + \sin^2(\theta_{\text{Latitude}}) \cdot d\phi_{\text{Longitude}}^2) \right]$$


The **Euclidean Space** in four-dimensional Cartesian vector space, with signature (+, +, +), (x, y, z).

1.2

$$ds^2 = \left[dx^2 + dy^2 + dz^2 \right] \rightarrow m^2$$

$$\left[dx'^2 + dy'^2 + dz'^2 \right] = \left[\left[dx^2 + dy^2 + dz^2 \right] + \left[dx_{\text{Map } \theta\phi}^2 + dy_{\text{Map } \theta\phi}^2 + dz_{\text{Map } \theta\phi}^2 \right] \right] \rightarrow m^2$$

The **Minkowski Metric** (ds^2) is a Spherically Symmetric Metric that is considered “Pseudo-Euclidean”. The **Minkowski Metric** ($ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2$) describes the differential geometry of a “Flat” space-time; and is “**net difference**” between the square differential radial ($dr^2 = c_{\text{Light}}^2 \cdot dt^2$) “flat” space-time component, and the “light cone” “space-time square differential surface component ($dr_{\text{Light}}^2 = c_{\text{Light}}^2 \cdot dt^2$), of a “Flat” space-time.



The **Minkowski Metric** (ds^2) is **independent** of the mass or energy present in a localized space, space-time, or gravitational field.

Spherically Symmetric Metric – Minkowski “Pseudo-Euclidean” Metric

1.3

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 \rightarrow m^2$$

$$ds^2 = \left[dr^2 - dr_{\text{Light}}^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

$$ds^2 = \left[dr^2 - c_{\text{Light}}^2 \cdot dt^2 + r^2 \cdot \left(d\theta_{\text{Latitude}}^2 + \sin^2(\theta_{\text{Latitude}}) \cdot d\phi_{\text{Longitude}}^2 \right) \right]$$

$$-c_{\text{Light}}^2 \cdot d\tau'^2 = \left[c_{\text{Light}}^2 \cdot d\tau^2 - c_{\text{Light}}^2 \cdot dt^2 + r^2 \cdot \left(d\theta_{\text{Latitude}}^2 + \sin^2(\theta_{\text{Latitude}}) \cdot d\phi_{\text{Longitude}}^2 \right) \right]$$

On an orthonormal basis the **Minkowski Space** is also a four-dimensional Cartesian vector space with signature $(-, +, +, +)$, $(-t, x, y, z)$.

1.4

$$ds^2 = \left[dr^2 - c_{\text{Light}}^2 \cdot dt^2 + r^2 \cdot \left(d\theta_{\text{Latitude}}^2 + \sin^2(\theta_{\text{Latitude}}) \cdot d\phi_{\text{Longitude}}^2 \right) \right] \rightarrow m^2$$

$$\left[dx'^2 + dy'^2 + dz'^2 \right] = \left[\left[dx^2 + dy^2 + dz^2 \right] - c_{\text{Light}}^2 \cdot dt^2 + \left[dx_{\text{Map } \theta\phi}^2 + dy_{\text{Map } \theta\phi}^2 + dz_{\text{Map } \theta\phi}^2 \right] \right] \rightarrow m^2$$


The Euclidean space has three ordinary dimensions of space, that are combined with a single dimension of “time” that is “space-like”; to form a four-dimensional manifold for representing a flat space-time.

In Euclidean space, two vectors are said to be orthogonal in a flat space. But Minkowski space differs by including hyperbolic-orthogonal actions in that flat space.

Hermann Minkowski (1864 - 1909) in his work “**The Fundamental Equations for Electromagnetic Processes in Moving Bodies**” [7], describes space-time as having three ordinary dimensions of space, that are combined with a single dimension of “time” that is “time-like”; to form a four-dimensional manifold for representing a space-time.

The Minkowski space is then considered a pseudo-Euclidean space, where the orthonormal rotation is also a representation of a hyperbolic rotation.

The Euclidean and Minkowski space, points in space, correspond to events in space-time; and events not on the light-cone are classified by their relation to the apex, of the space, as space-like or time-like.



Both the Euclidean and Minkowski space describe physical systems, over finite distances of a “flat space” geometry for space, space-time, or the gravitational field. Likewise, both the Euclidean and Minkowski space are typically applied to the weak Newtonian gravitational fields, and in the absence of, or independent of a large mass, which is a source of the gravity field.

When the Euclidean or Minkowski space, is being applied to strong gravitational fields, this is still considered “flat space”. The Euclidean or Minkowski space, is a condition of space, space-time, or the gravitational field where mass and the vacuum of space do not interact.

In a region of weak gravity fields, the Minkowski or Euclidean space space-time in that region becomes “flat” not just locally to the source, but is flat extended out to great distances very far away from the weak gravity source.

For this reason Euclidean and Minkowski space is often referred to as “flat” space or “flat” space-time. The Euclidean and Minkowski space is “flat” in the presence of mass and weak gravity fields, and in that same region the space-time is described by a curved 4-dimensional manifold for which the tangent space to any point is a 4-dimensional Minkowski space.

The Pseudo-Euclidean Spherically Symmetric Metrics, predicts that there is a “**Speed of Space or Space-time**” $((v(r,t))^2)$, and a **Speed of Light** (c_{Light}^2) constant.

Space-time – Square of the Speed of Light - Constant

1.5

$$c_{\text{Light}}^2 = \left(\frac{dr_{\text{Light}}^2}{dt^2} \right) = \left(\frac{dr^2}{d\tau^2} \right) = - \left(\frac{ds^2}{d\tau'^2} \right) = \text{Constant} \rightarrow m^2/s^2$$

Space-time – Square of the Speed of Space (Vacuum Energy Velocity)

1.6

$$(v(r,t))^2 = \left(\frac{dr^2}{dt^2} \right) = c_{\text{Light}}^2 \cdot \left(\frac{d\tau^2}{dt^2} \right) \rightarrow m^2/s^2$$

The **Minkowski Metric** (ds^2) is **independent** of the mass or energy present in a localized space, space-time, or gravitational field.

Spherically Symmetric Metric – Minkowski “Pseudo-Euclidean” Metric

1.7

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 \rightarrow m^2$$

$$ds^2 = \left[dr^2 - c_{\text{Light}}^2 \cdot dt^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

$$ds^2 = \left[\left(1 - \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right) \right) \cdot dr^2 + r^2 \cdot \left(d\theta_{\text{Latitude}}^2 + \sin^2(\theta_{\text{Latitude}}) \cdot d\phi_{\text{Longitude}}^2 \right) \right]$$

Although the Euclidean and Minkowski space describes “flat space” it is still a good description for describing the “curvature” of space, space-time, gravitational fields, and gravity forces. Thus the structure of Euclidean and Minkowski space is still valid in the general description of gravitational fields given by General Relativity (GR).

In Einstein’s [9] mathematical description of curvature of space-time of General Relativity Theory, the terms “metric” and “square of a line element” are used interchangeably.

Spherically Symmetric Metric – Einstein “Non-Euclidean” Metric

1.8

$$ds^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu \rightarrow m^2$$

The “Metric Coefficient” terms (μ & ν ; = [1, 2, 3, 4]), can take on values of one (1) to four (4). The value of one (1) represents “space”, the value of (2) represents “latitude direction space”, the value of three (3) represents “longitude direction space”, and the value of four (4) represents “time”.

According to the above “metric” equation, the “line element” describes information about the causal structure of the space-time.

The “metric” or “squared line element” is “time-like” when the **Minkowski/Einstein Metric** ($ds^2 < 0$).

$$ds^2 < 0 \quad ; \text{ when } ; \quad \left[c_{\text{Light}}^2 \cdot dt^2 > \left(dr^2 + r^2 \cdot d\Omega_{Map\ \theta\phi}^2 \right) \right]$$

The “metric” or “squared line element” is “light-like” when the **Minkowski/Einstein Metric** ($ds^2 = 0$).

$$ds^2 = 0 \quad ; \text{ when } ; \quad \left[c_{\text{Light}}^2 \cdot dt^2 = \left(dr^2 + r^2 \cdot d\Omega_{Map\ \theta\phi}^2 \right) \right]$$

The “metric” or “squared line element” is “space-like” when the **Minkowski/Einstein Metric** ($ds^2 > 0$).

$$ds^2 > 0 \quad ; \text{ when } ; \quad \left[c_{\text{Light}}^2 \cdot dt^2 < \left(dr^2 + r^2 \cdot d\Omega_{Map\ \theta\phi}^2 \right) \right]$$

In the case where there is a significant gravitational field present, and in the presence of a large mass, the space-time becomes curved or warped, in the local vicinity of the condensed matter, mass or energy.

Likewise, in the case where there is a significant gravitational field present, you have to abandon the “flat space” Euclidean and Minkowski, space of Special Relativity (SR), in favor of the “curved space” Einstein, Schwarzschild, and Riemannian, space of General Relativity (GR).

Thus in general there are the “flat space” metrics of Euclid and Minkowski, and the “curved space” “Non-Euclidean” metrics of Einstein and Schwarzschild.

The figure below describes a Euclidean Metric (ds^2), and various Spherically Symmetric Minkowski Metrics (ds^2); at various “**speed of space (Vacuum Energy Velocity)**” versus “**speed of light**” ratios $\left(\frac{v(r,t)}{c_{\text{Light}}}\right)$.

The larger the “**speed of space**” to the “**speed of light**” ratio $\left(\frac{v(r,t)}{c_{\text{Light}}}\right)$, the more closely the Minkowski space-time follows exactly the Euclidean space-time!

Next, will be presented a graph that shows the Euclidean Metric, in contrast with the Minkowski/Einstein Metrics at different values of $\left(\frac{v(r,t)}{c_{\text{Light}}}\right)$.

The Euclidean and the Minkowski “Metrics” (ds^2) and geodesic “line elements” (ds), are “**mass independent**” equations that describe the causality of “flat” space, space-time, or the gravitational field.

Spherically Symmetric Metric – Euclidean Metric


1.9

$$ds^2 = c_{\text{Light}}^2 \cdot d\tau'^2 = \left[dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

Spherically Symmetric Metric – Minkowski “Pseudo-Euclidean” Metric

1.10

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = \left[\left(1 - \left(\frac{c_{\text{Light}}^2}{v(r,t)^2} \right) \right) \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$



The graph of the **Euclidean** and **Minkowski Metrics** (ds^2), showing the radial integration limits, which extend from the center of the gradient gravitational field ($r \geq 0$), and into infinite distances ($r < \infty$), is given in this classical form below.


$$0 \leq r < \infty$$

In the graph below the Spherically Symmetric Minkowski Metrics (ds^2) are shown at various velocity ratios:

$$\left(\frac{v(r,t)}{c_{\text{Light}}} \right) = [0.1, 0.13, 0.7, 1.0, 1.3, 2.3, 500,000]$$

And the Surface Curvature components are given by:

$$d\theta_{\text{Latitude}} = \frac{\pi}{2} \quad ; \quad d\phi_{\text{Longitude}} = \pi \quad ; \quad d\Omega_{\text{Map } \theta\phi} = \frac{\pi}{2} \cdot \sqrt{5}$$




1.3. General Spherically Symmetric Metric – Schwarzschild Metric – Non-Euclidean Metric

In this section, it will be described a “new” generalized mathematical formalism for the Schwarzschild and Einstein “Non-Euclidean” Spherically Symmetric Metrics (ds^2). A “Non-Euclidean” Spherically Symmetric Metric (ds^2) is used to describe the causality of “gravity” or curvature of space, space-time, or gravitational field, in the presence of condensed matter, mass, and energy.


“Classical Formalism” Spherically Symmetric Metric – Euclidean Metric – Flat Space, Space-time, or Gravitational Field Metric – *Mass & Energy Independent*

$$ds^2 = \left[dr^2 + r^2 \cdot d\Omega_{Map\theta\phi}^2 \right] \rightarrow m^2 \quad 1.11$$



“New Formalism” Spherically Symmetric Metric – Non-Euclidean Metric – Curved/Warped Space, Space-time, or Gravitational Field – *Mass & Energy Dependent*

$$ds^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu = \left[\kappa_{Curvature} \cdot dr^2 + r^2 \cdot d\Omega_{Map\theta\phi}^2 \right] \rightarrow m^2 \quad 1.12$$



“New Formalism” – **“Metric “Curvature” Coefficient”** ($\kappa_{Curvature}$)

The Non-Euclidean Spherically Symmetric **Schwarzschild Metric** (ds^2) named after Karl Schwarzschild (1916) [10], is a mathematical equation that describes, a real curving or warping of physical space, space-time, or a gradient gravitational field, that is spherically symmetric, surrounding a condensed matter, mass (m_{Net}), or energy source. The **Schwarzschild Metric** (ds^2), is given in spherically symmetric mathematical formalism below;


Spherically Symmetric Metric – Schwarzschild (Non-Euclidean) Metric

1.13

$$ds^2 = \left[\left[\frac{dr^2}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r}\right)\right)} - \left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r}\right)\right) \cdot c_{\text{Light}}^2 \cdot dt^2 \right] + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right]$$

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu \rightarrow m^2$$


The above equation is an analogue of the classical Newtonian theory of gravitation, which corresponds to the gravitational field around a matter source; and in the classical Gauss theory of electricity, which corresponds to the electric field around a charge source.



In Newtonian ($\mathbf{F}_{\text{Self-Gravity-Force}} = \frac{m_{\text{Net}}^2 \cdot \mathbf{G}}{r^2}$), “Euclidean Flat Space” Gravitation

Theory, the gravitational field, and the attraction between bodies, of differing masses, causes the smaller mass body to orbit the larger mass body; and the smaller mass, has a greater acceleration towards the center of the gradient gravitational field system body.

In **Schwarzschild Metric** (ds^2) “*Non-Euclidean Curved Space*” Gravitation Theory, it similarly describes the gravitational field, and the attraction between bodies of differing mass, is caused by the curving or warping of space, space-time, or a gradient gravitational field, by the net inertial mass (m_{Net}), of the gravitation system, “carved” out by the inertial mass (m_{Net}) of the system.



The curvature and gradient gravitational field described by the **Schwarzschild Metric** (ds^2), and warping of space caused by the net inertial mass (m_{Net}) of the system, and this “forces” the smaller mass body to orbit the gradient gravitational field following a geodesic, in the curved space, space-time, or the gradient gravitational field.


The curvature described by the **Schwarzschild Metric** (ds^2), also predicts that the closer a test mass body is towards the center of the gradient gravitational field, the greater the “acceleration” the test mass (m_{Test}) experiences towards the center, of the system.

The **Schwarzschild Metric** (ds^2) is a mathematical equation which describes a “geodesic” world “**line element**” (ds), with spherically symmetric curvature of space-time, in the “presence” of matter or the Net (m_{Net}) Mass, of a localized gradient gravitational field; whose space-time extends from the surface of the Black Hole radius ($r = r_{\text{Schwarzschild}}$) and into infinite ($r = \infty$) distances of that localized space-time gravity field.

The **Schwarzschild Metric** (ds^2) predicts that the curvature of space surrounding a “Black Hole” gravity source is equivalent to an **Inhomogeneous Gradient Gravitational Field** which also obeys the three (3) laws of motion of Keplerian Mechanics, of the gradient gravity field.

The gradient gravitational field, described by the **Schwarzschild** (ds^2) **Metric** is comprised of an infinite series of “spherical shell potentials”, that originate at the “**Black Hole Event Horizon**” spherical volume, and extends into infinite distances, relative to the “Black Hole” source, and center of the gradient gravitational field. This model of the gradient gravity field is also a vortex system.

$$\mathbf{K}_{\text{Gravity}} = \mathbf{m}_{\text{Net}} \cdot \mathbf{G} = d\bar{v}_{\text{Gravity}}^2 \cdot d\bar{r} = d\bar{g}_{\text{Gravity}} \cdot d\bar{r}^2 = d\bar{\omega}_{\text{Gravity_Velocity}} \cdot d\bar{r}^3$$



Copyright © 2012 Super Principia Mathematica – The Rage to Master Conceptual & Mathematical Physics

In general, each and every one of the concentric spherical shells, or energy potentials of the “inhomogeneous” gradient gravitational field, of the gravitational vortex, is a “homogeneous” gravitational field.

The **Schwarzschild Metric** (ds^2), has a Black Hole Event Horizon, at the center of the gravitational field, with a fixed volume ($V_{ol\ B-Hole} = \frac{4 \cdot \pi}{3} \cdot r_{Schwarzschild}^3$) and Schwarzschild Semi-Major Radius ($r_{Schwarzschild} = \frac{2 \cdot m_{Net} \cdot G}{c_{Light}^2}$) of the inhomogeneous gradient gravitational field.

The **Schwarzschild Metric** (ds^2) predicts that the source of the inhomogeneous gradient gravitational field is given by Net inertial Mass (m_{Net}), and is directly proportional to the “Black Hole” Schwarzschild Semi-Major Radius ($r_{Schwarzschild} = \left(\frac{2 \cdot m_{Net} \cdot G}{c_{Light}^2} \right)$), and likewise is directly proportional to a localized infinite series, of extended inhomogeneous gradient gravitational field accelerations ($g_{Gravity} = \left(\frac{m_{Net} \cdot G}{r^2} \right)$), relative to the center of the Black Hole Event Horizon.

$$r_{Schwarzschild} \propto m_{Net} \propto g_{Gravity}$$

The exterior solution of the **Schwarzschild Metric** (ds^2), and exterior condition describes the curved physical space of the inhomogeneous gradient gravitational field, which extends, from the surface of the Black Hole Schwarzschild radius ($r = r_{Schwarzschild} = \left(\frac{2 \cdot m_{Net} \cdot G}{c_{Light}^2} \right)$), and extends away from the source, into infinite ($r = \infty$) distances of space, space-time or the gravity field.

The **Schwarzschild Metric** (ds^2) solution comes in two (2) forms, one is an “exterior solution” gradient gravitational field solution, and the other is an “interior solution” gravitational field solution.

The Exterior solution corresponds to the space outside of the Black Hole Event Horizon, and Schwarzschild Semi-Major Radius ($r_{Schwarzschild}$).

$$r_{Schwarzschild} \leq r \leq \infty \quad : \text{Exterior Solution}$$

The Interior solution corresponds to the space within or inside the Black Hole Event Horizon, and Schwarzschild Semi-Major Radius ($r_{Schwarzschild}$).

$$0 \leq r < r_{Schwarzschild} \quad : \text{Interior Solution}$$

The Schwarzschild and Einstein “Metrics” (ds^2) and geodesic “line elements” (ds), are “**mass dependent**” equations that describe the causality of “curved” space, space-time, or the gravitational field. The **Schwarzschild Metric** (ds^2) is described below.

Spherically Symmetric Metric – Schwarzschild (Non-Euclidean) Metric

$$ds^2 = \left[\left[\frac{dr^2}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r}\right)\right)} - \left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r}\right)\right) \cdot c_{\text{Light}}^2 \cdot dt^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2 \right] \quad 1.14$$

The **Schwarzschild Metric** (ds^2) must also fit the new mathematical formalism, when using the space-time relations:

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu \rightarrow m^2 \quad 1.15$$

Space-time – Square of the Speed of Light

$$c_{\text{Light}}^2 = \left(\frac{dr_{\text{Light}}^2}{dt^2}\right) = \left(\frac{dr^2}{d\tau^2}\right) = -\left(\frac{ds^2}{d\tau'^2}\right) = \text{Constant} \rightarrow m^2/s^2 \quad 1.16$$

Space-time – Square of the Speed of Space (Vacuum Energy Velocity)

$$(v(r,t))^2 = \left(\frac{dr^2}{dt^2}\right) = c_{\text{Light}}^2 \cdot \left(\frac{d\tau^2}{dt^2}\right) \rightarrow m^2/s^2 \quad 1.17$$

$$\left[\begin{array}{l} \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2}\right) = 1 \rightarrow \text{Speed of Space (equals) to Speed of Light} \\ \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2}\right) > 1 \rightarrow \text{Speed of Space (less than) Speed of Light} \\ \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2}\right) < 1 \rightarrow \text{Speed of Space (greater than) Speed of Light} \end{array} \right]$$

New Mathematical Formalism - Generalized Spherically Symmetric Metric

1.18

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu = \left[\kappa_{\text{Curvature}} \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

“New” Schwarzschild Metric “Curvature” Coefficient – ($\kappa_{\text{Curvature SC}}$)

1.19


$$\kappa_{\text{Curvature SC}} = \left[\frac{1 - \left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)^2 \cdot \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right)}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} \right] \rightarrow \text{Unitless}$$

New Mathematical Formalism - Generalized Spherically Symmetric Schwarzschild (Non-Euclidean) Metric

1.20

$$ds^2 = \left[\left[\frac{1 - \left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)^2 \cdot \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right)}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \right] \rightarrow m^2$$

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu = \left[\kappa_{\text{Curvature}} \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right]$$



The **Schwarzschild Metric** (ds^2) must also fit the new mathematical formalism, when the “Speed of Space (Vacuum Energy Velocity)” is equal to the “Speed of Light”; and using the space-time relation ($dt^2 = d\tau^2$):

1.21

$$\left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right) = 1$$

“New” Schwarzschild Metric “Curvature” Coefficient – ($\kappa_{\text{Curvature SC}}$)

1.22

$$\kappa_{\text{Curvature SC}} = \left[\left(\frac{r_{\text{Schwarzschild}}}{r} \right) \cdot \frac{\left(2 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} \right] \rightarrow \text{Unitless}$$

New Generalized Spherically Symmetric Schwarzschild (Non-Euclidean)


Metric - for condition $\left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} = 1 \right)$; Speed of Space = Speed of Light

1.23

Space & Angle (Curvature) – Schwarzschild Metric

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu = \left[\kappa_{\text{Curvature}} \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

$$ds^2 = \left[\left[\left(\frac{r_{\text{Schwarzschild}}}{r} \right) \cdot \frac{\left(2 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} \right] \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$



In all of the “modern literature” written on General Relativity, the **Schwarzschild Metric** (ds^2) yields predictable results in the “exterior solution” gravity field region; which extend from the Black Hole Event Horizon of the gradient gravitational field ($r > r_{\text{Schwarzschild}}$), and into infinite distances ($r < \infty$), is given in this classical form below.

$$r_{\text{Schwarzschild}} < r < \infty$$

The Exterior solution corresponds to the space outside of the Black Hole Event Horizon, and Schwarzschild Semi-Major Radius ($r_{\text{Schwarzschild}}$).

$$r_{\text{Schwarzschild}} \leq r \leq \infty \quad : \text{Exterior Solution}$$

The Interior solution corresponds to the space within or inside the Black Hole Event Horizon, and Schwarzschild Semi-Major Radius ($r_{\text{Schwarzschild}}$).

$$0 \leq r > r_{\text{Schwarzschild}} \quad : \text{Interior Solution}$$

This “modern literature” written on General Relativity, describes that the **Schwarzschild Metric** (ds^2) predicts “Two (2) Singularities” in the localized gradient gravitational field of the metric:

There is one “**Physical Singularity**” located at zero radius ($r = 0$) of the gradient gravitational field. ($ds^2 = \infty$)

And there is a second “**Coordinate Singularity**” located at the Black Hole Event Horizon, Schwarzschild Radius ($r = r_{\text{Schwarzschild}}$) – ($ds^2 = \infty$).


$$\left[\begin{array}{ll} ds^2 = \infty & \text{when } r = 0 & \text{Physical Singularity} \\ ds^2 = \infty & \text{when } r = r_{\text{Schwarzschild}} & \text{Coordinate Singularity} \\ ds^2 = \text{Exterior Solution} & \text{when } r > r_{\text{Schwarzschild}} & \\ ds^2 = \text{Interior Solution} & \text{when } r < r_{\text{Schwarzschild}} & \\ ds^2 = \left[dr^2 - c_{\text{Light}}^2 \cdot dt^2 \right] + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 & \text{when } r \Rightarrow \infty & \end{array} \right]$$

Because the **Schwarzschild Metric** (ds^2) predicts the **“Physical Singularity”** located at zero radius ($r = 0$), and the **“Coordinate Singularity”** located at the Black Hole Event Horizon, Schwarzschild Radius ($r = r_{\text{Schwarzschild}}$), of the gradient gravitational field, this has caused the mainstream physics community to reject the **Schwarzschild Metric** (ds^2), in favor of: Kruskal–Szekeres coordinates, Eddington–Finkelstein coordinates, and Rindler coordinate; and which neither have a **“Coordinate Singularity”**.

Below is a graph of the **Schwarzschild Metric** (ds^2).

The **Schwarzschild Metric** (ds^2) predicts the **“Physical Singularity”** located at zero radius, is a value that approaches zero, as the radius approaches zero. The **“Physical Singularity”** is a natural artifact for any Non-Euclidean metric.

$$(r \rightarrow 0 ; ds^2 \rightarrow 0) \text{ Then } (r = 0 ; ds^2 = \infty) \text{ And } \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right) = 1$$



The **“Coordinate Singularity”** is not a natural artifact for any Non-Euclidean metric. My goal is to find a solution to the **“Coordinate Singularity”** located at the Black Hole Event Horizon, Schwarzschild Radius ($r = r_{\text{Schwarzschild}}$), of the **Schwarzschild Metric** (ds^2).

1.4. General Spherically Symmetric Metric – Schwarzschild Metric – Non-Euclidean Metric – Dynamic Fluid Pressure Gradient Solution

The **Schwarzschild** “Non-Euclidean” Metric (ds^2) describes a “dynamic” differential geometry metric, and a line element, that describes the curvature of space, space-time, and the gravitational field; and must be used in conjunction, with a fluid mechanical model, **Perfect Fluid “Dynamic” Vacuum Energy Solution** for the causality gravitation.

The “**Schwarzschild Metric**” Spherically Symmetric gradient gravitational field vortex system body, describes a **dynamic** “Refraction/Condensing Pressure” ($\Delta P_{\text{Rarefaction-Pressure}}$), which changes in direct proportion to one third, the Inertial Volume Mass Density ($\frac{1}{3} \cdot \rho_{\text{Net}} \cdot \left(1 - \frac{\bar{r}_{\text{Schwarzschild}}}{\bar{r}}\right)$), of the gradient gravitational vortex system body.

$$\Delta P_{\text{Rarefaction-Pressure}} \propto \frac{1}{3} \cdot \rho_{\text{Net}} \cdot \left(1 - \frac{\bar{r}_{\text{Schwarzschild}}}{\bar{r}}\right)$$

The **Schwarzschild** solution, predicts the following equation for the **Isotropic Rarefaction Pressure of Gravitation** ($\Delta P_{\text{Rarefaction-Pressure}}$)

$$\Delta P_{\text{Rarefaction-Pressure}} = \frac{\Delta F_{\text{Rarefaction-Force}}}{\oint dA_{\text{Area}}} = \left[\begin{array}{c} P_{\text{Aether-Gravity-Pressure}} \\ - 2 \cdot P_{\text{Inertial-Gravity-Pressure}} \end{array} \right] \rightarrow \text{kg/m} \cdot \text{s}^2 \quad 1.24$$

$$\Delta P_{\text{Rarefaction-Pressure}} = \frac{\Delta F_{\text{Rarefaction-Force}}}{\oint dA_{\text{Area}}} = \frac{1}{3} \cdot (\rho_{\text{Net}} \cdot \bar{c}_{\text{Light}}^2) \cdot \left(1 - \frac{\bar{r}_{\text{Schwarzschild}}}{\bar{r}}\right) \rightarrow \text{kg/m} \cdot \text{s}^2$$

Substituting the Inertial Volume Mass Density – ($\rho_{\text{Net}} = \frac{m_{\text{Net}}}{V_{\text{ol}}} = \frac{m_{\text{Net}}}{\frac{4\pi}{3} \cdot \bar{r}^3}$)

$$\Delta P_{\text{Rarefaction-Pressure}} = \frac{\Delta F_{\text{Rarefaction-Force}}}{\oint dA_{\text{Area}}} = \frac{1}{4\pi} \cdot \left(\left[\frac{m_{\text{Net}}}{\bar{r}^3} \right] \cdot \bar{c}_{\text{Light}}^2 \right) \cdot \left(1 - \frac{\bar{r}_{\text{Schwarzschild}}}{\bar{r}}\right) \quad 1.25$$

The “Schwarzschild” Spherically Symmetric Metric (ds^2) corresponds to a gradient gravitational vortex system, where the, “Refraction/Condensing Pressure” ($\Delta P_{\text{Rarefaction-Pressure}} = 0$) on the exterior surface, of the Black Hole Event Horizon, is zero; ($\bar{r} = \bar{r}_{\text{Schwarzschild}}$).

Isotropic Rarefaction Pressure of Gravitation ($\Delta P_{\text{Rarefaction-Pressure}}$) at **Black Hole Event Horizon** ($\bar{r} = \bar{r}_{\text{Schwarzschild}}$)

$$\Delta P_{\text{Rarefaction-Pressure}} = \frac{\Delta F_{\text{Rarefaction-Force}}}{\oint dA_{\text{Area}}} = \left[\begin{array}{c} P_{\text{Aether-Gravity-Pressure}} \\ - 2 \cdot P_{\text{Inertial-Gravity-Pressure}} \end{array} \right] = 0 \quad 1.26$$

$$\Delta P_{\text{Rarefaction-Pressure}} = \frac{1}{3} \cdot (\rho_{\text{Net}} \cdot \bar{c}_{\text{Light}}^2) \cdot \left(1 - \frac{\bar{r}_{\text{Schwarzschild}}}{\bar{r}} \right) = 0$$

Therefore I believe that the Schwarzschild Solution is the correct solution where the, “Refraction/Condensing Pressure” ($\Delta P_{\text{Rarefaction-Pressure}} = 0$) on the exterior surface, of the Black Hole Event Horizon, is zero; ($\bar{r} = \bar{r}_{\text{Schwarzschild}}$); and is non-zero everywhere else.

The **Perfect Fluid “Dynamic” Vacuum Energy Solution** above satisfies the Schwarzschild Spherically Symmetric Metric (ds^2) and “line element” (ds); which is a **“mass dependent”** equation that describes the causality of “curved” space, space-time, or the gravitational field. The **Schwarzschild Metric** (ds^2) is described below.

Spherically Symmetric Metric – Schwarzschild (Non-Euclidean) Metric

$$ds^2 = \left[\left[\frac{dr^2}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} - \left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right) \cdot c_{\text{Light}}^2 \cdot dt^2 \right] + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2 \quad 1.27$$

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu = \left[\kappa_{\text{Curvature}} \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

1.5. General Spherically Symmetric Metric – Einstein Metric – Non-Euclidean Metric – Static Fluid Pressure Gradient Solution

Next, we will discuss the **Albert Einstein**, Spherically Symmetric (ds^2) Metric. The **Einstein Metric** (ds^2) is given by the **Static Vacuum Energy Solution** and condition.

The **Einstein** “**Non-Euclidean**” Metric (ds^2) describes a “**static**” geometry and a line element, that describes the curvature of space, space-time, and the gravitational field; and must be used in conjunction, with a fluid mechanical model, **Perfect Fluid “Static” Vacuum Energy Solution** for the causality gravitation.

The **Einstein** solution, predicts the following equation for the **Isotropic Aether Gravitational Field Pressure** ($P_{\text{Aether-Gravity-Pressure}}$)

1.28

$$P_{\text{Aether-Gravity-Pressure}} = \frac{F_{\text{Light-Force}}}{\oint dA_{\text{Area}}} = \frac{1}{3} \cdot (\rho_{\text{Net}} \cdot \bar{c}_{\text{Light}}^2) \rightarrow \text{kg}/\text{m} \cdot \text{s}^2$$

Substituting the Inertial Volume Mass Density – ($\rho_{\text{Net}} = \frac{m_{\text{Net}}}{V_{\text{ol}}} = \frac{m_{\text{Net}}}{\frac{4\pi}{3} \cdot \bar{r}^3}$)

1.29

$$P_{\text{Aether-Gravity-Pressure}} = \frac{F_{\text{Light-Force}}}{\oint dA_{\text{Area}}} = \frac{1}{4\pi} \cdot \left(\left[\frac{m_{\text{Net}}}{\bar{r}^3} \right] \cdot \bar{c}_{\text{Light}}^2 \right) \rightarrow \text{kg}/\text{m} \cdot \text{s}^2$$

The **Einstein** solution, predicts that the **Isotropic Aether Gravitational Field Pressure** ($P_{\text{Aether-Gravity-Pressure}}$) varies in direct proportional to one third the

Inertial Volume Mass Density ($\frac{1}{3} \cdot \rho_{\text{Net}} = \frac{1}{4\pi} \cdot \left(\frac{m_{\text{Net}}}{\bar{r}^3} \right)$) of the gradient gravity

field; and varies inversely proportional to the cube of the distance ($\frac{1}{\bar{r}^3}$), relative to

the center of the gradient gravitational field of a vortex system body.

$$P_{\text{Aether-Gravity-Pressure}} \propto \frac{1}{3} \cdot \rho_{\text{Net}} \propto \frac{1}{4\pi} \cdot \left(\frac{m_{\text{Net}}}{\bar{r}^3} \right)$$

In the **Einstein Metric** (ds^2) **Static Vacuum Energy Solution** the gradient “Rarefaction Pressure” ($P_{\text{Aether-Gravity-Pressure}} = \frac{1}{3} \cdot (\rho_{\text{Net}} \cdot \bar{c}_{\text{Light}}^2)$), is inhomogeneous and distributed in a normalized density gradient, of an infinite series of spherically symmetric shells, of potential energy; including on the surface of the Black Hole Event Horizon.

The “Einstein” Spherically Symmetric Metric (ds^2) corresponds to a gradient gravitational vortex system, where the, “Refraction/Condensing Pressure” ($\Delta P_{\text{Rarefaction-Pressure}} \neq 0$) on the exterior surface, of the Black Hole Event Horizon, is non zero; ($\bar{r} = \bar{r}_{\text{Schwarzschild}}$).

The fluid mechanical model, **Perfect Fluid “Static” Vacuum Energy Solution** describes a condition where there is inhomogeneous gradient “Inertial Volume Mass Density” ($P_{\text{Aether-Gravity-Pressure}} = 2 \cdot P_{\text{Inertial-Gravity-Pressure}} = \frac{1}{3} \cdot (\rho_{\text{Net}} \cdot \bar{c}_{\text{Light}}^2)$), equal to a “static” fixed value, **at the Black Hole Event Horizon**.

Isotropic Rarefaction Pressure of Gravitation ($\Delta P_{\text{Rarefaction-Pressure}} = 0$) at **Black Hole Event Horizon** ($\bar{r} = \bar{r}_{\text{Schwarzschild}}$)

1.30

$$\Delta P_{\text{Rarefaction-Pressure}} = \frac{\Delta F_{\text{Rarefaction-Force}}}{\oint dA_{\text{Area}}} = \left[\begin{array}{c} P_{\text{Aether-Gravity-Pressure}} \\ - 2 \cdot P_{\text{Inertial-Gravity-Pressure}} \end{array} \right] = 0$$

Isotropic Aether Gravitational Field Pressure ($P_{\text{Aether-Gravity-Pressure}}$) at **Black Hole Event Horizon** ($\bar{r} = \bar{r}_{\text{Schwarzschild}}$)

1.31

$$P_{\text{Aether-Gravity-Pressure}} = 2 \cdot P_{\text{Inertial-Gravity-Pressure}} \rightarrow \frac{kg}{m \cdot s^2}$$

$$\frac{1}{3} \cdot (\rho_{\text{Net}} \cdot \bar{c}_{\text{Light}}^2) = \frac{1}{2\pi} \cdot \left(\frac{m_{\text{Net}}^2 \cdot G}{\bar{r}^4} \right) = \frac{1}{4\pi} \cdot (m_{\text{Net}} \cdot \bar{c}_{\text{Light}}^2) \cdot \left(\frac{\bar{r}_{\text{Schwarzschild}}}{\bar{r}^4} \right)$$

$$P_{\text{Aether-Gravity-Pressure}} = \frac{1}{4\pi} \cdot \left(\frac{m_{\text{Net}} \cdot \bar{c}_{\text{Light}}^2}{\bar{r}_{\text{Schwarzschild}}^3} \right) = \frac{1}{32\pi} \cdot \left(\frac{\bar{c}_{\text{Light}}^8}{m_{\text{Net}}^2 \cdot G^3} \right)$$


In the **Static Vacuum Energy Solution** this condition is described by the Spherically Symmetric, mathematics of the **Einstein Metric** (ds^2).

Spherically Symmetric Metric – Einstein (Non-Euclidean) Metric

1.32

$$ds^2 = \left[\left[\frac{dr^2}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r}\right)\right)} - c_{\text{Light}}^2 \cdot dt^2 \right] + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu \rightarrow m^2$$



The **Einstein Metric** (ds^2) solution comes in two (2) forms, one is an “exterior solution” gradient gravitational field solution, and the other is an “interior solution” gravitational field solution.

The Exterior solution corresponds to the space outside of the Black Hole Event Horizon, and Schwarzschild Semi-Major Radius ($r_{\text{Schwarzschild}}$).

$$r_{\text{Schwarzschild}} \leq r \leq \infty \quad : \text{Exterior Solution}$$

The Interior solution corresponds to the space within or inside the Black Hole Event Horizon, and Schwarzschild Semi-Major Radius ($r_{\text{Schwarzschild}}$).

$$0 \leq r < r_{\text{Schwarzschild}} \quad : \text{Interior Solution}$$

The **Einstein Metric** (ds^2) must also fit the new mathematical formalism, when using the space-time relations, for the speed of light, and the speed of space; given below.

Space-time – Square of the Speed of Light

1.33

$$c_{\text{Light}}^2 = \left(\frac{dr_{\text{Light}}^2}{dt^2} \right) = \left(\frac{dr^2}{d\tau^2} \right) = - \left(\frac{ds^2}{d\tau'^2} \right) = \text{Constant} \rightarrow m^2/s^2$$

Space-time – Square of the Speed of Space (Vacuum Energy Velocity)

1.34

$$(v(r,t))^2 = \left(\frac{dr^2}{dt^2} \right) = c_{\text{Light}}^2 \cdot \left(\frac{d\tau^2}{dt^2} \right) \rightarrow m^2/s^2$$

$$\left[\begin{array}{l} \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right) = 1 \rightarrow \text{Speed of Space (equals) to Speed of Light} \\ \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right) > 1 \rightarrow \text{Speed of Space (less than) Speed of Light} \\ \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right) < 1 \rightarrow \text{Speed of Space (greater than) Speed of Light} \end{array} \right]$$

1.6. New Mathematical Formalism – General Spherically Symmetric Metric – Einstein Metric – Non-Euclidean Metric

New Generalized Spherically Symmetric Metric

1.35

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu = \left[\kappa_{\text{Curvature}} \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

“New” Einstein Metric “Curvature” Coefficient – ($\kappa_{\text{Curvature Einstein}}$)

1.36

$$\kappa_{\text{Curvature Einstein}} = \left[\frac{1 - \left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right) \cdot \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right)}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} \right] \rightarrow \text{Unitless}$$


New Generalized Spherically Symmetric Einstein (Non-Euclidean) Metric

1.37

Space & Angle (Curvature) – Einstein Metric

$$ds^2 = \left[\left[\frac{1 - \left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right) \cdot \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right)}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} \right] \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right]$$

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu = \left[\kappa_{\text{Curvature}} \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$



The **Einstein Metric** (ds^2) must also fit must also fit the new mathematical formalism, when the “Speed of Space (Vacuum Energy Velocity)” is equal to the “Speed of Light”; and using the space-time relation ($dt^2 = d\tau^2$).

1.38

$$\left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right) = 1$$

“New” Einstein Metric “Curvature” Coefficient – ($\kappa_{\text{Curvature Einstein}}$)

1.39

$$\kappa_{\text{Curvature Einstein}} = \left[\frac{\left(\frac{r_{\text{Schwarzschild}}}{r} \right)}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} \right] \rightarrow \text{Unitless}$$


New Generalized Spherically Symmetric Einstein (Non-Euclidean) Metric

1.40

Space & Angle (Curvature) – Einstein Metric

$$ds^2 = \left[\left[\frac{\left(\frac{r_{\text{Schwarzschild}}}{r} \right)}{\left(1 - \left(\frac{r_{\text{Schwarzschild}}}{r} \right) \right)} \right] \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$

$$ds^2 = -c_{\text{Light}}^2 \cdot d\tau'^2 = g_{\mu\nu} \cdot dx^\mu \cdot dx^\nu = \left[\kappa_{\text{Curvature}} \cdot dr^2 + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] \rightarrow m^2$$



The **Einstein Metric** (ds^2) yields predictable results in the “exterior solution” gravity field region; which extend from the Black Hole Event Horizon of the gradient gravitational field ($r > r_{\text{Schwarzschild}}$), and into infinite distances ($r < \infty$), is given in this classical form below.

$$r_{\text{Schwarzschild}} < r < \infty$$

The **Einstein Metric** (ds^2) predicts “Two (2) Singularities” in the localized gradient gravitational field of the metric:

There is one “**Physical Singularity**” located at zero radius ($r = 0$) of the gradient gravitational field. ($ds^2 = \infty$)

And there is a second “**Coordinate Singularity**” located at the Black Hole Event Horizon, Schwarzschild Radius ($r = r_{\text{Schwarzschild}}$) – ($ds^2 = \infty$).

$$\left[\begin{array}{lll} ds^2 = \infty & \text{when } r = 0 & \textit{Physical Singularity} \\ ds^2 = \infty & \text{when } r = r_{\text{Schwarzschild}} & \textit{Coordinate Singularity} \\ ds^2 = \textit{Exterior Solution} & \text{when } r > r_{\text{Schwarzschild}} & \\ ds^2 = \textit{Interior Solution} & \text{when } r < r_{\text{Schwarzschild}} & \\ ds^2 = \left[\left[dr^2 - c_{\text{Light}}^2 \cdot dt^2 \right] + r^2 \cdot d\Omega_{\text{Map } \theta\phi}^2 \right] & \text{when } r \Rightarrow \infty & \end{array} \right]$$


Because the **Einstein Metric** (ds^2) predicts a “**Physical Singularity**” located at zero radius ($r = 0$), and a “**Coordinate Singularity**” located at the Black Hole Event Horizon, Schwarzschild Radius ($r = r_{\text{Schwarzschild}}$), of the gradient gravitational field, this has caused the mainstream physics community to reject the **Einstein Metric** (ds^2), in favor of: Kruskal–Szekeres coordinates, Eddington–Finkelstein coordinates, and Rindler coordinate; and which neither have a “**Coordinate Singularity**”.

Below is a graph of the **Einstein Metric** (ds^2).

The **Einstein Metric** (ds^2) predicts the **“Physical Singularity”** located at zero radius, is a value that approaches zero, as the radius approaches zero. The **“Physical Singularity”** is a natural artifact for any Non-Euclidean metric.


$$(r \rightarrow 0 ; ds^2 \rightarrow 0) \quad \text{Then} \quad (r = 0 ; ds^2 = \infty)$$

$$\text{And} \left(\frac{c_{\text{Light}}^2}{(v(r,t))^2} \right) = 1$$




The **“Coordinate Singularity”** is not a natural artifact for any Non-Euclidean metric. My goal is to find a solution to the **“Coordinate Singularity”** located at the Black Hole Event Horizon, Schwarzschild Radius ($r = r_{\text{Schwarzschild}}$), of the **Schwarzschild Metric** (ds^2).

The **Einstein Metric** (ds^2) & **Schwarzschild Metric** (ds^2)



The **Euclidean Metric** (ds^2), **Minkowski Metric** (ds^2), **Einstein Metric** (ds^2), & **Schwarzschild Metric** (ds^2)



2. Conclusion

This work was written to physicists that are interested in understanding from a conceptual view, the rationale for selecting “Flat Geometry” Euclidean Space, or selecting a “Curved Geometry” Non-Euclidean Space; and whether to choose the Einstein Metric or the Schwarzschild Metric, as description for causality of gravity, or general motion in a gravitational field.

This paper described the conceptual and mathematical description of the differential geometry, of flat and curved space, space-time, or gravitational fields, using the “metric theory” mathematics of Euclidean, Minkowski, Einstein, and Schwarzschild, Spherically Symmetric metrics, and geodesic line elements.

This paper also gives a **new generalized mathematical formalism** for describing “Non-Euclidean” Spherically Symmetric Metrics, of space, space-time, or the gravitational field, using a generalized “**Metric “Curvature”** ($\kappa_{Curvature}$) **Coefficient”**.

It was demonstrated that the **Non-Euclidean Spherically Symmetric Metric** (ds^2) equations predicts that there is a “**Physical Singularity**” located at zero radius ($r = 0$), and a “**Coordinate Singularity**” located at the Black Hole Event Horizon, Schwarzschild Radius ($r = r_{Schwarzschild}$), of the gradient gravitational field, in consideration.

The “**Physical Singularity**” is the approaching of an infinity small number as “space and time” approaches zero in the center, of gradient gravitational field of any isolated and localized, gravitational vortex system body, and is a natural artifact of a **Non-Euclidean space, space-time, or gravitational field**.

The “**Coordinate Singularity**” is a “problem”, and is the approaching of an infinity large number as “space and time” approaches the Black Hole Event Horizon, Schwarzschild Radius ($r = r_{Schwarzschild}$), of the gradient gravitational field, of any isolated and localized, gravitational vortex system body. The “**Coordinate Singularity**” is an anomaly, in the mathematics of a particular choice of **Non-Euclidean Spherically Symmetric Metric** (ds^2) equation, used to describe the localized space, space-time, or gravitational field, in consideration.

In another paper I will present a solution to this “**Coordinate Singularity**” problem!

The **Schwarzschild** and the **Einstein “Non-Euclidean” Metrics** (ds^2) as discussed in this paper, describes the causality and geometry, of the curvature of space, space-time, and the gravitational field, and is used in conjunction, with a fluid mechanical model, **Perfect Fluid “Static or Dynamic” Vacuum Energy Solution** for the causality gravitation.

This paper postulates a “Vacuum Energy Perfect Fluid” model and a “Dark Matter Force and Pressure” associated with the Non-Euclidean Spherically Symmetric metric equations, and also gives a conceptual and mathematical description and rationale, for selecting the Schwarzschild Metric over the Einstein Metric, as a physical description of the gradient gravitational, field surrounding a localized net inertial mass/matter source.

In the next paper, I will discuss the specifics of the “Vacuum Energy Perfect Fluid” model and a “Dark Matter Force and Pressure” associated with the Non-Euclidean Spherically Symmetric metric equations.

Below are the topics that were discussed in this paper:

- **1.1 *Introduction to Basic Concepts of Euclidean and Non-Euclidean Geometry and Spherically Symmetric Metrics***
- **1.2 *General Spherically Symmetric Metric – Euclidean and Minkowski (Pseudo-Euclidean) Metrics***
- **1.3 *General Spherically Symmetric Metric – Schwarzschild Metric – Non-Euclidean Metric***
- **1.4 *General Spherically Symmetric Metric – Schwarzschild Metric – Non-Euclidean Metric – Dynamic Fluid Pressure Gradient Solution***
- **1.5 *General Spherically Symmetric Metric – Einstein Metric – Non-Euclidean Metric – Static Fluid Pressure Gradient Solution***
- **1.6 *New Mathematical Formalism – General Spherically Symmetric Metric – Einstein Metric – Non-Euclidean Metric***

References

- [1] **Physical Acceptability of Isolated, Static, Spherically Symmetric, Perfect Fluid Solutions of Einstein's Equations**, M.S.R. Delgaty and Kayll Lake¹, Department of Physics, Queen's University, Kingston Ontario, Canada K7L 3N6 (Dated: Septem 1998): <http://arxiv.org/pdf/gr-qc/9809013v1.pdf>

- [2] **Generating perfect fluid spheres in general relativity**, Petarpa Boonserm, * Matt Visser, and Silke Weinfurtner, School of Mathematics, Statistics, and Computer Science, Victoria University of Wellington, PO Box 600, Wellington, New Zealand (Dated: March 2005 - February 6, 2008): <http://arxiv.org/pdf/gr-qc/0503007v1.pdf> - (Pages: 13 - 16)

- [3] Karl Friedrich Gauss, **General Investigations of Curved Surfaces of 1827 and 1825**, The Project Gutenberg EBook of General Investigations of Curved Surfaces of 1827 and 1825, by Karl Friedrich Gauss: <http://www.gutenberg.org/files/36856/36856-pdf.pdf>

- [4] Gauss Law - Theorema Egregium: http://en.wikipedia.org/wiki/Theorema_Egregium

- [5] List of Differential Geometry Topics (Wikipedia): http://en.wikipedia.org/wiki/List_of_differential_geometry_topics

- [6] D.F. Lawden. **Introduction to Tensor Calculus, Relativity and Cosmology**; Dover Publications; 3 edition (January 27, 2003), ISBN 978-0486425405 pp. 6-10

- [7] Minkowski, Hermann (1907/8), "Die Grundgleichungen für die elektromagnetischen Vorgänge in bewegten Körpern", Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse: 53–111 *** Wikisource translation: **The Fundamental Equations for Electromagnetic Processes in Moving Bodies**. http://en.wikipedia.org/wiki/Differential_geometry_of_surfaces

- [8] List Of Differential Geometry Topics: http://en.wikipedia.org/wiki/List_of_differential_geometry_topics

- [9] Einstein's Paper: "Explanation of the Perihelion Motion of Mercury from General Relativity Theory"; Anatoli Andrei Vankov; IPPE, Obninsk, Russia; Bethany College, KS, USA:
<http://www.gsjournal.net/old/eeuro/vankov.pdf>
- [10] Schwarzschild, K. (1916). "Über das Gravitationsfeld eines Massenpunktes nach der Einstein'schen Theorie". Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften 1: 189–196: text of the original paper, in Wikisource:
[http://de.wikisource.org/wiki/%C3%9Cber_das_Gravitationsfeld_eines Massenpunktes nach der Einsteinschen Theorie](http://de.wikisource.org/wiki/%C3%9Cber_das_Gravitationsfeld_eines_Massenpunktes_nach_der_Einsteinschen_Theorie)
- [11] Schwarzschild Geodesics:
http://en.wikipedia.org/wiki/Schwarzschild_geodesics
- [12] Robert Louis Kemp. "Super Principia Mathematica – The Rage To Master Conceptual & Mathematical Physics – The General Theory of Relativity." ISBN 978-0-9841518-2-0, Flying Car Publishing Company, July 2010, pp. 20-21.
- [13] Robert Louis Kemp. "Super Principia Mathematica – The Rage To Master Conceptual & Mathematical Physics –
<http://superprincipia.wordpress.com/2011/12/23/inertial-mass-vortex-gravitation-theory-continued-part-3/>
- [14] Robert Louis Kemp. "Super Principia Mathematica – The Rage To Master Conceptual & Mathematical Physics –
<http://superprincipia.wordpress.com/2012/01/28/total-mechanical-energy-conservation-escape-velocity-binding-energy-einstein-field-equation/>
- [15] Robert Louis Kemp. "Super Principia Mathematica – The Rage To Master Conceptual & Mathematical Physics –
<http://superprincipia.wordpress.com/2012/01/16/a-theory-of-gravity-for-the-21st-century-the-gravitational-force-and-potential-energy-in-consideration-with-special-relativity-general-relativity/>

- [16] Robert Louis Kemp. "Super Principia Mathematica – The Rage To Master Conceptual & Mathematical Physics – <http://superprincipia.wordpress.com/2012/05/29/newtonian-self-gravitational-force-video-lecture/>
- [17] Robert Louis Kemp. "Super Principia Mathematica – The Rage To Master Conceptual & Mathematical Physics – <http://superprincipia.wordpress.com/2012/11/08/conditions-for-the-absence-and-presence-of-matter-in-general-relativity-gr-einstein-field-equations-efes/>