

*We can't solve problems by using the same kind of thinking we used when we created them.*

Albert Einstein

# World – Universe Model

---

## A Review

**Vladimir S. Netchitailo**  
[v.netchitailo@sbcglobal.net](mailto:v.netchitailo@sbcglobal.net)

Biolase, Inc.  
4 Cromwell, Irvine,  
CA 92618, USA

**2012 – 2013**

*The sciences do not try to explain, they hardly even try to interpret, they mainly make models. By a model is meant a mathematical construct, which, with addition of certain verbal interpretations describes observed phenomena. The justification of such a mathematical construct is solely and precisely that it is expected to work.*

John von Newmann

---

## ABSTRACT

---

World – Universe Model is based on the following primary assumptions:

- The World is finite and is expanding inside the Universe with speed equal to the electrodynamic constant  $c$ . The Universe serves as an unlimited source of energy that continuously flows into the World through the boundary.
- Medium of the World, consisting of protons, electrons, photons, neutrinos, and dark matter particles, is an active agent in all physical phenomena in the World.
- Two fundamental parameters in various rational exponents define all macro and micro features of the World: Fine-Structure Constant  $\alpha$ , and dimensionless quantity  $Q$ . While  $\alpha$  is constant,  $Q$  increases with time, and is in fact a measure of the size and the age of the World.

The World – Universe Model explains experimental data accumulated in the field of Cosmology over the last decades: the size and age of the World; critical energy density and the gravitational parameter; temperatures of the cosmic microwave background radiation and black body radiation of cosmic dust; fractal structure of the World and its evolution; World expansion and cosmological redshift. Additionally, the Model makes predictions pertaining to masses of dark matter particles, photons, axions, and neutrinos; proposes new types of particle interactions and the fundamental physical parameters of the World; and resolves paradoxes like “Matter – Antimatter Asymmetry” and “Faint Young Sun”.

**Keywords:** Fractal Cosmology; World – Universe Model; Fine-Structure Constant; Dark Matter; Microwave Background Radiation; Maxwell-Lorentz Equations; Dirac’s Equations; Fractal Particle Structure; Grand Unified Theory

## CONTENTS

---

Abstract	1
1. Instead of an Introduction	3
2. Cosmology	4
2.1. The Beginning	4
2.2. Time Varying Parameters	5
2.3. The Size and the Age of the World	5
2.4. Critical Energy Density, Gravitational Parameter, Front Surface Enthalpy	7
2.5. Microwave Background Radiation	8
2.6. Mass Varying Photons, Speed of Light	11
2.7. Mass Varying Neutrinos, Distribution of the World's Energy Density	12
2.8. Fine-Structure Constant	17
2.9. Dark Matter Particles	19
2.10. Macroobject Cores built up from Fermionic Dark Matter	22
2.11. Fractal Cosmology	27
2.12. Fractal structure of the World	31
2.13. Evolution of the World's fractal structure	34
2.14. Extrasolar Systems	37
2.15. Brown Dwarfs	42
2.16. Planets	43
2.17. Cosmic Dust	45
2.18. World Expansion	46
2.19. Cosmological Redshift	49
3. Particle Physics	51
3.1. Analysis of Maxwell's Equations	51
3.2. Magnetic Monopole, Magnetic Dipole	54
3.3. Dirac's Equation for Gravitoelectromagnetic Field	56
3.4. Elementary Charge	58
3.5. Model of an Electron	61
3.6. Fractal Structure of Particles	63
3.7. Grand Unified Theory	65
4. The World	70

## 1. INSTEAD OF AN INTRODUCTION

---

*In science one tries to tell people, in such a way as to be understood by everyone, something that no one knew before.*

Paul Dirac

I am a Doctor of Sciences in Laser Physics, specializing in interaction of laser radiation with matter. I belong to the school of physicists established by Alexander Prokhorov – Nobel Prize Laureate in Physics and co-inventor of masers and lasers. Most of my past papers were published under V.S. Nechitailo alias.

I have been developing the World – Universe Model for over 10 years, and am now presenting it for the first time in this Review.

A number of ideas presented in this paper are not new, and I don't claim credit for them. In fact, several ideas belonging to classical physicists such as Tesla, Dirac, McCullagh, and Bjerknes, are examined in a new light. The World – Universe Model unifies and simplifies existing theories and results in Cosmology into a single coherent picture.

World – Universe Model, at its present state, requires significant further elaboration and validation. I welcome criticism of the overall Model and individual ideas underpinning it.

The Model is developed around two fundamental parameters: Fine-Structure Constant  $\alpha$ , and dimensionless quantity  $Q$ . While  $\alpha$  is a constant,  $Q$  increases with time, and in fact defines the size and the age of the World.

Intermediate results in subsequent analysis will often be obtained using classical notions and parameters. All final formulas, however, will be expressed in terms of  $\alpha$  and  $Q$  in various rational exponents, as well as small integer numbers and  $\pi$ .

Numerical values are provided in SI for convenience.  $\alpha$ -dependent quantities are calculated to 8 significant digits, and  $Q$ -dependent quantities – to 5 significant digits.

We will use  $\Leftrightarrow$  symbol to describe ranges of values.

## 2. COSMOLOGY

---

### 2.1. THE BEGINNING

---

*The Big Bang is a mythical “creation event”, before that there was nothing. Can we really believe in nothing turning to something out of the blue? Where did all that energy for the rapid expansion of the Universe and the forming of Galaxies and such come from?*

O.A. van den Berg

In the beginning, there was nothing but the Universe – unlimited field of energy. Our World was started by a fluctuation in the Universe, and the Core of the World was born. The World has since been expanding through the Universe, consuming energy as the World – Universe boundary advances.

The World is expanding with speed equal to electrodynamic constant  $c$  for time  $t$ , and has the radius of  $R = ct$ . Subsequently, we will refer to the moving World – Universe boundary as the Front.

The Front has a temperature invariant surface enthalpy  $\sigma_0$ . Generation of particle – antiparticle pairs is occurring at the Front due to high surface energy density of the Universe. Antiparticles remain at the Front, and particles continue on into the World. In other words, all antimatter is concentrated at the Front, and equal amount of matter exists in the World, resolving the long-standing “Matter – Antimatter Asymmetry” paradox.

Amount of energy added to the World is proportional to the increase of the area of the Front. The total amount of the World energy is thus

$$E_W = 4\pi R^2 \sigma_0 \tag{2.1.1}$$

The energy density of the World  $\rho_W$  is inversely proportional to the radius of the World  $R$ :

$$\rho_W = \frac{3\sigma_0}{R} \tag{2.1.2}$$

---

## 2.2. TIME VARYING PARAMETERS

---

Recall the well-known equation for the critical energy density of the World  $\rho_{cr}$ :

$$\rho_{cr} = \frac{3H_0^2 c^2}{8\pi G} \quad 2.2.1$$

where  $H_0$  is the Hubble parameter:

$$H_0 = \frac{1}{t} = \frac{c}{R} \quad 2.2.2$$

Equation 2.2.1 can be rewritten as

$$\frac{4\pi G}{c^2} \times \frac{2}{3} \rho_{cr} = H_0^2 = \frac{c^2}{R^2} \quad 2.2.3$$

The principal idea of our Model is that the energy density of the World  $\rho_W$  equals to the critical energy density  $\rho_{cr}$ :

$$\rho_{cr} = \rho_W = \frac{3\sigma_0}{R} \propto \frac{1}{R} \quad 2.2.4$$

We see that the gravitational parameter  $G$  is also proportional to  $\frac{1}{R}$  and is decreasing in time as  $G \propto \frac{1}{t}$ . This property of gravitational parameter  $G$  was originally hypothesized by Paul Dirac in 1937.<sup>1</sup>

Since  $M_p^2 = \frac{hc}{2\pi G}$ , the Planck mass  $M_p$  is proportional to  $t^{\frac{1}{2}}$ , and the Planck length  $L_p = \frac{h}{M_p c}$  is proportional to  $t^{-\frac{1}{2}}$ , where  $h$  is the Planck constant.

---

## 2.3. THE SIZE AND THE AGE OF THE WORLD

---

Let's introduce a dimensionless time-varying quantity  $Q$ :

$$Q = \frac{R}{2\pi a_0} = \frac{R}{a} \quad 2.3.1$$

where  $a_0$  is the classical electron radius and  $a$  is the radius of the World's Core at the Beginning ( $Q=1$ ):

$$a = 2\pi a_0 = 1.7705645 \times 10^{-14} \text{ m} \quad 2.3.2$$

Let us additionally introduce a basic unit of time  $t_0$ :

$$t_0 = \frac{a}{c} = 5.9059674 \times 10^{-23} \text{ s} \quad 2.3.3$$

---

<sup>1</sup> P. A. M. Dirac, Nature **139**, 323 (1937).

We will subsequently use  $a$  as the basic unit of measure of length and  $t_0$  – as the unit of time. Quantity  $Q$  is then the radius of the World measured in terms of  $a$ .

Let's introduce a length parameter  $L_g$  that is the geometric mean of the World's current radius  $R$  and its Core radius  $a$ :

$$L_g = \sqrt{aR} \quad 2.3.4$$

In our Model,  $L_g$  is a basic unit of measure of macroobjects size (see Section 2.10), and it satisfies the following equation:

$$2L_g L_p = a^2 \quad 2.3.5$$

The radius of the World  $R$  is then

$$R = \frac{a^3}{4L_p^2} = a \times Q \quad 2.3.6$$

Substituting values of  $a$  and  $L_p$  into 2.3.6 we obtain:

$$R = 1.3459 \times 10^{26} \text{ m} \quad 2.3.7$$

We can now calculate the age of the World  $A_t$  at current time  $t$ :

$$\begin{aligned} A_t &= \frac{R}{c} = 4.4894 \times 10^{17} \text{ s} = \\ &= 14.226 \text{ billion years (By)} \end{aligned} \quad 2.3.8$$

The age of the World calculated based on the gravitational parameter  $G$  (14.226 By) is somewhat greater than the commonly adopted value of 13.772 By.

Calculating the value of Hubble parameter  $H_0$  based on  $A_t$ , we obtain

$$H_0 = \frac{1}{A_t} = 2.2275 \times 10^{-18} \text{ s}^{-1} = 68.733 \frac{\text{km/s}}{\text{Mpc}} \quad 2.3.9$$

which is in good agreement with  $H_0 = 69.32 \pm 0.8 \frac{\text{km/s}}{\text{Mpc}}$  obtained using WMAP data [Wikipedia, Hubble's Law].

From 2.3.6 we calculate the value of the dimensionless parameter  $Q$ :

$$Q = 0.76014 \times 10^{40} \quad 2.3.10$$

Parameter  $Q$  defines both the size and the age of the World: radius of the World  $R = a \times Q$ , and age of the World  $A_t = \frac{R}{c} = t_0 \times Q$ .

## 2.4. CRITICAL ENERGY DENSITY, GRAVITATIONAL PARAMETER, FRONT SURFACE ENTHALPY

---

The gravitational parameter  $G$  equals to

$$\begin{aligned} G &= \frac{L_p^2 c^4}{2\pi h c} = \frac{a^2 c^4}{8\pi h c} \times Q^{-1} = \\ &= 6.6726 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2} \end{aligned} \quad 2.4.1$$

Using equation 2.2.3 we calculate the value of critical density  $\rho_{cr}$ :

$$\begin{aligned} \rho_{cr} &= \frac{3hc}{a^4} \times Q^{-1} = \rho_{cr0} \times Q^{-1} = \\ &= 7.9773 \times 10^{-10} \frac{\text{J}}{\text{m}^3} \end{aligned} \quad 2.4.2$$

$\rho_{cr0}$  is the energy density of the World at the Beginning:

$$\rho_{cr0} = 3\rho_0 = 6.0638901 \times 10^{30} \frac{\text{J}}{\text{m}^3} \quad 2.4.3$$

$\rho_0$  is the basic unit of energy density:

$$\rho_0 = \frac{hc}{a^4} = 2.0212967 \times 10^{30} \frac{\text{J}}{\text{m}^3} \quad 2.4.4$$

The gravitomagnetic parameter of the World's Medium  $\mu_g$  is

$$\mu_g = \frac{4\pi G}{c^2} = \frac{a^2 c^2}{2hc} \times Q^{-1} = (2\rho_0 t_0^2)^{-1} \times Q^{-1} \quad 2.4.5$$

and the gravitoelectric parameter of the Medium  $\varepsilon_g$  is

$$\varepsilon_g = \frac{1}{\mu_g c^2} \quad 2.4.6$$

The surface enthalpy of the World – Universe Front is

$$\sigma_0 = \frac{hc}{a^3} = (\rho_0^2 E_0)^{\frac{1}{3}} = 3.5788363 \times 10^{16} \frac{\text{J}}{\text{m}^2} \quad 2.4.7$$

The total energy of the World  $E_W$  at current time  $t$  then equals to

$$\begin{aligned} E_W &= \frac{4\pi R^2 hc}{a^3} = 4\pi E_0 \times Q^2 = 4\pi E_0 \left(\frac{A_t}{t_0}\right)^2 = \\ &= 8.1464 \times 10^{69} \text{ J} \end{aligned} \quad 2.4.8$$

where basic energy unit  $E_0$  equals to

$$E_0 = \frac{hc}{a} = 1.1219288 \times 10^{-11} \text{ J} = 70.025267 \text{ MeV} \quad 2.4.9$$



The proportionality of total energy in the World to its age squared ( $E_W \propto A_t^2$ ) was also hypothesized by Paul Dirac. <sup>1</sup>

In our Model, Length, Time, Energy, and Energy Density are measured in units of basic length  $a$ , time  $t_0$ , energy  $E_0$ , and energy density  $\rho_0$  respectively. All other physical characteristics are calculated in terms of these basic units.

Plugging the values of  $G$  and  $E_W$  into the formula of Schwarzschild radius,

$$R_{SH} = \frac{2GE_W}{c^4} = \frac{2}{c^4} \frac{a^2 c^4}{8\pi h c} \times Q^{-1} \times 4\pi \frac{h c}{a} \times Q^2 = R \quad 2.4.10$$

we conclude that the World is a black hole.

The hypothesis *that the universe may not only be a closed structure (as perceived by its inhabitants at the present epoch) but may also be a black hole, confined to a localized region of space which cannot expand without limit* was proposed by Raj Pathria in 1972. <sup>2</sup> In our Model, the World expands in the Universe without limit, because the Universe is an unlimited source of energy.

Residing inside of a black hole, we can conduct no observations of the outside Universe, and learn nothing about its characteristics. We can only observe and measure the Universe's interaction with the World that occurs at the World - Universe Front: the temperature invariant surface enthalpy  $\sigma_0$ .

---

## 2.5. MICROWAVE BACKGROUND RADIATION

---

In our Model, the World consists of stable elementary particles with lifetimes longer than the age of the World. Protons with mass  $m_p$  and energy  $E_p = m_p c^2$  and electrons with mass  $m_e$  and energy  $E_e = \alpha E_0$  have identical concentrations in the World:  $n_p = n_e$ , where  $E_0$  is the basic energy and  $\alpha$  is the fine-structure constant.

Low density plasma consisting of protons and electrons has plasma frequency  $\omega_{pl}$ :

$$\omega_{pl}^2 = \frac{4\pi n_e e^2}{4\pi \epsilon_0 m_e} = 4\pi n_e \alpha \frac{h}{2\pi m_e c} c^2 = 2n_e \alpha c^2 \quad 2.5.1$$

where  $e$  is the elementary charge and  $\epsilon_0$  is the permittivity of the Medium.

Let's choose  $\omega_{pl}$  that satisfies the following equation:

$$\omega_{pl} = \frac{m_e 2\pi c}{m_p L_g} = \frac{m_e}{m_p} t_0^{-1} \times Q^{-\frac{1}{2}} \quad 2.5.2$$

---

<sup>2</sup> R. K. Pathria, Nature, **240**, 298 (1972).

$\omega_{pl}^2$  is then proportional to  $Q^{-1}$ . Energy densities of protons and electrons are then proportional to  $\frac{1}{R}$ , similar to the critical energy density  $\rho_{cr} \propto \frac{1}{R}$  (2.2.4).

Since the formula calculating the potential energy of interaction of protons and electrons contain the same parameter  $k_{pe}$ :

$$k_{pe} = m_p \omega_{pl}^2 = m_e \left( \frac{2\pi c}{L_g} \right)^2 \quad 2.5.3$$

we substitute  $\omega_{pl}^2 = \frac{m_e}{m_p} \left( \frac{2\pi c}{L_g} \right)^2$  into 2.5.1 and calculate concentrations of protons and electrons:

$$n_p = n_e = \frac{2\pi^2 m_e}{a^3 m_p} \times Q^{-1} \quad 2.5.4$$

$\rho_p = n_p E_p$  is the energy density of protons in the Medium. The relative energy density of protons  $\Omega_p$  is then the ratio of  $\frac{\rho_p}{\rho_{cr}}$ :

$$\Omega_p = \frac{\rho_p}{\rho_{cr}} = \frac{2\pi^2 \alpha}{3} = 0.048014655 \quad 2.5.5$$

The above value is in good agreement with estimations of baryonic matter in the World  $\Omega_p \cong 0.046$  [Wikipedia, Dark Matter].

From equations 2.5.1 and 2.5.4 we obtain the value of the lowest radio-wave frequency  $\nu_{pl}$ :

$$\nu_{pl} = \frac{\omega_{pl}}{2\pi} = \left( \frac{m_e}{m_p} \right)^{\frac{1}{2}} t_0^{-1} \times Q^{-\frac{1}{2}} = 4.5322 \text{ Hz} \quad 2.5.6$$

Note that this value is close to the low limit of the standard Extremely low frequency band  $3 \Leftrightarrow 30 \text{ Hz}$  [Wikipedia, Radio Spectrum].

Substituting radius of the World  $R$  obtained in 2.3.6, we use equation 2.5.4 to calculate the proton and electron concentrations in the Medium:

$$n_p = n_e = 8\pi^2 \frac{m_e L_p^2}{m_p a^5} = 0.25480 \text{ m}^{-3} \quad 2.5.7$$

which is in good agreement with their estimated concentration in the intergalactic medium  $n_p \cong 0.25 \text{ m}^{-3}$  [Wikipedia, Outer space].

$\rho_e = n_e E_e$  is the energy density of electrons in the Medium. The relative energy density of electrons  $\Omega_e$  is then the ratio of  $\frac{\rho_e}{\rho_{cr}}$ :

$$\Omega_e = \frac{\rho_e}{\rho_{cr}} = \frac{2\pi^2\alpha m_e}{3 m_p} \quad 2.5.8$$

Let's assume that the energy density of Microwave Background Radiation  $\rho_{MBR}$  is twice larger than  $\rho_e$  (due to two polarizations of photons):

$$\rho_{MBR} = 4\pi^2\alpha \frac{m_e}{m_p} \rho_0 \times Q^{-1} = \frac{8\pi^5}{15} \frac{k_B^4}{(hc)^3} T_{MBR}^4 \quad 2.5.9$$

where  $k_B$  is the Boltzmann constant and  $T_{MBR}$  is MBR temperature.

We can now calculate the value of  $T_{MBR}$ :

$$T_{MBR} = \frac{E_0}{k_B} \left( \frac{15\alpha m_e}{2\pi^3 m_p} \right)^{\frac{1}{4}} \times Q^{-\frac{1}{4}} = 2.7250 K \quad 2.5.10$$

Thus calculated value of  $T_{MBR}$  is in excellent agreement with experimentally measured value of  $2.72548 \pm 0.00057 K$  [Wikipedia, Cosmic microwave background radiation].

At the Beginning of the World, the extrapolated value of  $T_{MBR0}$  at  $Q = 1$  is

$$T_{MBR0} = 2.1927 \text{ MeV} = 2.5445 \times 10^{10} K \quad 2.5.11$$

Note that  $T_{MBR0}$  is significantly smaller than values commonly discussed in literature.

Let's proceed to calculate the value of  $T_{MBR}$  at different Ages of the World  $A_t$ :

Age	$T_{MBR}$
1 s	6.0785 eV = 70,537 K
$10^8 \text{ s} \cong 3.2 \text{ y}$	705.37 K
$10^{16} \text{ s} \cong 0.32 \text{ By}$	7.0537 K
$3 \times 10^{17} \text{ s} \cong 9.6 \text{ By}$ (birth of Solar system)	3.0140 K
$4.49 \times 10^{17} \text{ s} \cong 14.23 \text{ By}$ (present)	2.7250 K

Observe that practically all macroobjects – galaxies, stars, planets, etc. – have arisen in a cold World. Our Solar system, for instance, was created when the temperature of MBR was about 3 K. Therefore, any Model describing creation of macroobjects must hold true in cold World conditions.

## 2.6. MASS VARYING PHOTONS, SPEED OF LIGHT

---

Photons with energy smaller than  $E_{ph} = hv_{pl}$  cannot propagate in plasma, thus  $hv_{pl}$  is the smallest amount of energy a photon may possess. This amount of energy can be viewed as a particle (we'll name it axion), whose frequency-independent effective "rest mass" equals to

$$\begin{aligned} m_a &= E_a/c^2 = \left(\frac{m_e}{m_p}\right)^{\frac{1}{2}} m_0 \times Q^{-\frac{1}{2}} = 3.6680 \times 10^{-20} m_e = \\ &= 1.8743 \times 10^{-14} \text{ eV}/c^2 \end{aligned} \quad 2.6.1$$

where  $E_a$  is a rest energy of the axion and  $m_0$  is a basic unit of mass that equals to

$$m_0 = \frac{E_0}{c^2} = 70.025267 \text{ MeV}/c^2 = 1.2483143 \times 10^{-28} \text{ kg} \quad 2.6.2$$

The calculated mass of an axion is in agreement with  $m_a \sim 10^{-15} \text{ eV}/c^2$  discussed by C. Csaki *et al*<sup>3</sup>, and with experimental checks of Coulomb's law on photon mass  $m_{ph}$ . *A null result of such an experiment has set a limit of  $m_{ph} \lesssim 10^{-14} \text{ eV}/c^2$ . If the photon mass is generated via the Higgs mechanism then the upper limit of  $m_{ph} \lesssim 10^{-14} \text{ eV}/c^2$  from the test of Coulomb's law is valid [Wikipedia, Photon].*

Energy of an axion moving with a group velocity  $v_{gr}$  is given by

$$E_a(v_{gr}) = \frac{hv_{pl}}{\sqrt{1 - \frac{v_{gr}^2}{c^2}}} \quad 2.6.3$$

Taking into account the dispersion relation for plasma:

$$v_{gr} v_{ph} = c^2 \quad 2.6.4$$

and the value of phase velocity  $v_{ph} = \frac{c}{n_{pl}}$ , where  $n_{pl}$  is the index of plasma refraction,

$$n_{pl} = \sqrt{1 - \frac{v_{pl}^2}{v^2}} \quad 2.6.5$$

from equation 2.6.4 it follows that

$$\frac{v_{gr}^2}{c^2} = 1 - \frac{v_{pl}^2}{v^2} \quad 2.6.6$$

and we calculate moving axion energy  $E_a(v_{gr})$  to be

$$E_a(v_{gr}) = hv = E_{ph} \quad 2.6.7$$

---

<sup>3</sup> C. Csaki, N. Kaloper, and J. Terning (2001), arXiv: hep-ph/0112212.

where  $\nu$  is photon frequency.

In our Model, the total energy of a moving particle consists of two components: rest energy and “coat” energy. A particle’s coat is the response of the Medium to the particle’s movement. A photon is then a constituent axion with rest energy  $E_a = h\nu_{pl}$  and total energy  $E_{ph} = h\nu$ . In most cases  $\nu \gg \nu_{pl}$ , and practically all of the photon’s energy is concentrated in the axion’s coat that is the part of the Medium surrounding the axion. Axions are fully characterized by their four-momentum.

Rest energy of the axion is decreasing with time:  $E_a \propto t^{-\frac{1}{2}}$  (see 2.6.1), and total energy remains constant (2.6.7).

The higher the photon’s energy, the closer its speed approaches  $c$ . But the fact that axions possess non-zero rest masses means that photons can never reach that speed.

## 2.7. MASS VARYING NEUTRINOS, DISTRIBUTION OF THE WORLD’S ENERGY DENSITY

---

It is now established that there are at least three different types of neutrinos: electronic  $\nu_e$ , muonic  $\nu_\mu$ , and tauonic  $\nu_\tau$ , and their antiparticles. Pontecorvo and Smorodinskii discussed the possibility of energy density of neutrinos exceeding that of baryonic matter.<sup>4</sup> Neutrino oscillations imply that neutrinos have non-zero masses.

Let’s take neutrinos masses  $m_{\nu_e}$ ,  $m_{\nu_\mu}$ ,  $m_{\nu_\tau}$  that are near

$$m_\nu = m_0 \times Q^{-\frac{1}{4}} \cong 7.5 \times 10^{-3} \text{ eV}/c^2 \quad 2.7.1$$

Their concentrations  $n_\nu$  are then proportional to

$$n_\nu \propto \frac{1}{a^3} \times Q^{-\frac{3}{4}} = \frac{1}{L_F^3} \quad 2.7.2$$

where Fermi length parameter  $L_F$

$$L_F = a \times Q^{\frac{1}{4}} \quad 2.7.3$$

is a characteristic of neutrino density (2.7.2), and also of critical energy density:

$$\rho_{cr} = \frac{3hc}{L_F^4} \quad 2.7.4$$

---

<sup>4</sup> B. Pontecorvo and Ya. Smorodinskii, Sov. Phys. JETP, **14**, 173 (1962).

Energy densities of neutrinos are proportional to  $Q^{-1}$ , and consequently to  $\frac{1}{R}$ , since critical energy density  $\rho_{cr}$  is proportional to  $\frac{1}{R}$ .

Experimental results obtained by M. Sanchez<sup>5</sup> show  $\nu_e \rightarrow \nu_{\mu,\tau}$  neutrino oscillations with parameters given by

$$2.3 \times 10^{-5} eV^2/c^4 \leq \Delta m_{sol}^2 \leq 9.3 \times 10^{-5} eV^2/c^4 \quad 2.7.5$$

and  $\nu_{\mu} \rightarrow \nu_{\tau}$  neutrino oscillations with parameters

$$1.6 \times 10^{-3} eV^2/c^4 \leq \Delta m_{atm}^2 \leq 3.9 \times 10^{-3} eV^2/c^4 \quad 2.7.6$$

where  $\Delta m_{sol}^2$  and  $\Delta m_{atm}^2$  are mass splitting for solar and atmospheric neutrinos respectively.

Significantly more accurate results were obtained by P. Kaus *et al*<sup>6</sup> for the following ratio:

$$\sqrt{\frac{\Delta m_{sol}^2}{\Delta m_{atm}^2}} \cong 0.16 \quad 2.7.7$$

Let's assume that muonic neutrino's mass indeed equals to

$$m_{\nu_{\mu}} = m_{\nu} = m_0 \times Q^{-\frac{1}{4}} \cong 7.5 \times 10^{-3} eV/c^2 \quad 2.7.8$$

From equation 2.7.7 it then follows that

$$m_{\nu_{\tau}} = 6m_{\nu} \cong 4.5 \times 10^{-2} eV/c^2 \quad 2.7.9$$

Based on equation for Fermi Coupling Parameter  $G_F$  (see Section 3.7),

$$m_{\nu_e} = \frac{1}{24} m_{\nu} \cong 3.1 \times 10^{-4} eV/c^2 \quad 2.7.10$$

Then the squared values of the muonic and tauonic masses fall into ranges 2.7.5 and 2.7.6:

$$\begin{aligned} m_{\nu_{\mu}}^2 &\cong 5.6 \times 10^{-5} eV^2/c^4 \\ m_{\nu_{\tau}}^2 &\cong 2 \times 10^{-3} eV^2/c^4 \end{aligned} \quad 2.7.11$$

Considering that all elementary particles, including neutrinos, are fully characterized by their four-momentum  $(\frac{E_{\nu i}}{c}, \mathbf{p}_{\nu i})$ :

$$\left(\frac{E_{\nu i}}{c}\right)^2 - \mathbf{p}_{\nu i}^2 = (m_{\nu i}c)^2$$

---

<sup>5</sup> M. Sanchez, Oscillation Analysis of Atmospheric Neutrinos in Soudan 2, Dissertation, 2003.

<sup>6</sup> P. Kaus, S. Meshkov, Neutrino Mass Matrix and Hierarchy, AIP Conf. Proc. **672**, 117 (2003).

$$i = e, \mu, \tau \quad 2.7.12$$

we obtain the following neutrino energy densities in accordance with theoretical calculations made by L. D. Landau and E. M. Lifshitz <sup>7</sup>:

$$\begin{aligned} \rho_{\nu i} &= \frac{8\pi c}{h^3} \int_0^{p_F} p^2 \sqrt{p^2 + m_{\nu i}^2 c^2} dp = \\ &= \frac{2\pi(p_F c)^4}{(hc)^3} \times F(x_{\nu i}) \end{aligned} \quad 2.7.13$$

where  $p_F$  is Fermi momentum,

$$F(x_{\nu i}) = \frac{x_{\nu i}^{\frac{1}{2}}(2x_{\nu i}+1)(x_{\nu i}+\frac{1}{2})^{\frac{1}{2}} - \ln \left[ x_{\nu i}^{\frac{1}{2}} + (x_{\nu i}+1)^{\frac{1}{2}} \right]}{2x_{\nu i}^2} \quad 2.7.14$$

$$x_{\nu i} = \left( \frac{p_F}{m_{\nu i} c} \right)^2 \quad 2.7.15$$

$$m_{\nu i} = A_i m_0 \times Q^{-\frac{1}{4}} \quad 2.7.16$$

$$A_i = \frac{1}{24}; 1; 6 \quad 2.7.17$$

Let's take the following value for Fermi momentum  $p_F$ :

$$p_F^2 = \frac{h^2}{2\pi^2 L_F^2} = \frac{h^2}{2\pi^2 a^2} \times Q^{-\frac{1}{2}} = p_0^2 \times Q^{-\frac{1}{2}} \quad 2.7.18$$

where  $p_0^2 = \frac{h^2}{2\pi^2 a^2}$  is the extrapolated value of  $p_F$  at the Beginning when  $Q = 1$ . As a side note, the equation for surface enthalpy of the World – Universe Front  $\sigma_0$  can be rewritten as  $\sigma_0 a_0^2 = \frac{p_0^2}{2m_0}$ .

Using 2.7.13, we obtain neutrino relative energy densities  $\Omega_{\nu i}$  in the Medium in terms of the critical energy density  $\rho_{cr}$ :

$$\Omega_{\nu i} = \frac{\rho_{\nu i}}{\rho_{cr}} = \frac{1}{6\pi^3} F(y_{\nu i}) \quad 2.7.19$$

where

$$y_{\nu i} = (2\pi^2 A_i^2)^{-1} \quad 2.7.20$$

---

<sup>7</sup> L. D. Landau, E. M. Lifshitz, Statistical Physics, Third Edition, Part 1: Volume 5 (Course of Theoretical Physics, Volume 5).

It's commonly accepted that concentrations of all types of neutrinos are equal. This assumption allows us to calculate the total neutrino relative energy density in the Medium:

$$\Omega_\nu = \frac{\rho_\nu}{\rho_{cr}} = \frac{\rho_{\nu e} + \rho_{\nu \mu} + \rho_{\nu \tau}}{\rho_{cr}} = 0.45801647 \quad 2.7.21$$

One of the principal ideas of World – Universe Model holds that energy densities of Medium particles are proportional to proton energy density in the World's Medium (2.5.5):

$$\Omega_p = \frac{2\pi^2\alpha}{3} = 0.048014655 \quad 2.7.22$$

Let's take  $\Omega_\nu = \frac{30}{\pi}\Omega_p$ . We obtain

$$\Omega_\nu = 20\pi\alpha = 0.45850618 \quad 2.7.23$$

which is close to the value calculated in 2.7.21 (the difference is  $\cong 0.1\%$ ). The slight increase of neutrinos energy density can be attributed to the additional temperature-dependent part of neutrinos energy density at the Medium temperature  $T_M > 0$ :  $T_M = T_{MBR} = 2.7250 \text{ K}$ .

The electron relative energy density in the Medium  $\Omega_e$  is

$$\Omega_e = \frac{m_e}{m_p}\Omega_p \quad 2.7.24$$

The sum of MBR photons, black body radiation from cosmic dust, X-rays, and Gamma rays relative energy densities (see Section 2.9) is

$$\Omega_{rad} = \left(\frac{8}{3} + \frac{2}{15\pi}\right)\Omega_e \quad 2.7.25$$

Dark Matter (DM) energy density in the Medium (see Section 2.9) is

$$\Omega_{DM} = \frac{10}{3}\Omega_p \quad 2.7.26$$

The total Medium relative energy density  $\Omega_M$  then equals to

$$\begin{aligned} \Omega_M &= \Omega_p + \Omega_e + \Omega_{rad} + \Omega_\nu + \Omega_{DM} = \\ &= \left[\frac{13}{3} + \left(\frac{11}{3} + \frac{2}{15\pi}\right)\frac{m_e}{m_p} + \frac{30}{\pi}\right]\Omega_p = \frac{2}{3} \end{aligned} \quad 2.7.27$$

Let's recall that equation 2.2.3 contains the gravitomagnetic parameter of the World's Medium  $\mu_g = \frac{4\pi G}{c^2}$  multiplied by  $\frac{2}{3}\rho_{cr}$ . It follows that the World's Medium has energy density  $\rho_M = \frac{2}{3}\rho_{cr}$ , and the remaining energy  $E_{MO} = \frac{1}{3}\rho_{cr}V_W$  resides in the World's macroobjects (galaxies, stars, planets, cosmic dust, etc.). The World relative energy density  $\Omega_W$  is then



$$\Omega_W = 1.5\Omega_M = 1 \quad 2.7.28$$

The obtained result means that the calculated energy density of the World  $\rho_W$  equals to the critical energy density  $\rho_{cr}$  that is in accordance with the principal idea of our Model (see Section 2.2).

The total neutrinos energy density (in the Medium and in macroobjects) equals to

$$\Omega_{vtot} = \frac{45}{\pi}\Omega_p = 0.68775927 \quad 2.7.29$$

The total Dark Matter energy density is

$$\Omega_{DMtot} = 5\Omega_p = 0.24007327 \quad 2.7.30$$

The total baryonic energy density  $\Omega_B$  is

$$\Omega_B = 1.5\Omega_p = 0.072021982 \quad 2.7.31$$

To summarize:

- The World's energy density is proportional to  $Q^{-1}$ ;
- The particles relative energy densities are proportional to  $\alpha$ ;
- The total neutrinos energy density is almost 10 times greater than baryonic energy density, and about 3 times greater than Dark Matter energy density.

## 2.8. FINE-STRUCTURE CONSTANT

---

*The mystery about  $\alpha$  is actually a double mystery. The first mystery - the origin of its numerical value  $\approx 1/137$  has been recognized and discussed for decades. The second mystery – the range of its domain – is generally unrecognized.*

Malcolm H. Mac Gregor

The Fine-structure constant (FSC)  $\alpha$  is a fundamental physical constant. Wikipedia has this to say about the FSC [Fine-structure constant]:

*Arnold Sommerfeld introduced the Fine-Structure Constant in 1916, as part of his theory of the relativistic deviations of atomic spectral lines from the predictions of the Bohr model. The first physical interpretation of the fine-structure constant  $\alpha$  was as the ratio of the velocity of the electron in the first circular orbit of the relativistic Bohr atom to the speed of light in vacuum. Equivalently, it was the quotient between the maximum angular momentum allowed by relativity for a closed orbit, and the minimum angular momentum allowed for it by the quantum mechanics. It appears naturally in Sommerfeld's analysis, and determines the size of the splitting or fine-structure of the hydrogenic spectral lines.*

*The fine-structure constant  $\alpha$  has several physical interpretations.  $\alpha$  is:*

- *The square of  $\alpha$  is the ratio between the Hartree energy (27.2 eV = twice the Rydberg energy) and the electron rest mass (511 keV);*
- *The ratio of three characteristic lengths: the classical electron radius  $a_0$ , the Bohr radius  $a_B$  and the Compton wavelength of electron  $L_{ce}$  over  $2\pi$ :  $a_0 = \frac{\alpha L_{ce}}{2\pi} = \alpha^2 a_B$ ;*
- *The ratio of the electromagnetic impedance of the free space  $\frac{1}{\epsilon_0 c} \cong 377 \Omega$ , to the quantum of Resistance,  $\frac{h}{e^2} \approx 25.8 \text{ k}\Omega$  is  $2\alpha$ , etc.*

The Fine-structure constant  $\alpha$  plays a central role in the World – Universe Model.

Recall that by definition, the classical radius of an electron  $a_0$  is

$$a_0 = \frac{e^2}{4\pi\epsilon_0 m_e c^2} \tag{2.8.1}$$

Using the following equation:

$$\frac{e^2}{4\pi\epsilon_0} = \frac{2\pi a_0}{L_{ce}} \frac{hc}{2\pi} = \alpha \frac{hc}{2\pi} \tag{2.8.2}$$

we can conclude that  $\alpha$  is really the ratio of the classical electron radius to the electron Compton length over  $2\pi$ .

Equivalently,  $\alpha$  is the rest mass of an electron  $m_e$  measured in terms of basic units  $m_0$

$$m_0 = \frac{h}{ac} = 70.025267 \text{ MeV}/c^2 \quad 2.8.3$$

which is related to the basic energy unit  $E_0$ :

$$E_0 = m_0 c^2 \quad 2.8.4$$

Masses of all stable elementary particles of the World can be expressed in terms of  $m_0$  as follows:

$$m_e = \alpha m_0$$

$$m_p = \beta m_0$$

$$m_a = \left(\frac{\alpha}{\beta}\right)^{\frac{1}{2}} m_0 \times Q^{-\frac{1}{2}}$$

$$m_\nu = m_0 \times Q^{-\frac{1}{4}} \quad 2.8.5$$

$\beta = 13.399053$  is the ratio of proton mass  $m_p$  to the basic mass  $m_0$ . The ratio of the electron mass to the proton mass can then be expressed as follows:

$$\frac{m_e}{m_p} = \frac{\alpha}{\beta} \quad 2.8.6$$

Additionally,  $m_0$  plays a key role when masses of Dark Matter particles are discussed in Section 2.9.

## 2.9. DARK MATTER PARTICLES

---

Dark Matter (DM) is among the most important open problems in both cosmology and particle physics.

There are three prominent hypotheses on nonbaryonic DM, namely Hot Dark Matter (HDM), Warm Dark Matter (WDM), and Cold Dark Matter (CDM).

The most widely discussed models for nonbaryonic DM are based on the CDM hypothesis, and corresponding particles are most commonly assumed to be Weakly Interacting Massive Particles (WIMPs).

A neutralino with mass  $m_N$  in  $100 \Leftrightarrow 10,000 \text{ GeV}/c^2$  range is the leading DM candidate [Wikipedia, Neutralino]. Light Dark Matter Particles that are heavier than WDM and HDM but lighter than the traditional forms of CDM (neutralino) are DM candidates too. Their masses  $m_{WIMP}$  fall into  $1 \Leftrightarrow 10 \text{ GeV}/c^2$  range [Wikipedia, Light dark matter]. Subsequently, we will refer to the light dark matter particles as WIMPs.

It is known that a sterile neutrino with mass  $m_{\nu_s}$  in  $1 \Leftrightarrow 10 \text{ keV}/c^2$  range is a good WDM candidate [Wikipedia, Warm dark matter]. The best candidate for the identity of HDM is neutrino [Wikipedia, Hot dark matter]. In our opinion, a tauonic neutrino is a good HDM candidate.

In addition to fermions discussed above, we offer another type of Dark Matter particles – bosons, consisting of two fermions each. There are two types of Dark Matter bosons: DIRACs possessing mass of  $m_0$  that are in fact magnetic dipoles, and ELOPs having mass of  $\frac{2}{3}m_e$  – preon dipoles.

Dissociated DIRACs can only exist at nuclear densities or at high temperatures. A DIRAC breaks into two Dirac monopoles with mass  $\frac{m_0}{2}$  and charge  $\mu = \frac{e}{2\alpha}$  (see Section 3.2). These monopoles are the smallest building blocks of fractal structure of constituent quarks and hadrons (mesons and baryons).

ELOPs break into two preons whose mass  $m_{pr}$  equals to one third of an electron's mass:

$$m_{pr} = \frac{1}{3}m_e \tag{2.9.1}$$

and charge  $e_{pr}$  – to one third of an electron's charge:

$$e_{pr} = \frac{1}{3}e \tag{2.9.2}$$

Preons are the smallest building blocks of a fractal structure of quarks and leptons.

According to Wikipedia [Preon]: *In particle physics, preons are postulated “point-like” particles, conceived to be subcomponents of quarks and leptons.*<sup>8</sup>

We did not take into account the binding energies of DIRACs and ELOPs, and thus the values of their masses are approximate. They have negligible electrostatic and electromagnetic charges because the separation between charges is very small. They do however possess electrostatic and electromagnetic dipole momentum (see Section 3.2).

In our Model, DM particle masses are proportional to  $m_0$  multiplied by different exponents of  $\alpha$ . Consequently, we can predict the masses of various types of DM particles:

CDM particles (Neutralinos and WIMPs):

$$m_N = \alpha^{-2}m_0 = 1.3149950 \text{ TeV}/c^2 \quad 2.9.3$$

$$m_{WIMP} = \alpha^{-1}m_0 = 9.5959823 \text{ GeV}/c^2 \quad 2.9.4$$

DIRACs:

$$m_{DIRAC} = 2\alpha^0 \frac{m_0}{2} = 70.025267 \text{ MeV}/c^2 \quad 2.9.5$$

ELOPs:

$$m_{ELOP} = 2\alpha^1 \frac{m_0}{3} = 340.66606 \text{ keV}/c^2 \quad 2.9.6$$

WDM particles (sterile neutrinos):

$$m_{\nu_s} = \alpha^2 m_0 = 3.7289402 \text{ keV}/c^2 \quad 2.9.7$$

These values fall into the ranges estimated in literature.

Our Model holds that the energy densities of all types of DM particles are proportional to the proton energy density in the World’s Medium:

$$\rho_p = \frac{2\pi^2\alpha}{3}\rho_{cr} \quad 2.9.8$$

In all, there are 5 different types of DM particles. Then the total energy density of DM is

$$\rho_{DM} = 5\rho_p = 0.24007327\rho_{cr} \quad 2.9.9$$

which is close to the DM energy density discussed in literature:  $\rho_{DM} \cong 0.23\rho_{cr}$  [Wikipedia, Dark Matter].

The total neutrino energy density (in the Medium and in macroobjects, see Section 2.7) equals to

---

<sup>8</sup> I. A. D’Souza, C. S. Kalman, Preons: Models of Leptons, Quarks and Gauge Bosons as Composite Objects. World Scientific (1992). ISBN 978-981-02-1019-9.

$$\rho_{vtot} = \frac{45}{\pi} \rho_p \quad 2.9.10$$

The total baryonic energy density  $\rho_B$  is:

$$\rho_B = 1.5\rho_p \quad 2.9.11$$

The sum of MBR photons, black body radiation from cosmic dust, X-rays, and Gamma rays energy densities equals to

$$\rho_{rad} = \left(4 + \frac{1}{5\pi}\right) \frac{\alpha}{\beta} \rho_p \quad 2.9.12$$

We chose the above value of  $\rho_{rad}$  so that the energy density of the World  $\rho_W$  equals to the theoretical critical energy density  $\rho_{cr}$  in accordance with the principal idea of our Model:

$$\rho_W = \left[\frac{13}{2} + \left(\frac{11}{2} + \frac{1}{5\pi}\right) \frac{\alpha}{\beta} + \frac{45}{\pi}\right] \rho_p = \rho_{cr} \quad 2.9.13$$

From equation 2.9.13 we can calculate the value of the FSC, using electron-to-proton mass ratio

$$\frac{1}{\alpha} = \frac{\pi}{15} \left[450 + 65\pi + (55\pi + 2) \frac{m_e}{m_p}\right] = 137.03600 \quad 2.9.14$$

which is in an excellent agreement with the commonly adopted value:

$$\frac{1}{\alpha} = 137.035999074(44) \quad 2.9.15$$

It follows that there are direct correlations between constants  $\alpha$ ,  $\beta$ , and  $\frac{m_e}{m_p}$  expressed by equation of the total energy density of the World (2.9.13).

As shown above,  $\beta$  and  $\frac{m_e}{m_p}$  are not independent constants, but are instead derived from  $\alpha$ . We will, however, continue to use  $\beta$  for convenience.

The main suggestion for experimentalists dealing with observations of Dark Matter is to concentrate their efforts on particles possessing masses shown above.

## 2.10. MACROOBJECT CORES BUILT UP FROM FERMIONIC DARK MATTER

---

According to Wikipedia [Compact star]: *In astronomy, the term Compact Star (sometimes compact object) is used to refer collectively to white dwarfs, neutron stars, other exotic dense stars, and black holes. The term compact star is often used when the exact nature of the star is not known, but evidence suggests that it is very massive and has a small radius.*

In this section, we discuss the possibility of all macroobject cores consisting of Dark Matter particles introduced in section 2.9. In our view, all macroobjects of the World (including galaxy clusters, galaxies, star clusters, extrasolar systems, and planets) possess the following properties:

- Macroobject cores are made up of DM particles;
- Macroobjects consist of all particles under consideration, in the same proportion as they exist in the World's Medium;
- Macroobjects contain other particles, including DM and baryonic matter, in shells surrounding the cores.

The first phase of stellar evolution in the history of the World may be dark stars, powered by Dark Matter heating rather than fusion. Neutralinos and WIMPs, which are their own antiparticles, can annihilate and provide an important heat source for the stars and planets in the World.

Taking into account the main principle of World – Universe Model (all equations should contain  $\alpha$ ,  $\beta$ ,  $Q$ , small integer numbers and  $\pi$ ) we modify the published theory of fermionic compact stars developed by G. Narain *et al*<sup>9</sup> as follows. We'll take a scaling solution for a free Fermi gas consisting of fermions with mass  $m_f$  in accordance with following equations:

$$\text{Maximum mass: } M_{max} = A_1 M_F; \tag{2.10.1}$$

$$\text{Minimum radius: } R_{min} = A_2 R_F; \tag{2.10.2}$$

$$\text{Maximum density: } \rho_{max} = A_3 \rho_0 \tag{2.10.3}$$

where

$$M_F = \frac{M_P^3}{m_f^2}; \quad R_F = \frac{M_P L_{cf}}{m_f 2\pi}; \quad \rho_0 = \frac{hc}{a^4} \tag{2.10.4}$$

and  $L_{cf}$  is a Compton length of the fermion.  $A_1$ ,  $A_2$ , and  $A_3$  are parameters.

---

<sup>9</sup> G. Narain, J. Schaffner-Bielich, and I. N. Mishustin (2006), astro-ph/0605724.

Let us choose  $\pi$  as the value of  $A_2$  (instead of  $A_2 = 3.367$  taken by G. Narain *et al* <sup>9</sup>). Then diameter of CS is proportional to the fermion Compton length  $L_{cf}$ . We use  $\frac{\pi}{6}$  as the value of  $A_1$  (instead of  $A_1 = 0.384$  taken by G. Narain *et al* <sup>9</sup>). Then  $A_3$  will equal to

$$A_3 = \left(\frac{m_f}{m_0}\right)^4 \quad 2.10.5$$

Table 1 summarizes the parameter values for Compact Stars made up of various fermions:

**Table 1**

<b>Fermion</b>	<b>Fermion relative mass</b>	<b>Macroobject relative mass</b>	<b>Macroobject relative radius</b>	<b>Macroobject relative density</b>
	$m_f/m_0$	$M_{max}/M_0$	$R_{min}/L_g$	$\rho_{max}/\rho_0$
<b>Muonic neutrino</b>	$Q^{-\frac{1}{4}}$	$Q^{\frac{1}{2}}$	$Q^{\frac{1}{2}}$	$Q^{-1}$
<b>Tauonic neutrino</b>	$6 \times Q^{-\frac{1}{4}}$	$6^{-2} \times Q^{\frac{1}{2}}$	$6^{-2} \times Q^{\frac{1}{2}}$	$6^4 \times Q^{-1}$
<b>Sterile neutrino</b>	$\alpha^2$	$\alpha^{-4}$	$\alpha^{-4}$	$\alpha^8$
<b>Preon</b>	$3^{-1}\alpha^1$	$3^2\alpha^{-2}$	$3^2\alpha^{-2}$	$3^{-4}\alpha^4$
<b>Electron-proton (white dwarf)</b>	$\alpha^1, \beta$	$\beta^{-2}$	$(\alpha\beta)^{-1}$	$\alpha^3\beta$
<b>Monopole</b>	$2^{-1}$	$2^2$	$2^2$	$2^{-4}$
<b>WIMP</b>	$\alpha^{-1}$	$\alpha^2$	$\alpha^2$	$\alpha^{-4}$
<b>Neutralino</b>	$\alpha^{-2}$	$\alpha^4$	$\alpha^4$	$\alpha^{-8}$
<b>Interacting WIMPs</b>	$\alpha^{-1}$	$\beta^{-2}$	$\beta^{-2}$	$\beta^4$
<b>Interacting neutralinos</b>	$\alpha^{-2}$	$\beta^{-2}$	$\beta^{-2}$	$\beta^4$
<b>Neutron (star)</b>	$\approx \beta$	$\beta^{-2}$	$\beta^{-2}$	$\beta^4$

where

$$M_0 = \frac{4\pi m_0}{3} \times Q^{\frac{3}{2}} \quad 2.10.6$$

$$L_g = a \times Q^{\frac{1}{2}} \quad 2.10.7$$

The maximum density of neutron stars equals to the nuclear density

$$\rho_{max} = \left(\frac{m_p}{m_0}\right)^4 \rho_0 = \beta^4 \rho_0 \quad 2.10.8$$

which is the maximum possible density of any macroobject in the World.

A Compact Star made up of heavier particles – WIMPs and neutralinos – could in principle have a much higher density. In order for such a star to remain stable and not exceed the



nuclear density, WIMPs and neutralinos must partake in an annihilation interaction whose strength equals to  $\frac{1}{\alpha}$  and  $\frac{1}{\alpha^2}$  respectively.

Scaling solution for interacting WIMPs can also be described with equations 2.10.1 – 2.10.3 and the following values of  $A_1$ ,  $A_2$  and  $A_3$ :

$$A_{1max} = \frac{\pi}{6} (\alpha\beta)^{-2} \quad 2.10.9$$

$$A_{2min} = \pi (\alpha\beta)^{-2} \quad 2.10.10$$

$$A_{3max} = \beta^4 \quad 2.10.11$$

The maximum mass and minimum radius increase about two orders of magnitude each and the maximum density equals to nuclear density. Note that parameters of a CS made up of strongly interacting WIMPs are identical to those of neutron stars.

In accordance with the paper by G. Narain *et al*<sup>9</sup>, the most attractive feature of the strongly interacting Fermi gas of WIMPs is practically constant value of CS minimum radius in the large range of masses  $M_{WIMP}$  from

$$M_{WIMPmax} = \frac{\pi}{6} (\alpha\beta)^{-2} M_F = \frac{1}{\beta^2} M_0 \quad 2.10.12$$

down to

$$M_{WIMPmin} = \alpha^4 M_{WIMPmax} \quad 2.10.13$$

$M_{WIMPmin}$  is more than eight orders of magnitude smaller than  $M_{WIMPmax}$ . It makes strongly interacting WIMPs good candidates for stellar and planetary cores (see Sections 2.14, 2.16).

When the mass of a CS made up of WIMPs is much smaller than the maximum mass, the scaling solution yields the following equation for parameters  $A_1$  and  $A_2$ :

$$A_1 A_2^3 = \pi^4 \quad 2.10.14$$

Compare  $\pi^4 \cong 97.4$  with the value of 91 used by G. Narain *et al*<sup>9</sup>.

When the density of a CS is lower than the nuclear density and WIMP annihilation reaction does not get initiated, we obtain the following values of minimum mass and maximum radius:

$$A_{1min} = \frac{\pi}{6} \sqrt{6} (\alpha\beta)^2 \quad 2.10.15$$

$$A_{2max} = \pi \sqrt[6]{6} (\alpha\beta)^{-\frac{2}{3}} \quad 2.10.16$$

It follows that the range of stellar masses ( $A_{1min} \Leftrightarrow A_{1max}$ ) spans about three orders of magnitude, and the range of star core radii ( $A_{2min} \Leftrightarrow A_{2max}$ ) – one order of magnitude. It makes WIMPs good candidates for brown dwarf cores too (see Section 2.15).

Scaling solution for interacting neutralinos can be described with the same equations (2.10.1 – 2.10.3) and the following values of  $A_1^*$ ,  $A_2^*$  and  $A_3^*$ :

$$A_{1max}^* = \frac{\pi}{6} (\alpha^2 \beta)^{-2} \quad 2.10.17$$

$$A_{2min}^* = \pi (\alpha^2 \beta)^{-2} \quad 2.10.18$$

$$A_{3max}^* = \beta^4 \quad 2.10.19$$

In this case, the maximum mass and minimum radius increase about four orders of magnitude each and the maximum density equals to the nuclear density. Note that parameters of a CS made up of strongly interacting neutralinos are identical to those of neutron stars.

Practically constant value of CS minimum radius takes place in the huge range of masses  $M_N$  from

$$M_{Nmax} = \frac{\pi}{6} (\alpha \beta)^{-2} \alpha^2 M_F = \frac{1}{\beta^2} M_0 \quad 2.10.20$$

down to

$$M_{Nmin} = \alpha^8 M_{Wmax} \quad 2.10.21$$

$M_{WIMPmin}$  is more than seventeen orders of magnitude smaller than  $M_{Nmax}$ . It makes strongly interacting neutralinos good candidates for stellar and planetary cores (see Sections 2.14, 2.16).

When the mass of a CS made up of neutralinos is much smaller than the maximum mass, the scaling solution yields the following equation for parameters  $A_1^*$  and  $A_2^*$ :

$$A_1^* A_2^{*3} = \pi^4 \quad 2.10.22$$

When the density of a CS is lower than the nuclear, we obtain the following values of minimum mass and maximum radius:

$$A_{1min}^* = \frac{\pi}{6} \sqrt{6} (\alpha^2 \beta)^2 \quad 2.10.23$$

$$A_{2max}^* = \pi \sqrt[6]{6} (\alpha^2 \beta)^{-\frac{2}{3}} \quad 2.10.24$$

It means that the range of stellar masses ( $A_{1min}^* \Leftrightarrow A_{1max}^*$ ) is about twelve orders of magnitude, and the range of star core radii ( $A_{2min}^* \Leftrightarrow A_{2max}^*$ ) is about four orders of magnitude.

The numerical values for CS masses and radii will be given in Section 2.11.

Fermionic Compact Stars (FCS) have the following properties:

- The maximum potential of interaction  $U_{max}$  between any particle or macroobject and FCS made up of any fermions

$$U_{max} = \frac{GM_{max}}{R_{min}} = \frac{c^2}{6} \quad 2.10.25$$

does not depend on the nature of the fermions;

- The minimum radius of FCS made of any fermion

$$R_{min} = 3R_{SH} \quad 2.10.26$$

equals to three Schwarzschild radii and does not depend on the nature of the fermion;

- FCS density does not depend on  $M_{max}$  and  $R_{min}$  and does not change in time while  $M_{max} \propto t^{\frac{3}{2}}$  and  $R_{min} \propto t^{\frac{1}{2}}$ .

Boson stars made up of bosonic DM are discussed in literature (see, for example, the paper by J. Ho *et al*<sup>10</sup>) as an alternative to black holes. Axions with mass  $m_a$  introduced in Section 2.6. are good candidates for such compact macroobjects:

$$m_a = \left(\frac{\alpha}{\beta}\right)^{\frac{1}{2}} m_0 \times Q^{-\frac{1}{2}} \quad 2.10.27$$

We calculate maximum mass  $M_{Bmax}$ , radius  $R_B$ , and density  $\rho_{Bmax}$ :

$$M_{Bmax} \sim \frac{M_P^2}{m_a} = 4 \left(\frac{\beta}{\alpha}\right)^{\frac{1}{2}} m_0 \times Q^{\frac{3}{2}} = \frac{3}{\pi} \left(\frac{\beta}{\alpha}\right)^{\frac{1}{2}} M_0 \quad 2.10.28$$

$$R_B \sim \frac{h}{m_a c} = \left(\frac{\beta}{\alpha}\right)^{\frac{1}{2}} L_g \quad 2.10.29$$

$$\rho_{Bmax} \sim \frac{\alpha}{\beta} \rho_0 \quad 2.10.30$$

Boson stars made up of axions are good candidates for the cores of star clusters. These stars have a constant density in time, similar to fermionic compact stars.

---

<sup>10</sup> J. Ho, S. Kim, and B. Lee (1999), arXiv: gr-qc/9902040.

## 2.11. FRACTAL COSMOLOGY

---

*All attempts to explain the workings of the universe without recognizing the existence of the ether and the indispensable function it plays in the phenomena are futile and destined to oblivion.*

*There is no energy in matter other than that received from the environment.*

Nikola Tesla

Yu. Baryshev and P. Teerikorpi <sup>11</sup> have this to say about fractal cosmology:

*A fundamental task of practical cosmology is to study how matter is distributed in space and how it has evolved in cosmic time. The discovery of the strongly inhomogeneous spatial distribution of galaxies, at scales from galaxies to Superclusters, i.e. over four orders of magnitude in scale, was of profound cosmological significance.*

*The debate on the fractality of the large scale galaxy distribution has been going on around two new fundamental empirical cosmic numbers, - the fractal dimension  $D$ , which determines the global mass-radius behavior of the Universe:*

$$M(r) \propto r^D \qquad 2.11.1$$

*and the bordering scale where fractality transforms into homogeneity  $R_{hom}$ . Their values have been debated, and  $D = 1.2$  indirectly deduced from angular catalogues has been replaced by  $D = 2.2 \pm 0.2$  obtained from 3-d maps. The discussion of galaxy clustering started from scales 1 Mpc – 10 Mpc, then observations of the large scale structure have shifted to the scales of 10 Mpc – 100 Mpc, and now we are entering gigantic scales of 100 Mpc – 1000 Mpc.*

*In the realm of physics real structures usually have a lower scale  $R_{min}$  and an upper scale  $R_{max}$  between which the physical system follows fractal self-similar behavior. These scales are called lower and upper cutoffs.*

*For different cosmological problems there could be different choices of the lower cutoff: dark matter clumps of  $(10^6 - 10^8)M_{Sun}$ , stars, comet-size objects, atoms, elementary particles.*

*The upper cutoff presents a much more complicated problem in studies of the galaxy distribution. Is there an upper cutoff for the large-scale galaxy distribution and what is its value? These are the primary questions around which the most acute discussion is going on.*

*The fractal mass-radius law of galaxy clustering has become a key phenomenon in observational cosmology. It creates novel challenges for theoretical understanding of the origin and evolution of the galaxy distribution, including the role of dark matter.*

---

<sup>11</sup> Yu. Baryshev, and P. Teerikorpi (2005), astro-ph/0505185.

Walls and filaments are the largest known structures in the World. The Great Wall is a sheet of galaxies more than 500 million light-years long and 200 million wide (but only 15 million light-years thick). The Sloan Great Wall is up to 1.5 billion light-years across. On January 11, 2013, a large quasar group, the Huge-LOG, was discovered. It was measured to be four billion light-years across, and is presently the largest known structure in the World [Wikipedia, Observable universe].

*In astronomy, voids are the vast empty spaces between filaments (the largest-scale structures in the Universe), which contain very few, or no, galaxies [Wikipedia, Void (astronomy)]. A Supervoid in the constellation Eridanus is possibly a billion light-years across. In our opinion, voids are the Medium of the World in its purest.*

Superclusters are largest known grouping of galaxies. The Local Supercluster (Virgo Supercluster), for example, contains over 47,000 galaxies, is about  $10^{24}$  m across and weighs  $\sim 10^{17}$  solar masses ( $10^{47}$  kg) [Wikipedia, Supercluster];

Galaxy clusters contain 50 to 1,000 galaxies. Galaxy clusters have diameters of  $\sim 10^{23}$  m and total masses of  $10^{14}$  to  $10^{15}$  solar masses ( $10^{44} \leftrightarrow 10^{45}$ ) kg [Wikipedia, Galaxy cluster];

Groups of galaxies typically contain no more than 50 galaxies, and have a diameter of  $\sim 10^{22}$  m and weigh in at  $\sim 10^{13}$  solar masses ( $\sim 10^{43}$  kg) [Wikipedia, Galaxy groups and clusters];

Galaxies range from dwarfs with as few as  $10^7$  stars to giants containing  $10^{14}$  stars, each orbiting their galaxy's own center of mass. There are more than  $1.7 \times 10^{11}$  galaxies in the World. Most galaxies are 3,000 to 300,000 light-years in diameter. Galaxies are usually separated by distances on the order of 3 million light-years. Ultra-compact dwarf galaxies have recently been discovered that are only 300 light-years across [Wikipedia, Galaxy].

Two types of Star Clusters can be distinguished: globular clusters are tight groups of hundreds of thousands of very old stars which are gravitationally bound, while open clusters, more loosely clustered groups of stars, generally contain fewer than a few hundred members, and are often very young [Wikipedia, Star cluster].

Extrasolar systems range from brown dwarfs with minimum mass of about 0.013 solar masses ( $2.6 \times 10^{28}$  kg) [Wikipedia, List of least massive stars], red dwarfs with the minimum mass about of 0.075 solar masses ( $1.5 \times 10^{29}$  kg) [Wikipedia, Red dwarf], to giant stars that are 150 times as massive as the Sun ( $3 \times 10^{32}$  kg) [Wikipedia, Star].

The following table summarizes the various macroobjects:

Macroobject	Size (m)	Mass (kg)
World	$10^{26}$	$3 \times 10^{52}$
Walls, Filaments	$10^{24} \Leftrightarrow 10^{25}$	$10^{49} \Leftrightarrow 10^{51}$
Supercluster	$10^{24}$	$10^{47} \Leftrightarrow 10^{48}$
Galaxy cluster	$10^{23}$	$10^{45} \Leftrightarrow 10^{46}$
Group of galaxies	$10^{22}$	$10^{43} \Leftrightarrow 10^{44}$
Galaxy	$10^{19} \Leftrightarrow 10^{21}$	$10^{39} \Leftrightarrow 10^{43}$
Star cluster	$10^{17} \Leftrightarrow 10^{18}$	$10^{35} \Leftrightarrow 10^{37}$
Extra solar system	$10^{14} \Leftrightarrow 10^{16}$	$10^{29} \Leftrightarrow 10^{33}$

According to World – Universe Model, the total macroobject energy  $E_{MO}$  enclosed in surface  $S_{MO}$  is proportional to the area of that surface:

$$E_{MO} = \sigma_0 S_{MO} \quad 2.11.2$$

where  $\sigma_0$  is the surface enthalpy defined in Section 2.4. All the energy contained in macroobjects was received from the environment.

In case when the stars and galaxies are distributed in a hierarchy of spherical clusters of radius  $R_{MO}$ , the energy  $E_{MO}$  equals to

$$E_{MO} = 4\pi\sigma_0 R_{MO}^2 \quad 2.11.3$$

Comparing this result with equation 2.11.1 we conclude that the World has a fractal structure with the theoretical fractal dimension  $D = 2$ , which is in good agreement with the value of  $D = 2.2 \pm 0.2$  experimentally obtained by P. Teerikorpi *et al*<sup>12</sup>. Note that the Olbers' paradox (dark night sky) can be explained only if the fractal dimension of the World  $D \leq 2$  [Wikipedia, Olbers' paradox].

The upper cutoff of the fractal structure is the entire World, with its total mass of  $M_W$  and radius  $R = R_{hom}$ :

$$M_W = \frac{4\pi\sigma_0}{c^2} R^2 = 4\pi m_0 \times Q^2 = 9.0640 \times 10^{52} \text{ kg} \quad 2.11.4$$

$$R = a \times Q = 1.3459 \times 10^{26} \text{ m} \quad 2.11.5$$

The lower cutoff of the fractal structure is an extrasolar system (ESS) with total mass  $M_{ESS}$ , radius  $R_{ESS}$ , and number  $N_{ESS}$  in the following ranges:

$$M_{ESS} = 4\pi m_0 \times Q^{\frac{3}{2}} \times (\alpha^2 \Leftrightarrow 1) =$$

$$(5.5360 \times 10^{28} \Leftrightarrow 1.0396 \times 10^{33}) \text{ kg} \quad 2.11.6$$

---

<sup>12</sup> P. Teerikorpi, M. Hanski, G. Theureau, Yu. Baryshev, G. Paturel, L. Bottinelli, L. Gouguenheim, A&A 334, 395 (1998).

$$R_{ESS} = a \times Q^{\frac{3}{4}} \times (\alpha \Leftrightarrow 1) =$$

$$(1.0518 \times 10^{14} \Leftrightarrow 1.4414 \times 10^{16}) m \quad 2.11.7$$

$$N_{ESS} \sim Q^{\frac{1}{2}} \times (1 \Leftrightarrow \alpha^{-2}) \sim (10^{20} \Leftrightarrow 10^{24}) \quad 2.11.8$$

$N_{ESS}$  is the total number of extra-solar systems in the World. Note that an ESS receives all of its energy from its environment (Galaxy).

According to our Model, all macroobjects of the World (galaxies, stars, planets) have cores made up of Dark Matter particles. The theory of fermion compact stars (FCS) made up of Dark Matter particles is well developed. Scaling solutions are derived for a free and an interacting Fermi gas in Section 2.10.

Table 2 describes the parameters of FCS made up of different fermions:

**Table 2**

<b>Fermion</b>	<b>Fermion mass</b>	<b>Macroobject mass</b>	<b>Macroobject radius</b>	<b>Macroobject density</b>
	$m_f, MeV/c^2$	$M_{max}, kg$	$R_{min}, m$	$\rho_{max}, kg/m^3$
Muonic neutrino	$7.50 \times 10^{-9}$	$3.0 \times 10^{52}$	$1.3 \times 10^{26}$	$3.0 \times 10^{-27}$
Tauonic neutrino	$4.50 \times 10^{-8}$	$8.4 \times 10^{50}$	$3.7 \times 10^{24}$	$3.8 \times 10^{-24}$
Sterile neutrino	$3.73 \times 10^{-3}$	$1.2 \times 10^{41}$	$5.4 \times 10^{14}$	$1.8 \times 10^{-4}$
Preon	0.170	$5.9 \times 10^{37}$	$2.6 \times 10^{11}$	$7.8 \times 10^2$
Monopole	35.01	$1.4 \times 10^{33}$	$6.2 \times 10^6$	$1.4 \times 10^{12}$
Interacting WIMPs	9,596	$1.9 \times 10^{30}$	$8.6 \times 10^3$	$7.2 \times 10^{17}$
Interacting neutralinos	$1,315 \times 10^3$	$1.9 \times 10^{30}$	$8.6 \times 10^3$	$7.2 \times 10^{17}$
Electron-proton (white dwarf)	0.511-938.3	$1.9 \times 10^{30}$	$1.6 \times 10^7$	$1.2 \times 10^8$
Neutron (star)	939.6	$1.9 \times 10^{30}$	$8.6 \times 10^3$	$7.2 \times 10^{17}$

The calculated parameters of FCS show that

- White Dwarf Shell (WDS) around the core made of strongly interacting WIMPs or neutralinos (see Section 2.14) compose the nucleus of stars in extrasolar systems;
- Dissociated DIRACs to Monopoles form cores of star clusters;
- Dissociated ELOPs to Preons constitute nuclei of galaxies;

- Sterile neutrinos make up cores of galaxy clusters;
- Tauonic neutrinos reside in the cores of galaxy superclusters.

Interestingly, the calculated radius of an FCS made up of muonic neutrinos exactly equals to the radius of the World, while its mass would equal the combined mass of all the World macroobjects.

Although there are no free Dirac's monopoles and preons in the World, they can arise in the cores of FCS as the result of DIRACs and ELOPs gravitational collapse with density increasing up to the nuclear density ( $\sim 10^{17} \frac{kg}{m^3}$ ) and/or at high temperatures, with subsequent dissociation of dipoles to monopoles and preons.

To summarize, stars macroobjects of the World have cores made up of the discussed DM particles. Other particles, including DM and baryonic matter, form shells surrounding the cores. In our Model, all macroobjects consist of all particles under consideration, in the same proportion as they exist in the World's Medium. There are no compact stars made up solely of DM fermionic particles, for instance.

---

## 2.12. FRACTAL STRUCTURE OF THE WORLD

---

The Model provides us with a facility to calculate the masses, sizes, and numbers of the World's cluster structures that follows fractal self-similar behavior. Galaxy clusters (GC) have total mass  $M_{GC}$ , radius  $R_{GC}$ , and number  $N_{GC}$  in the following ranges:

$$\begin{aligned}
 M_{GC} &= 4\pi m_0 \times Q^{\frac{15}{8}} \times (\alpha^2 \Leftrightarrow 1) = \\
 &= (4.9950 \times 10^{43} \Leftrightarrow 9.3801 \times 10^{47}) \text{ kg}
 \end{aligned} \tag{2.12.1}$$

$$\begin{aligned}
 R_{GC} &= a \times Q^{\frac{15}{16}} \times (\alpha \Leftrightarrow 1) = \\
 &= (3.1595 \times 10^{21} \Leftrightarrow 4.3297 \times 10^{23}) \text{ m}
 \end{aligned} \tag{2.12.2}$$

$$N_{GC} \sim Q^{\frac{1}{8}} \times (1 \Leftrightarrow \alpha^{-2}) \sim (10^5 \Leftrightarrow 10^9) \tag{2.12.3}$$

Galaxies (G) have total mass  $M_G$ , radius  $R_G$ , and number  $N_G$  in the following ranges:

$$\begin{aligned}
 M_G &= 4\pi m_0 \times Q^{\frac{7}{4}} \times (\alpha^2 \Leftrightarrow 1) = \\
 &= (5.1692 \times 10^{38} \Leftrightarrow 9.7073 \times 10^{42}) \text{ kg}
 \end{aligned} \tag{2.12.4}$$

$$\begin{aligned}
 R_G &= a \times Q^{\frac{7}{8}} \times (\alpha \Leftrightarrow 1) = \\
 &= (1.0164 \times 10^{19} \Leftrightarrow 1.3928 \times 10^{21}) \text{ m}
 \end{aligned} \tag{2.12.5}$$



$$N_G \sim Q^{\frac{1}{4}} \times (1 \Leftrightarrow \alpha^{-2}) \sim (10^{10} \Leftrightarrow 10^{14}) \quad 2.12.6$$

For star clusters (SC) we obtain total mass  $M_{SC}$ , radius  $R_{SC}$ , and number  $N_{SC}$  in the following ranges:

$$\begin{aligned} M_{SC} &= 4\pi m_0 \times Q^{\frac{13}{8}} \times (\alpha^2 \Leftrightarrow 1) = \\ &= (5.3496 \times 10^{33} \Leftrightarrow 1.0046 \times 10^{38}) \text{ kg} \end{aligned} \quad 2.12.7$$

$$\begin{aligned} R_{SC} &= a \times Q^{\frac{13}{16}} \times (\alpha \Leftrightarrow 1) = \\ &= (3.2696 \times 10^{16} \Leftrightarrow 4.4806 \times 10^{18}) \text{ m} \end{aligned} \quad 2.12.8$$

$$N_{SC} \sim Q^{\frac{3}{8}} \times (1 \Leftrightarrow \alpha^{-2}) \sim (10^{15} \Leftrightarrow 10^{19}) \quad 2.12.9$$

When stars and galaxies are distributed in a hierarchy of disk-shape clusters, the calculated radii  $R_{MO}$  should be multiplied by  $\sqrt{2}$ .

The calculated ranges of radii, masses, and numbers of the World, GC, G, SC, and ESS are in good agreement with literature estimates. Our calculations show that the distance separating the galaxies is approximately  $10^{21}$  m, which is in good agreement with experimentally measured distances. The distance from the Milky Way to the Large Magellanic Cloud, for instance, is about  $1.5 \times 10^{21}$  m [Wikipedia, List of nearest galaxies].

Within a galaxy, we calculate the distances between the stars to be about  $10^{16}$  m. The distance from the Sun to the Proxima Centauri is about  $4 \times 10^{16}$  m [Wikipedia, List of nearest stars].

The central macroobject (CMO) of a galaxy has a core made up of preons. Our calculations show that its mass is smaller than  $5.9 \times 10^{37}$  kg, and its radius is greater than  $2.6 \times 10^{11}$  m. From the movement of S2 star it was estimated that our own Milky Way's central object mass is about 4.1 million solar masses ( $8.2 \times 10^{36}$  kg), and its radius is no larger than  $6.7 \times 10^{12}$  m [Wikipedia, Sagittarius A\*].

In our Model it is natural to define surface  $S_{MO}$  as the boundary between macroobject and surrounding environment. In case of our Solar system such a surface is named Heliosphere [Wikipedia, Heliosphere]. We will refer to such surfaces as Macroobject Boundary (MOB). The radii  $R$ ,  $R_{GC}$ ,  $R_G$ ,  $R_{SC}$ ,  $R_{ESS}$  introduced above are really radii of corresponding Macroobject boundaries.

According to the developed Model, CMOs have cores made up of fermionic DM particles possessing radii  $R_{CORE}$  described in Tables 1 & 2. In case of extrasolar systems, the cores are made up of interacting neutralinos or WIMPs surrounded with white dwarf shells (WDS).

Surrounding the cores, there is a transitional region in which the density decreases rapidly to the point of the zero level of the fractal structure<sup>13</sup> characterized by radius  $R_f$  and energy density  $\rho_f$  that satisfy the following equation for  $r \geq R_f$ :

$$\rho(r) = \frac{\rho_f R_f}{r} \quad 2.12.10$$

According to Yu. Baryshev: *For a structure with fractal dimension  $D = 2$  the constant  $\rho_f R_f$  may be actually viewed as a new fundamental physical constant.*<sup>13</sup> Our Model allows us to calculate the value of this constant. Recall that 1/3 of the total macroobject energy resides inside of the CMO (star in case of an ESS), and 2/3 of it belongs to the fractal structure above CMO. It follows that

$$2\pi\rho_f R_f R_{MO}^2 = \frac{2}{3} \times 4\pi\sigma_0 R_{MO}^2 \quad 2.12.11$$

and the value of  $\rho_f R_f$  is:

$$\rho_f R_f = \frac{4}{3} \sigma_0 \quad 2.12.12$$

As for the values of  $R_f$  and  $\rho_f$ , let us take

$$R_f = \alpha^{-1} R_{CORE} \quad 2.12.13$$

and

$$\rho_f = \frac{4}{3} \sigma_0 \frac{\alpha}{R_{CORE}} \quad 2.12.14$$

Equation 2.12.10 fits naturally into our Model, since the evolution of all spherical structures of the World is progressing in a quasi-stationary mode (see Section 2.13). The ball of radius  $R_f$  is absorbing energy from the environment, and the distribution of energy outside of the ball follows equation 2.12.10.

The calculations carried out for our Sun using equations 2.12.13 and 2.12.14 are in agreement with the experimentally measured characteristics of the Sun. Taking the value of the solar core radius  $R_{CORESun} \cong 1.6 \times 10^8 \text{ m}$  (see 2.14.16) we obtain

$$R_f \cong 2.2 \times 10^{10} \text{ m} \quad 2.12.15$$

which is in agreement with estimated sizes of the Heliosphere. *The Heliosphere, which is the cavity around the Sun filled with the solar wind plasma, extends from approximately 20 solar radii ( $\sim 1.4 \times 10^{10} \text{ m}$ ) to the outer fringes of the Solar System* [Wikipedia, Sun].

According to 2.12.14, the mass density  $\rho_{fm}$  at radius  $R_f$  is

---

<sup>13</sup> Yu. Baryshev, Practical Cosmology, 2, 60 (2008).

$$\rho_{fm} = \frac{4}{3} \frac{\sigma_0}{c^2 R_f} \cong 2.4 \times 10^{-11} \frac{kg}{m^3} \quad 2.12.16$$

and the minimum mass density  $\rho_{fmin}$  at the boundary of a macroobject is:

$$\rho_{fmin} = \frac{4}{3} \sigma_0 \left( \frac{3M_{Sun} c^2}{4\pi\sigma_0} \right)^{-\frac{1}{2}} \cong 4.9 \times 10^{-16} \frac{kg}{m^3} \quad 2.12.17$$

Mass of the fractal structure around Sun  $M_V$  at distances  $R_V \gg R_f$  is

$$M_V = \frac{8\pi}{3} \sigma_0 R_V^2 \quad 2.12.18$$

At distance  $R_V = 1.8 \times 10^{13} m$  away from the Sun (approximate distance to Voyager 1<sup>14</sup>),

$$M_V \cong 1.1 \times 10^{27} kg \quad 2.12.19$$

that is  $\sim 0.05\%$   $M_{Sun}$ . This additional mass can explain the observed deceleration of Voyagers. Note that the distances traveled by Voyagers ( $\sim 10^{13} m$ ) are much smaller than the radius of the MOB  $R_{MOB}$ :

$$R_{MOB} = \left( \frac{3M_{Sun}}{4\pi\sigma_0} \right)^{\frac{1}{2}} \cong 1.1 \times 10^{15} m \quad 2.12.20$$

The strongly inhomogeneous fractal spatial distribution of matter at scales from extrasolar system to the World, i.e. over twenty orders of magnitude in scale, has profound cosmological significance.

---

### 2.13. EVOLUTION OF THE WORLD'S FRACTAL STRUCTURE

---

We will analyze the evolution of the World's fractal structure concentrating on three important types of macroobjects: extrasolar systems, galaxies, and the World.

As discussed in Section 2.11, the total macroobject energy  $E_{MO}$  enclosed in surface  $S_O$  is proportional to the area of that surface:

$$E_{MO} = \sigma_0 S_{MO} \quad 2.13.1$$

where  $\sigma_0$  is the surface enthalpy. All macroobjects receive all of their energy from their environment.

When stars and galaxies are distributed in a hierarchy of spherical clusters of radius  $R_{MO}$ , the energy  $E_{MO}$  is:

$$E_{MO} = 4\pi\sigma_0 R_{MO}^2 \quad 2.13.2$$

---

<sup>14</sup> Voyager. The interstellar Mission, June 14, 2012.

It was shown above (see Sections 2.10, 2.11) that masses of all macroobject cores and ESS are proportional to  $Q^{\frac{3}{2}}$  and are increasing in time  $\propto t^{\frac{3}{2}}$ . The total energy arriving to extrasolar systems from the environment (enclosing galaxies) is consumed solely by ESS.

All larger cosmological cluster structures are receiving more energy than required for increase of the mass of their components (CMO and ESS). The remainder of that energy is spent on creation of new macroobjects. Consider a galaxy. Its total mass  $M_G$  is proportional to  $Q^{\frac{7}{4}}$  and is increasing in time  $\propto t^{\frac{7}{4}}$ :

$$M_G \sim 4\pi m_0 \times Q^{\frac{7}{4}} \quad 2.13.3$$

CMO and ESS, however, are consuming energy  $\propto t^{\frac{3}{2}}$  only.

The World, galaxies, and extrasolar systems have the following volumes:

$$V_W = V_0 \times Q^3 \propto t^3 \quad 2.13.4$$

$$V_G \propto V_0 \times Q^{\frac{21}{8}} \propto t^{\frac{21}{8}} \quad 2.13.5$$

$$V_{ESS} \propto V_0 \times Q^{\frac{9}{4}} \propto t^{\frac{9}{4}} \quad 2.13.6$$

where  $V_0 = \frac{4\pi}{3} a^3$  is the volume of the World's Core at the Beginning ( $Q = 1$ ).

The quasi-stationary expansion of them is taking place at different rates:

$$\frac{dV_W}{dt} = 3 \frac{V_W}{t} \quad 2.13.7$$

$$\frac{dV_G}{dt} = \frac{21}{8} \frac{V_G}{t} \quad 2.13.8$$

$$\frac{dV_{ESS}}{dt} = \frac{9}{4} \frac{V_{ESS}}{t} \quad 2.13.9$$

The World, galaxies, and ESS have the following total energies:

$$E_W = 4\pi E_0 \times Q^2 \propto t^2 \quad 2.13.10$$

$$E_G \propto 4\pi E_0 \times Q^{\frac{7}{4}} \propto t^{\frac{7}{4}} \quad 2.13.11$$

$$E_{ESS} \propto 4\pi E_0 \times Q^{\frac{3}{2}} \propto t^{\frac{3}{2}} \quad 2.13.12$$

and are consuming energy at the following rates:

$$\frac{dE_W}{dt} = 2 \frac{E_W}{t} \quad 2.13.13$$

$$\frac{dE_G}{dt} = \frac{7}{4} \frac{E_G}{t} \quad 2.13.14$$

$$\frac{dE_{ESS}}{dt} = \frac{3}{2} \frac{E_{ESS}}{t} \quad 2.13.15$$

We see that the expansion rates are 1.5 times greater than energy consumption rates. Hence average densities of galaxies and extrasolar systems are decreasing with time:

$$\rho_G \propto \rho_0 \times Q^{-\frac{7}{8}} \propto t^{-\frac{7}{8}} \propto \rho_{cr} \times Q^{\frac{1}{8}} \quad 2.13.16$$

$$\rho_{ESS} \propto \rho_0 \times Q^{-\frac{3}{4}} \propto t^{-\frac{3}{4}} \propto \rho_{cr} \times Q^{\frac{1}{4}} \quad 2.13.17$$

and are about 5 and 10 orders of magnitude higher than the critical density  $\rho_{cr}$ , respectively.

The energy consumption rates are greater for galaxies relative to ESS, and for the World relative to galaxies. It follows that new stars and star clusters can be created inside of a galaxy, and new galaxies and galaxy clusters can arise in the World. Formation of galaxies and stars is not a process that concluded ages ago; instead, it is ongoing.

The amount of time  $\Delta t_{DG}$  necessary for the World to accumulate sufficient energy to create a new dwarf galaxy with mass  $M_{DG}$

$$M_{DG} = 4\pi m_0 \alpha^2 \times Q^{\frac{7}{4}} \quad 2.13.18$$

is:

$$\Delta t_{DG} = \frac{1}{2} t \alpha^2 \times Q^{-\frac{1}{4}} = 1280.2 \text{ s} \quad 2.13.19$$

Similarly, the amount of time  $\Delta t_G$  necessary to accumulate enough energy for a large new galaxy having maximum possible mass

$$M_{Gmax} = 4\pi m_0 \times Q^{\frac{7}{4}} \quad 2.13.20$$

is:

$$\Delta t_G = \frac{1}{2} t \times Q^{-\frac{1}{4}} = 2.4040 \times 10^7 \text{ s} \cong 0.76 \text{ y} \quad 2.13.21$$

Similar calculations carried out for extrasolar systems show that minimum time  $\Delta t_{BD}$  to create brown dwarf with mass  $M_{BD}$

$$M_{BD} = 4\pi m_0 \alpha^2 \times Q^{\frac{3}{2}} \quad 2.13.22$$

and minimum time  $\Delta t_{ESS}$  needed to create an extrasolar system with maximum mass

$$M_{ESS} = 4\pi m_0 \times Q^{\frac{3}{2}} \quad 2.13.23$$

are:

$$\Delta t_{BD} = \frac{4}{7} t \alpha^2 \times Q^{-\frac{1}{4}} = 1463.1 \text{ s} \quad 2.13.24$$

$$\Delta t_{ESS} = \frac{4}{7} t \times Q^{-\frac{1}{4}} = 2.7475 \times 10^7 \text{ s} \cong 0.87 \text{ y} \quad 2.13.25$$

The time needed for creation of a main sequence star like our Sun is about  $100 \Delta t_{BD} \cong 40 \text{ hrs}$ , which is consistent with the estimates of star generation in MS1358arc Galaxy made by M. Swinbank *et al.*<sup>15</sup> Within the star-forming regions of this infant galaxy, new stars were being created at a rate of about 50 main sequence stars per year – around 100 times faster than had been previously thought.

---

## 2.14. EXTRASOLAR SYSTEMS

---

There are two primary types of stars: main-sequence stars and red stars. They differ in their surface temperatures and radii:

- Red stars have cool surface temperatures:  $3,500 \Leftrightarrow 4,500 \text{ K}$  for Hypergiants, Supergiants, Giants [Wikipedia, Hypergiant, Red supergiant, Red giant], lower for Red dwarfs ( $2,300 \Leftrightarrow 3,800 \text{ K}$ ) [Wikipedia, Red dwarf], and significantly lower for Brown dwarfs ( $300 \Leftrightarrow 1,000 \text{ K}$ ) [Wikipedia, Brown dwarfs]. These stars have enormous range of radii: from  $1,650 R_{Sun}$  for Hypergiants down to  $0.08 R_{Sun}$  for Red dwarfs, and lower still for Brown dwarfs.
- Main-sequence stars have surface temperatures in the range of  $3,000 \Leftrightarrow 45,500 \text{ K}$ , and radii in the range from  $35 R_{Sun}$  for the most massive known star R136a1 [Wikipedia, R136a1] down to  $0.1 R_{Sun}$  for least heavy stars [Wikipedia, Main sequence].

As we have shown above (2.13.17), extrasolar systems (ESS) have average density  $\rho_{ESS}$  that is about 10 orders of magnitude higher than the critical density:

$$\rho_{ESS} \propto \rho_{cr} \times Q^{\frac{1}{4}} \quad 2.14.1$$

The range of ESS masses  $M_{ESS}$  is about four orders of magnitude:

$$\begin{aligned} M_{ESS} &= 4\pi m_0 \times Q^{\frac{3}{2}} (\alpha^2 \Leftrightarrow 1) \sim \\ &\sim (5 \times 10^{28} \Leftrightarrow 10^{33}) \text{ kg} \end{aligned} \quad 2.14.2$$

One third of this mass resides in macroobjects constituting an extra-solar system. Most of that mass lies in the star itself. The star and other macroobjects are composed of all particles under consideration.

---

<sup>15</sup> M. Swinbank, et al, Monthly Notices of the Royal Astronomical Society, 2009.

Extrasolar systems form from clouds of particles. Due to gravitational instability, a gravitational collapse takes place. The heaviest particles, neutralinos or WIMPs, sink first and form the core of a new star.

In our opinion, the difference between main-sequence stars and red stars lies in composition of stellar cores. Main-sequence cores are made up of neutralinos, while red star cores consist of WIMPs. As we have shown in Section 2.10, in both cases the cores' maximum mass and minimum radius equals to that of a neutron star. The fermions, however, have drastically different interaction strength of annihilation:  $\frac{1}{\alpha}$  in case of WIMPs and  $\frac{1}{\alpha^2}$  in case of neutralinos.

The Core temperature is therefore much higher in main-sequence stars whose cores are made up of neutralinos. Ignition of chain reactions developing in the surrounding shells happens much more efficiently in these stars.

Let's analyze red stars with cores made up of WIMPs, with the surrounding white dwarf and preon shells.

Taking into account the 100x increase of maximum stable mass of cores made up of strongly interacting WIMPs (see 2.10.9), we calculate the total maximum core mass  $M_{CORE}$ :

$$M_{CORE} = \frac{1}{\beta^2} M_S \cong 1.93 \times 10^{30} kg \quad 2.14.3$$

It follows that the energy density of WIMPs in the World  $\rho_{WIMP}$  equals to

$$\rho_{WIMP} = \frac{1}{\beta^2} \rho_{cr} \quad 2.14.4$$

Calculations based on results of Section 2.10 show that the maximum stellar mass  $M_{Smax}$  is

$$M_{Smax} = M_0 = 3.4654 \times 10^{32} kg (\cong 174 M_{Sun}) \quad 2.14.5$$

Stars must be massive enough to support core densities equal to the nuclear density in order to initiate strong interaction between WIMPs. The minimum stellar mass  $M_{Smin}$  equals to

$$\begin{aligned} M_{Smin} &= \sqrt{6} \alpha^4 \beta^4 M_{Smax} \cong \\ &\cong 7.8 \times 10^{28} kg (\cong 0.039 M_{Sun}) \end{aligned} \quad 2.14.6$$

$M_{Smin}$  is over four orders of magnitude smaller than  $M_{Smax}$ . These numbers are in good agreement with the commonly accepted range of red stellar masses ( $0.075 \Leftrightarrow 150 M_{Sun}$ ).

The smallest true stars (red dwarfs) have masses of *less than half that of the Sun (down to about 0.075 solar masses, below which stellar objects are brown dwarfs) and a surface*

temperature of less than 4,000 K. Red dwarfs are by far the most common type of star in the Galaxy [Wikipedia, Red dwarf].

Minimum radius of a stellar core  $R_{COREmin}$  is:

$$R_{COREmin} = \frac{1}{\beta^2} L_g \cong 8.6 \text{ km} \quad 2.14.7$$

The next heaviest particles – protons, joined by electrons – will follow WIMPs during the gravitational collapse, and form the White Dwarf Shell (WDS) around the core made of strongly interacting WIMPs. The mass of the WDS is proportional to the ratio of protons in the World:

$$M_{WDS} = 1.5 \frac{\rho_p}{\rho_{cr}} M_S \quad 2.14.8$$

Using the following equation (see 2.10.4 for reference):

$$M_{WDS} = \frac{\pi M_P^3}{6 m_p^2} \quad 2.14.9$$

we obtain the maximum mass  $M_{WDScold}$  of a cold WDS:

$$M_{WDScold} \cong 1.93 \times 10^{30} \text{ kg} (\cong M_{Sun}) \quad 2.14.10$$

Taking into account the proton-proton chain reaction with the interaction strength equal to  $\beta$ , we can estimate the increase of the maximum stable mass of the WDS in accordance with theory developed by G. Narain *et al* <sup>9</sup>:

$$M_{WDSshot} = \left(1 + \frac{\beta}{2}\right) M_{WDScold} \cong 1.49 \times 10^{31} \text{ kg} \quad 2.14.11$$

Calculated value of  $M_{WDSshot}$  is consistent with the expected protons mass obtained from the maximum star mass ( $3.4654 \times 10^{32}$  kg) with 7.2% concentration of protons ( $\cong 2.50 \times 10^{31}$  kg).

The minimum radius of cold WDS is

$$R_{WDScold} = \frac{L_g}{\alpha\beta} \cong 1.6 \times 10^7 \text{ m} \quad 2.14.12$$

Taking into account the proton-proton chain reaction for the minimum radius of WDS in accordance with the paper of G. Narain *et al* <sup>9</sup> we obtain:

$$R_{WDSshot} = \left(1 + \frac{\beta}{4}\right) R_{WDScold} \cong 7.0 \times 10^7 \text{ m} \quad 2.14.13$$

The calculated parameters of red stars can explain the characteristics of brown dwarfs, red dwarfs, and subgiants that are slightly brighter than main-sequence stars, but not as bright as true giant stars. As a side note, *subgiants are the only type of stars other than main-sequence stars believed capable of hosting life-bearing planets* [Wikipedia, Subgiant].



Enormous radii of Hypergiants (up to  $1,650 R_{Sun} \cong 10^{12} m$ ) and huge luminosity of giant stars can be explained by an additional shell of preons – particles whose charge equals to  $\frac{1}{3}e$ . They compose hot high density plasma with surface temperature in the range of 3,500  $\Leftrightarrow$  4,500 K. The minimum radius of preon shell  $R_{min} \cong 2.6 \times 10^{11} m$  (see Table 2).

The analysis of main-sequence stars whose cores are made up of neutralinos with surrounding white dwarf and preon shells shows that their cores have the same maximum mass and minimum radius as those of red stars, but much higher temperature, due to considerably greater interaction strength of annihilation of neutralinos as compared to WIMPs. The characteristics of the white dwarf shell are close to those of red stars. Much higher core temperature, however, enables main-sequence stars to have much greater surface temperature. The hottest observed star has a surface temperature of 45,500 K [Wikipedia, Main sequence].

The maximum stellar mass remains the same ( $\cong 174 M_{Sun}$ ). According to Wikipedia [List of most massive stars]: *Studying the Arches cluster, which is the densest known cluster of stars in our galaxy, astronomers have confirmed that stars in that cluster do not occur any larger than about  $150 M_{Sun}$ . One theory to explain rare ultramassive stars that exceed this limit, for example in the R136 star cluster (up to  $265 M_{Sun}$ ), is the collision and merger of two massive stars in a close binary system. If any stars still exist above (150 – 200)  $M_{Sun}$ , they would challenge current theories of stellar evolution.*

Strongly interacting Fermi gas of neutralinos has practically constant value of minimum radius in the huge range of masses  $M_N$  from

$$M_{Nmax} = \frac{\pi}{6} (\alpha\beta)^{-2} \alpha^2 M_F = \frac{1}{\beta^2} M_0 \quad 2.14.14$$

down to

$$M_{Nmin} = \alpha^8 M_{Wmax} \quad 2.14.15$$

$M_{Nmin}$  is more than seventeen orders of magnitude smaller than  $M_{Nmax}$ .

We use equations 2.10.2 and 2.10.22 to calculate WDS radius of the Sun, keeping in mind that its mass is 174 smaller than the maximum stellar mass:

$$R_{WDSsun} = \sqrt[3]{6 \frac{M_{Smax}}{M_{Sun}}} R_{WDScold} \cong 1.6 \times 10^8 m \quad 2.14.16$$

$R_{WDSsun}$  is about 0.23 solar radii, which is in good agreement with solar core radius discussed in literature (0.2  $\Leftrightarrow$  0.25 solar radii).

The developed star model explains the very low power production density produced by fusion inside of the Sun. Wikipedia humorously notes that the power output of the Sun *more nearly approximates reptile metabolism than a thermonuclear bomb* [Wikipedia, Sun]. In our Model, the core made up of strongly interacting neutralinos is the supplier of proton-

electron pairs into WDS and igniter of the proton-proton chain reaction developing in the surrounding WDS with small interaction strength  $\beta \cong 13.4$ . The energy to support neutralinos annihilation and proton fusion is coming from outside of the star (Galaxy).

With respect to the developed model of FCS (Section 2.10), the masses of the core and WDS are increasing in time  $\propto t^{\frac{3}{2}}$ :

- New neutralinos and WIMPs freely penetrate through the entire stellar envelope and get absorbed into the core.
- New protons and electrons are generated in the core as the result of neutralinos and WIMPs annihilation, and enter the WDS.

The radii of the core and WDS are increasing in time  $\propto t^{\frac{1}{2}}$ . Consequently, the density and fusion power production density remain constant in time.

Consider the closed spherical surface around the WDS. Its radius is increasing in time  $\propto t^{\frac{1}{2}}$ , and its area is increasing in time  $\propto t$ . Stellar luminosity is thus increasing in time  $\propto t$ . Taking into account that the age of the World is  $\cong 14.2$  By and the age of solar system is  $\cong 4.6$  By, it is easy to find that the young Sun's output was only 67.6% of what it is today. Literature commonly refers to the value of 70%. So-called "Faint young Sun paradox" is thus resolved [Wikipedia, Faint young Sun paradox].

The described star creation picture is consistent with a new image from ESO (European Southern Observatory) which *shows a dark cloud where new stars are forming, along with a cluster of brilliant stars that have already emerged from their dusty stellar nursery. This cloud is known as Lupus 3 and it lies about 600 light-years from Earth in the constellation of Scorpius (The Scorpion) which is one of the closest such stellar nurseries to the Sun.*

*The bright stars are young stars that have not yet started to shine by nuclear fusion in their cores and are still surrounded by glowing gas. They are probably less than one million years old. The Lupus 3 region is both fascinating and a beautiful illustration of the early stages of the life of stars.*<sup>16</sup>

An important consequence for Solar system, and in fact for all other stars in the World, is that they will never burn their "fuel" out. On the contrary, stars accumulate more fuel with time, and output more power.

The existence of supermassive objects in galactic centers is now commonly accepted. Although it is believed that the central mass is a supermassive black hole, it has not yet been firmly established. Alternative models for the supermassive dark objects in galactic centers, formed by self-gravitating non-baryonic matter composed of fermions and bosons, are widely discussed in literature.

---

<sup>16</sup> Light from the Darkness, eso 1303 (2013).

The heaviest macroobjects include a high-density preon plasma shell around their cores:

- Macroobjects with a cold preon shell emit strong radio waves. Such objects are good candidates for the compact astronomical radio sources at centers of galaxies like Sagittarius A\* in the Milky Way Galaxy [Wikipedia, Sagittarius A\*].
- Red Giants are macroobjects with hot preon shells.
- Macroobjects with a very hot preon shell are candidates for Active Galactic Nuclei (AGN).

Note that the temperature of the preon shell depends on the composition of the macroobject core. Macroobjects whose cores are made up solely of preons remain cold. Macroobjects with cores made up of WIMPs and WDS produce hot preon shells. Macroobjects whose cores consist of neutralinos and WDS have very hot preon shells.

The radius of the AGN is about four orders of magnitude larger than the radius of WDS (see Table 2). The area of the closed spherical surface around the AGN is more than 8 orders of magnitude greater than the surface area of WDS. Luminosity of the AGN is then at least 8 orders of magnitude higher than the luminosity of the largest star.

The described model of AGN can explain the fact that *the most luminous quasars radiate at a rate that can exceed the output of average galaxies, equivalent to two trillion ( $2 \times 10^{12}$ ) suns* [Wikipedia, Quasar].

New protons and electrons are penetrating from the core into WDS as the result of neutralinos and WIMPs annihilation, and then emanating from the star itself. The Sun produces solar wind; hottest macroobjects such as an AGN may be emitting protons at relativistic speeds.

---

## 2.15. BROWN DWARFS

---

According to Wikipedia, Brown Dwarfs are sub-stellar objects whose masses range from 13 to 80 Jupiter masses ( $M_J$ ) [Wikipedia, Brown Dwarfs].

In our opinion, Brown Dwarfs (BD) differ from red stars in that the density of their cores is smaller than nuclear density. Consequently, WIMP annihilation does not take place.

As we have shown in Section 2.10, the minimum mass and maximum radius of a compact star made up of weakly interacting WIMPs can be calculated with the following parameters  $A_1$  and  $A_2$ :

$$A_1 = \frac{\pi}{6} \sqrt{6} (\alpha\beta)^2 \tag{2.15.1}$$

$$A_2 = \pi \sqrt[6]{6} (\alpha\beta)^{\frac{2}{3}} \tag{2.15.2}$$

Parameter  $A_1$  in the scaling solution defines the maximum mass of the core  $M_{COREmax}$  made up of warm WIMPs, and consequently the maximum mass of brown dwarf  $M_{BDmax}$ :

$$M_{COREmax} = \sqrt{6}\alpha^4\beta^2 M_0 \cong 4.33 \times 10^{26} \text{ kg} \quad 2.15.3$$

$$M_{BDmax} = \beta^2 M_{COREmax} \cong 7.77 \times 10^{28} \text{ kg} (\cong 41 M_J) \quad 2.15.4$$

Parameter  $A_2$  defines the minimum radius of the BD core  $R_{COREmin}$  made up of warm WIMPs:

$$R_{COREmin} = A_2\alpha^2 \frac{L_g}{\pi} \cong 280 \text{ m} \quad 2.15.5$$

The minimum mass of the BD  $M_{BDmin}$  equals to the minimum star mass:

$$M_{BDmin} = M_{Smin} = \alpha^{-2} M_0 \cong 1.8 \times 10^{28} \text{ kg} (\cong 9.7 M_J) \quad 2.15.6$$

---

## 2.16. PLANETS

---

By Wikipedia definition, *Planet is a celestial body orbiting a star or stellar remnant that is massive enough to be rounded by its own gravity, is not massive enough to cause thermonuclear fusion, and has cleared its neighbouring region of planetesimals.* [Wikipedia, Planet]

Let's see how planetary system formation occurs.

As described above, extrasolar systems arise from clouds of all particles under consideration with mass in the range of

$$M_{Cl} = 4\pi m_0 \times Q^{\frac{3}{2}} \times (\alpha^2 \leftrightarrow 1) \quad 2.16.1$$

and density

$$\rho_{Cl} \sim \rho_{cr} \times Q^{\frac{1}{4}} \quad 2.16.2$$

As a result of gravitational instability, gravitational collapse takes place and one third of  $M_{Cl}$  is concentrating at the center of the cloud, increasing the density of the core up to the nuclear density.

The heaviest particles – neutralinos and WIMPs – are the first in this stream of matter. When their density achieves the nuclear density, self-annihilation process ignites. As the result, the Stellar Core (SC) grows up to  $10^4$  and  $10^2$  times respectively, taking additional mass of neutralinos and WIMPs from oncoming stream.

Concurrently, White Dwarf Shell (WDS) forms around the SC. WDS is comprised of next heaviest particles – protons, accompanied by electrons. The total mass of WDS equals to  $\cong 2.4 \%$  of the cloud mass.

Expansion of the hot SC and WDS is progressing explosively fast, in a process not unlike boiling. Drops of the boiling SC and WDS are ejected from the forming star, and give birth to planets.

The following two facts support the creation picture outlined above:

- The analysis of a mass – radius ratios for compact stars made of strongly interacting fermions shows that the radius remains approximately constant for a wide range of compact stars masses;
- The analysis of a mass – radius ratios for the lowest mass white dwarfs shows the same behavior – radius does not depend on mass. It happens because at the low mass end the Coulomb pressure (which is characterized by constant density  $\propto \frac{M}{R^3}$  and thus  $R \propto M^{\frac{1}{3}}$ ) starts to compensate the degeneracy:  $R \propto M^{-\frac{1}{3}}$ . The two effects nearly cancel each other out, so  $R \propto M^0$  – no dependency at all.

As discussed above, the maximum mass of the hot neutralinos and WIMPs core  $M_{CORE} \cong 1.93 \times 10^{30} \text{ kg}$ ; the maximum mass of a star  $M_{Smax} = 3.4654 \times 10^{32} \text{ kg}$ ; and the minimum radius  $R_{min} \cong 8.6 \text{ km}$ .

The radius of the hot core remains practically constant ( $\cong 8.6 \text{ km}$ ) whether the core belongs to a star or to a planet. The masses of planets formed around red stars and main-sequence stars differ:

- Planets formed around red stars have the smallest mass of  $\sim 10^{-6} M_{Sun}$ , 8 orders of magnitude smaller than maximum star mass  $\cong 174 M_{Sun}$ .
- Planets formed around main-sequence stars may be as light as  $\sim 10^{-15} M_{Sun}$ , 17 orders of magnitude smaller than the maximum star mass. Consequently, all spherically-shaped objects, down to Mimas in Solar system, contain hot neutralinos cores with WDS.

Planets can arise only around main sequence and red stars. Due to the less violent nature of their formation, brown dwarfs do not create planets. There have been observations of a number of BDs possessing planets; with respect to our Model, the masses of such BDs should exceed  $0.039 M_{Sun}$ , which would classify them as red stars.

## 2.17. COSMIC DUST

---

According to Wikipedia, *Cosmic dust was once solely an annoyance to astronomers, as it obscures objects they wish to observe. When infrared astronomy began, those previously annoying dust particles were observed to be significant and vital components of astrophysical processes* [Wikipedia, Cosmic dust].

In our own Solar system, not only does cosmic dust play a major role in the zodiac light, but it also produces thermal emission, which is the most prominent feature of the night-sky light in the  $5 \Leftrightarrow 50$  micrometer wavelength domain.

The grains characterizing the  $2.9 K$  black body infrared radiation have typical sizes of  $10 \Leftrightarrow 100$  micrometers with an average density  $\sim 2.5 \times 10^3 \frac{kg}{m^3}$ .

P. H. Siegel has this to say about cosmic dust:

*Results from the NASA Cosmic Background Explorer (COBE) Diffuse Infrared Background Experiment (DIRBE) and examination of the spectral energy distributions in observable galaxies, indicate that approximately one-half of the total luminosity and 98% of the photons emitted since the Big Bang fall into sub-millimeter and far-IR. Much of this energy is being radiated by cool interstellar dust.*<sup>17</sup>

The sizes of cosmic grains  $D_{gr}$  are roughly equal to the Fermi length  $L_F$ :

$$D_{gr} \sim L_F = a \times Q^{\frac{1}{4}} = 1.6532 \times 10^{-4} m \quad 2.17.1$$

and their mass  $m_{gr}$  is close to the Planck mass:

$$m_{gr} \sim (10^{-9} \Leftrightarrow 10^{-7}) kg$$

$$(M_p = 2.1767 \times 10^{-8} kg) \quad 2.17.2$$

As discussed in Section 3.3, masses of two gravitationally interacting objects  $m_1$  and  $m_2$  must satisfy the following expression:

$$m_1 m_2 \geq \frac{1}{2} M_p^2 = 2m_0^2 \times Q \quad 2.17.3$$

Cosmic grains with masses around  $M_p$  are the smallest building blocks that participate in star creation. Formation of a new star starts with a gravitational instability of the dust cloud and subsequent gravitational collapse, with the resulting macroobject (Nucleus) possessing mass about  $M_{Nuc}$ :

$$M_{Nuc} = m_o \times Q \cong 10^{12} kg \quad 2.17.4$$

---

<sup>17</sup> P. H. Siegel, IEEE Transactions on Microwave Theory and Techniques, Vol. 50, No. 3, 910 (2002).

Then all particles heavier than  $m_0$  (neutralinos, WIMPs, protons, DIRACs) will be attracted to this Nucleus, increasing its mass and attracting lighter particles as described above. The size of this Nucleus is:

$$R_{Nuc} \sim 10^{-2} m \quad 2.17.5$$

A dust particle of mass  $B_1 M_P$  and radius  $B_2 L_F$  is absorbing energy from the Medium at the following rate:

$$\frac{d}{dt}(B_1 M_P c^2) = \frac{B_1 M_P c^2}{2t} \quad 2.17.6$$

where  $B_1$  and  $B_2$  are parameters.

The absorbed energy will increase the particle's temperature, until equilibrium is achieved: power absorption equals to the power irradiated by the surface in accordance with the Stefan-Boltzmann law

$$\frac{B_1 M_P c^2}{2t} = \sigma_{SB} T_{st}^4 \times 4\pi B_2^2 L_F^2 \quad 2.17.7$$

where  $\sigma_{SB}$  is the Stefan-Boltzmann constant and  $k_B$  is the Boltzmann constant:

$$\sigma_{SB} = \frac{2\pi^5 k_B^4}{15h^3 c^3} \quad 2.17.8$$

Applying the World equation 2.11.2 to our particle:

$$B_1 M_P c^2 = 4\pi B_2^2 L_F^2 \sigma_0 \quad 2.17.9$$

we calculate its stationary temperature  $T_{st}$  to be

$$k_B T_{st} = \left(\frac{15}{4\pi^5}\right)^{\frac{1}{4}} \frac{hc}{L_F} \quad 2.17.10$$

$$T_{st} = 28.95 K$$

This result is in an excellent agreement with experimentally measured value of 29 K.

---

## 2.18. WORLD EXPANSION

One of the long-standing questions of Cosmology revolves around the observed expansion of the World. Furthermore, the expansion of the World appears to be accelerating. It is commonly accepted that the mysterious Dark Energy is the agent responsible for this acceleration. In this section we introduce an alternative explanation of this phenomenon.

In our Model, there is no gravitational interaction between microobjects (protons, electrons, DM particles, etc.). At least one of the parties participating in gravitational

interaction must be a macroobject. More precisely, the product of the two objects' masses must equal to at least Planck mass squared (see Section 3.3):

$$m_1 m_2 \geq \frac{1}{2} M_P^2 \quad 2.18.1$$

Consequently, the particles constituting the Medium of the World do not participate in gravitational interaction with each other.

The motion of macroobjects in the Medium is governed by three separate forces.

- Outward force: the force that the Medium applies to macroobjects, pushing the objects away from the World center.
- Gravity: the force that macroobjects exert on each other, pulling them back together.
- Friction: the force due to the motion of macroobjects through the Medium, slowing them down.

The three forces are in equilibrium, and macroobjects maintain constant momentums.

From the preceding chapters recall that the World consists of homogenous Medium with mass density

$$\rho_M = \frac{2\rho_{cr}}{3c^2} \quad 2.18.2$$

and the World's Boundary with the Universe has an antimatter surface density

$$\rho_S = \frac{\sigma_0}{c^2} \quad 2.18.3$$

Let us find the gravitoelectric potential  $U_W$  of the World. For a homogeneously charged ball with radius  $R$ , potential  $U_B$  equals to

$$U_B = \frac{2\pi\rho_M R^2}{4\pi\epsilon_g} \left(1 - \frac{r^2}{3R^2}\right) = \frac{c^2}{2} \left(1 - \frac{r^2}{3R^2}\right) \quad 2.18.4$$

where  $r$  is the distance from the center of the ball.

For a homogeneously charged surface of the ball, potential  $U_S$  inside of the sphere with the radius  $R$  is:

$$U_S = \frac{4\pi R^2 \rho_S}{4\pi\epsilon_g R} = \frac{c^2}{2} \quad 2.18.5$$

The potential of the World is:

$$U_W = c^2 \left(1 - \frac{r^2}{6R^2}\right) \quad 2.18.6$$

The gravitoelectric field  $\mathbf{E}_g$  is:



$$\mathbf{E}_g = \frac{r}{3t^2} \frac{r}{r} = a_g(r) \frac{r}{r} \quad 2.18.7$$

where  $a_g(r) = \frac{r}{3t^2}$  is acceleration at distance  $r$  from the center of the ball.

The further away an object from the center of the World, the higher the acceleration. Its maximum value occurs at the Front where  $r = R$ ,

$$a_g(R) = \frac{c}{3t} \cong 2.2 \times 10^{-10} \frac{m}{s^2} \quad 2.18.8$$

The accelerated movement of macroobjects does not however imply an accelerated expansion of the World. The Front of the World is advancing with constant speed  $c$ !

Since all interactions propagate with finite speed that does not exceed  $c$ , the above effect manifests itself mostly close to the Front, and is negligible in the vicinity of the center of the World.

The idealized Medium of the World considered in the above equations is frictionless. Macroobjects moving through such Medium do not lose momentum, and are indeed accelerating. We ignored the effects of friction, as well as gravity between individual macroobjects.

In the actual Medium, the outward force equals to the sum of gravity and friction. Macroobjects will then move with constant speeds.

Let's calculate the friction coefficient  $k_{fr}$  of the Medium:

$$mE_g - G \frac{m}{r^2} \times \frac{4\pi}{3} r^3 \times \frac{1}{3} \frac{\rho c r}{c^2} = k_{fr} v = k_{fr} \frac{r}{R} c \quad 2.18.9$$

$$k_{fr} = \frac{m}{6t} \quad 2.18.10$$

The friction force  $F_{fr}$  for any object with momentum  $p$  then equals to

$$F_{fr} = \frac{p}{6t} \quad 2.18.11$$

The dependence of the Medium friction coefficient on time  $k_{fr} \propto t^{-1}$  can be easily explained by the dependence of a dynamic viscosity of the Medium  $\eta_M$

$$\eta_M = \rho_M v_M \quad 2.18.12$$

on its density  $\rho_M \propto t^{-1}$ , while kinematic viscosity of the Medium  $v_M$  remains constant:

$$v_M = ac \quad 2.18.13$$

Consequently, macroobjects maintain constant momentum during the expansion of the World.

## 2.19. COSMOLOGICAL REDSHIFT

---

Wikipedia has this to say about cosmological redshift [Redshift]:

*In the early part of the twentieth century, Hubble and others made the first measurements of the galaxies redshifts beyond the Milky Way. They initially interpreted these redshifts as due solely to the Doppler Effect, but later Hubble discovered a rough correlation between the increasing redshifts and the increasing distance of galaxies. Theorists immediately realized that these observations could be explained by different mechanisms for producing redshifts*

Let us analyze the movement of photons as they travel from distant galaxies to Earth in the time-varying Medium.

As we have shown in Section 2.6, energy of photons remains constant in the ideal frictionless Medium. In the actual Medium with a friction coefficient for photons

$$k_{ph} \sim t^{-1} \tag{2.19.1}$$

the equation for the photons momentum  $p_{ph}$  is:

$$\frac{dp_{ph}}{dt} = -\delta \frac{p_{ph}}{t} \tag{2.19.2}$$

where  $\delta$  is a parameter. Solving equation 2.19.2 we obtain

$$p_{ph}t^\delta = const \tag{2.19.3}$$

Consider a photon with initial momentum  $p_{emit}$  emitted at time  $t_{emit}$ . The photon is continuously losing momentum as it moves through the Medium until time  $t_{obsv}$  when it is observed. The observer will measure  $\lambda_{obsv}$ , compare it with well-known wavelength  $\lambda_{emit}$ , and calculate a redshift:

$$z = \frac{\lambda_{obsv} - \lambda_{emit}}{\lambda_{emit}} \tag{2.19.4}$$

By definition,  $\lambda = \frac{h}{p}$ . When  $\delta = 1$  we obtain:

$$p_{obsv}t_{obsv} = p_{emit}t_{emit} \tag{2.19.5}$$

$$1 + z = \frac{\lambda_{obsv}}{\lambda_{emit}} = \frac{p_{emit}}{p_{obsv}} = \frac{t_{obsv}}{t_{emit}} \tag{2.19.6}$$

Recall that  $t_{emit}$  and  $t_{obsv}$  are cosmic times (ages of the World at the moments of emitting and observing), both measured from the Beginning of the World.  $t_{obsv}$  equals to the present age of the World  $A_t$ . If the photon travelled for time  $t_{ph}$ , then

$$t_{obsv} = t_{emit} + t_{ph} \tag{2.19.7}$$

$$t_{ph} = t_{obsv} - t_{emit} = t - t_{emit} \tag{2.19.8}$$

The cosmological redshift is then described by a nonlinear equation on  $t_{ph}$ :

$$1 + z = \frac{1}{1 - \frac{t_{ph}}{t}} \quad 2.19.9$$

As an example, a photon travelling for 7.11 *By* (half of the World's age) will have a redshift of  $1 + z = 2$ . Photon travelling for 12.64 *By* will have a redshift of  $1 + z = 9$ . The difference is due to the dependence of the Medium friction on time: it was 9 times greater at  $t_{emit} = 1.58 \text{ By}$  than it is now at  $t \approx 14.22 \text{ By}$ .

Experimental observations measuring light from distant galaxies and supernovae seem to imply that the World is expanding at an accelerated pace, as is evident from the observed redshift. The time varying friction of the Medium offered above provides an alternative interpretation of these observations.

M. Lopez-Corredoira has this to say about the loss of energy by photons:

*The idea of loss of energy of the photon in the intergalactic medium was first suggested in 1929 by Zwicky. Nernst in 1937 had developed a model which assumed that radiation was being absorbed by luminifereous ether. But there are two problems: 1) all images of distant objects look blurred if the intergalactic space produces scattering; 2) the scattering effect and the consequent loss of energy is frequency dependent.* <sup>18</sup>

Different mechanisms were proposed to avoid blurring and scattering. A paper by M. Lopez-Corredoira provides an excellent review of such mechanisms. <sup>18</sup>

Laio A. *et al* showed that the shift of photon frequency in low density plasma (which is the case in our Model) could come from quantum effects derived from standard quantum electrodynamics. <sup>19</sup>

According to E. J. Lerner, quantum mechanics indicates that a photon gives up a tiny amount of energy as it collides with an electron, but its trajectory does not change. <sup>20</sup>

There is another way to explain the absence of the blurring and scattering. Back in 1839 James McCullagh proposed a theory of rotationally elastic medium, i.e. the medium in which every particle resists absolute rotation. <sup>21</sup> This theory produces equations analogous to Maxwell's electromagnetic equations. In our opinion, the Medium of the World is in fact such a rotationally elastic medium. We propose to review the interaction of photons with the Medium in light of this very unique theory.

---

<sup>18</sup> M. Lopez-Corredoira (2003), astro-ph/0310214v2.

<sup>19</sup> A. Laio, G. Rizzi, and A. Tartaglia, Phys. Rev. E, 55, 7457 (1997).

<sup>20</sup> E. J. Lerner, The Big Bang never happened: a startling refutation of the dominant theory of the origin of the universe, Random House, Toronto, 1991.

<sup>21</sup> J. McCullagh, Transactions of the Royal Irish Academy, 21, 17 (1839).

## 3. PARTICLE PHYSICS

---

### 3.1. ANALYSIS OF MAXWELL'S EQUATIONS

---

*In speaking of the Energy of the field, however, I wish to be understood literally. All energy is the same as energy, whether it exists in the form of motion or in that of elasticity, or in any other form. The energy in electromagnetic phenomena is mechanical energy.*

James Clerk Maxwell

Maxwell's equations, together with the Lorentz force law, form the foundation of classical electrodynamics, classical optics, and electric circuits [Wikipedia, Maxwell's equations]. We'll subsequently refer to Maxwell equations and Lorentz law jointly as Maxwell-Lorentz equations (MLE).

The value of MLE is even greater because J. Swain showed that *linearized general relativity admits a formulation in terms of gravitoelectric and gravitomagnetic fields that closely parallels the description of the electromagnetic field by Maxwell's equation.*<sup>22</sup>

Hans Thirring pointed out this analogy in his "On the formal analogy between the basic electromagnetic equations and Einstein's gravity equations in first approximation" paper published in 1918.<sup>23</sup> It allows us to use formal analogies between the electromagnetism and relativistic gravity.

*The equations for Gravitoelectromagnetism were first published in 1893, before general relativity, by Oliver Heaviside as a separate theory expanding Newton's law* [Wikipedia, Gravitomagnetism].

Maxwell's equations vary with the unit system used. Although the general shape remains the same, various definitions are changed, and different constants appear in different places.

We'll start our discussion with MLE in SI units. We will not rewrite well-known equations, but only provide the relationships between physical quantities used in MLE for electromagnetism and gravitoelectromagnetism in the Tables 3 and 4:

---

<sup>22</sup> J. Swain (2010), arXiv: 1006.575.

<sup>23</sup> H. Thirring, *Physikalische Zeitschrift* **19**, pp. 204-205 (1918).

**Table 3.** Electromagnetism

Charge	Impedance of Electromagnetic Field	Magnetic Flux
$q, C$	$Z_0 = \sqrt{\frac{\mu_0}{\epsilon_0}} = \mu_0 c, \Omega$	$\phi_q, Wb$
Electric Current	Magnetic Parameter	Electric Potential
$I_q, A$	$\mu_0, Hm^{-1}$	$U_q, V$
Magnetic Field Intensity	Electric Parameter	Electric Field
$H_q, Am^{-1}$	$\epsilon_0 = (\mu_0 c^2)^{-1}, \phi m^{-1}$	$E_q, Vm^{-1}$
Electric Flux Density	Electrodynamic Constant	Magnetic Flux Density
$D_q, Cm^{-2}$	$c, ms^{-1}$	$B_q, Wbm^{-2}$

**Table 4.** Gravitoelectromagnetism

Mass	Impedance of Gravitational Field	Gravitomagnetic Flux
$m, kg$	$Z_g = \sqrt{\frac{\mu_g}{\epsilon_g}} = \mu_g c$	$\phi_m, m^2 s^{-1}$
Mass Current	Gravitomagnetic Parameter	Gravitoelectric potential
$I_m, kgs^{-1}$	$\mu_g = \frac{4\pi G}{c^2}$	$U_m, m^2 s^{-2}$
Gravitomagnetic Field Intensity	Gravitoelectric Parameter	Gravitoelectric Field
$H_m, kgm^{-1}s^{-1}$	$\epsilon_g = (\mu_g c^2)^{-1}$	$E_m, ms^{-2}$
Gravitoelectric Flux Density	Gravitoelectrodynamic Constant	Gravitomagnetic Flux Density
$D_m, kgm^{-2}$	$c, ms^{-1}$	$B_m, s^{-1}$

In Maxwell-Lorentz equations, electrodynamic constant  $c$  is defined as the ratio of the absolute electromagnetic unit of charge to the absolute electrostatic unit of charge. It is easy to see that the dimension of products (Charge  $\times$  Magnetic Flux) and (Impedance  $\times$  Charge squared) equals to that of the Plank constant.

From the above Tables it becomes clear that the dimensions of all physical quantities depend on the choice of the charge and mass dimensions (Coulomb & kilogram in SI units). In other unit systems the dimensions are different. For instance, in Gaussian units (CGSE):

- $[q_e] = cm^{\frac{3}{2}}g^{\frac{1}{2}}s^{-1}$
- $[Z_e] = cm^{-1}s$

In CGSM:

- $[q_m] = cm^{\frac{1}{2}}g^{\frac{1}{2}}$
- $[Z_m] = cms^{-1}$

We seem to possess a substantial degree of freedom when it comes to choosing the dimension of charge. For an arbitrary charge transformation parameter  $K$ , we can

- Multiply the charge and mass and all physical quantities on the left side of Tables 3 and 4 by an arbitrary parameter  $K$
- Divide impedances by  $K^2$
- Divide magnetic fluxes and all physical quantities on the right side of Tables 3 and 4 by  $K$ .

Following such a transformation, all physically measurable parameters such as force, energy density, and energy flux density remain the same, and have the same mechanical dimensions.

By definition, 1 Coulomb equals to one tenth of the absolute electromagnetic unit of charge. It follows that in SI we use electromagnetic unit of charge  $e$  in the electrostatic Coulomb law instead of the electrostatic unit  $\frac{e}{c}$ . This seems a bit odd.

Likewise, when describing Newtonian Law of gravitation, we use  $m$  – the inertial mass, instead of gravitoelectrostatic charge  $mc$  – the gravitational mass. The gravitoelectromagnetic charge is then  $mc^2$ . Similarly to the electromagnetic field, the gravitoelectrodynamics constant  $c$  is the ratio of the absolute gravitoelectromagnetic unit of charge to the absolute gravitoelectrostatic unit of charge.

All elementary particles in the World are fully characterized by their four-momentum  $\left(\frac{E}{c}, \mathbf{p}\right)$  that satisfies the following equation:

$$\left(\frac{E}{c}\right)^2 - \mathbf{p}^2 = Inv = (mc)^2 \quad 3.1.1$$

where the invariant is, in fact, the gravitoelectrostatic charge  $mc$  squared, and  $E$  is the gravitoelectromagnetic charge.

The inertial mass and the gravitational mass are not the same physical quantity. Instead, they are proportional to each other, and their ratio equals to the gravitoelectrodynamics constant  $c$ . The classical theory offers no compelling reason why the gravitational mass  $mc$  has to equal the inertial mass  $m$ , commonly referred to as “rest mass.”

Analogous to electromagnetism, we can think of  $m$  as a gravitocapacitor. Then,  $E = mc^2$  describes the accumulation of energy by gravitocapacitor with capacity  $m$ , rather than transformation of energy to mass.

When a gravitoelectrostatic charge of a particle equals to momentum  $p_{DB}$ , gravitomagnetic flux  $\phi_{DB}$  is

$$\phi_{DB} = \frac{h}{p_{DB}} = \lambda_{DB} \quad 3.1.2$$

known as de Broglie wavelength.

The notion of “wavelength” is thus a macroscopic notion, namely, gravitomagnetic flux of particles characterized by four-momentum only.

### 3.2. MAGNETIC MONOPOLE, MAGNETIC DIPOLE

---

*Maxwell's equations of electromagnetism relate the electric and magnetic fields to each other and to the motions of electric charges. The standard equations provide for electric charges, but they posit no magnetic charges [Wikipedia, Magnetic monopole].*

Let's start from the original equations. The Dirac's equation introduces the magnetic monopole:

$$\frac{e\mu}{4\pi\epsilon_0} = n \frac{hc}{4\pi} \quad 3.2.1$$

where  $n$  is an integer, and  $e$  and  $\mu$  are electromagnetic charges. Taking into account the following well-known equation

$$\frac{e^2}{4\pi\epsilon_0} = \frac{\alpha hc}{2\pi} \quad 3.2.2$$

for  $n = 1$  we obtain the minimum magnetic charge  $\mu = \frac{e}{2\alpha}$ .

Impedance of electromagnetic field  $Z_0$  equals to

$$Z_0 = \frac{1}{\epsilon_0 c} = \frac{h}{e\mu} \quad 3.2.3$$

Using the equations for  $Z_0$  and  $\mu$  derived above, we obtain the magnetic parameter  $\mu_0$ :

$$\mu_0 = \frac{h}{e\mu c} \quad 3.2.4$$

It is well-known that the dimension of the magnetic field intensity  $[H_q] = Am^{-1}$ . We can rewrite it in the following way:

$$[H_q] = \frac{cm}{m^2s} = \frac{[d_m]}{m^2s} \quad 3.2.5$$

where  $d_m$  is electromagnetic dipole of the electromagnetic charge  $q$ . It looks like magnetic field intensity  $H_q$  is, in fact, the current density of electromagnetic dipoles  $d_m$ . Using the constitutive relation

$$B_q = \mu_0 H_q \quad 3.2.6$$

we can express the magnetic flux with the following equation:

$$\phi_q = \mu_0 H_q S = \mu_0 I_{d_m} \quad 3.2.7$$

where  $S$  is a magnetic flux area,  $I_{d_m}$  is current, and  $H_q$  is the current density of electromagnetic dipoles  $d_m$ .

Magnetic flux quantum  $\phi_0$  can then be expressed as follows:

$$\phi_0 = \frac{h}{2e} = \mu_0 I_{d_m} = \frac{h}{e\mu c} \frac{\mu c}{2} \quad 3.2.8$$

and the quant of electromagnetic dipole current  $I_{d_m}$  is:

$$I_{d_m} = \frac{\mu c}{2} = \frac{\mu a_0}{\tau_0} \quad 3.2.9$$

where  $\tau_0$  is the atomic time:

$$\tau_0 = \frac{a_0}{c} \quad 3.2.10$$

It means that the magnetic flux  $\phi_q$  is the magnetic current of the electromagnetic dipoles:

$$d_m = \frac{\mu a_0}{2} \quad 3.2.11$$

While the magnetic field intensity  $H_q$  is the current density of electromagnetic dipoles  $d_m$ , the electric flux density  $D_q$  is the current density of electrostatic dipoles

$$d_e = \frac{\mu a_0}{2c} \quad 3.2.12$$

To summarize, electrostatic and electromagnetic monopoles are not the subjects of Maxwell-Lorentz equations; instead, currents and current densities of the electrostatic and electromagnetic dipoles are. We will subsequently refer to electrostatic and electromagnetic dipoles as simply Magnetic Dipoles (MDs). Previously, we have also used the term "DIRAC" to refer to these particles (see Section 2.9.)

So-called "auxiliary" magnetic field intensity and electric flux density are indeed real physical characteristics of the electromagnetic field. They refer to current density of magnetic dipoles.



DIRACs have negligible electrostatic and electromagnetic charges, since the separation between charges is very small ( $\frac{a_0}{2}$ ). They do, however, possess a substantial electromagnetic dipole momentum  $d_{em}$  that equals to half of the Bohr magneton  $\mu_B$ :

$$d_{em} = \frac{\mu_B}{2} = 4.63700484(10) \times 10^{-24} \frac{J}{T} \quad 3.2.13$$

The same conclusion can be derived for ELOPs – magnetic dipoles made of two preons: they have negligible charge and a dipole momentum that equals to the nuclear magneton  $\mu_N$ :

$$\mu_N = \frac{eh}{4\pi m_p} = 5.05078324(13) \times 10^{-27} \frac{J}{T} \quad 3.2.14$$

It is interesting to proceed with the same approach for the gravitoelectromagnetic field. It turns out that:

- The quant of the gravitoelectromagnetic dipole is  $\frac{hc}{4\pi}$  ;
- The quant of the gravitoelectrostatic dipole is  $\frac{h}{4\pi}$  (spin of fermions);
- The gravitomagnetic field intensity is the current density of MDs  $\frac{hc}{4\pi}$ ;
- The gravitoelectric flux density is the current density of the MDs  $\frac{h}{4\pi}$ .

### 3.3. DIRAC'S EQUATION FOR GRAVITOELECTROMAGNETIC FIELD

---

Recall Dirac's magnetic monopole equation:

$$\frac{e\mu}{4\pi\epsilon_0} = n \frac{hc}{4\pi} \quad 3.3.1$$

This equation is known as the Dirac quantization condition. Wikipedia [Magnetic monopole] says: *The hypothetical existence of a magnetic monopole would imply that the electric charge must be quantized in certain units; also, the existence of the electric charges implies that the magnetic charges of the hypothetical magnetic monopoles, if they exist, must be quantized in units inversely proportional to the elementary electric charge.*

The charge of Dirac's monopole equals to

$$\mu = n \frac{\epsilon_0 hc}{e} = n \frac{e}{2\alpha} \quad 3.3.2$$

and the minimum charge ( $n=1$ ) is:

$$\mu = \frac{e}{2\alpha} \quad 3.3.3$$

Taking into account the analogy between electromagnetic and gravitoelectromagnetic fields, we can rewrite the same equation for masses of a gravitoelectromagnetic field:

$$\frac{mM}{4\pi\epsilon_g} = \frac{hc}{2\pi} \frac{mM}{M_p^2} = n \frac{hc}{4\pi} \quad 3.3.4$$

Taking  $n = 1$ , we obtain the minimum product of the masses

$$mM = \frac{1}{2} M_p^2 = 2.36904 \times 10^{-16} \text{ kg}^2 \quad 3.3.5$$

Neutralino is the heaviest particle in our model:  $m_N = 2.3441924 \times 10^{-24} \text{ kg}$ . The smallest mass of a macroobject with which a neutralino would interact  $M_{min}$  equals to

$$M_{min} = \frac{1}{2} \frac{M_p^2}{m_N} = 1.0106 \times 10^8 \text{ kg} \quad 3.3.6$$

$M_{min}$  is then the smallest mass of a macroobject produced by a Medium fluctuation that could initiate a gravitational collapse of particles.

Two smaller objects whose masses are close to  $M_p$  could initiate a gravitation collapse as well. Cosmic dust particles discussed above could be playing a significant role in this process.

Let's calculate the magnitudes of fluctuations required to produce macroobjects possessing the minimum mass  $M_{min}$ .

For galaxies with an average density

$$\rho_G \propto \rho_{cr} \times Q^{\frac{1}{8}} \quad 3.3.7$$

the minimum size of a fluctuation is

$$R_{Gmin} \cong 3 \times 10^9 \text{ m} \quad 3.3.8$$

which is much smaller than the size of a galaxy ( $\sim 10^{21} \text{ m}$ ).

For extra-solar systems with an average density

$$\rho_{ESS} \propto \rho_{cr} \times Q^{\frac{1}{4}} \quad 3.3.9$$

the minimum size of a fluctuation is

$$R_{ESSmin} \cong 6.6 \times 10^7 \text{ m} \quad 3.3.10$$

which is much smaller than the size of extra-solar system ( $\sim 10^{16} \text{ m}$ ).  $R_{ESSmin}$  is about 10 times smaller than the radius of the Sun  $\cong 7 \times 10^8 \text{ m}$ .

To produce a dust particle with mass  $m_{dmin}$

$$m_{dmin} = M_p = 2.1767 \times 10^{-8} \text{ kg} \quad 3.3.11$$

the minimum size of a fluctuation in galaxies is:

$$r_{Gmin} \cong 1.8 \times 10^4 \text{ m} \quad 3.3.12$$

and in ESS

$$r_{ESSmin} \cong 4 \times 10^2 \text{ m} \quad 3.3.13$$

These calculations are true for present conditions of the Medium, but the properties of the Medium are changing with time. Let's consider how the first macroobjects arose.

At the very Beginning when the radius of the World was equal to  $a$  and the density  $\rho_{cr0}$  was equal to

$$\rho_{cr0} = 3\rho_0 \quad 3.3.14$$

the total energy inside of the Core of the World was equal to

$$E_{W0} = 4\pi E_0 \quad 3.3.15$$

which is sufficient to produce DIRACs and lighter particles only. The conditions for generating the first macroobjects (which are in fact particles) actualized when the size of the World was about the Compton length of a preon. The total energy at that time was equal to:

$$E_W \left( Q = \frac{3}{\alpha} \right) = \frac{36\pi E_0}{\alpha^2} \quad 3.3.16$$

and was sufficient to generate the very first ensemble of particles.

---

### 3.4. ELEMENTARY CHARGE

---

*Conceptually, Maxwell's equations describe how electric charges and electric currents act as sources for the electric and magnetic fields. Further, it describes how a time varying electric field generates a time varying magnetic field and vice versa [Wikipedia, Maxwell's equations].*

Maxwell's equations produce only two physically measurable quantities: energy density  $\rho_E$ , and energy flux density  $\mathbf{j}_E$ . Other notions – electrical field, magnetic field, etc. – describe the behavior of the Medium, and are only used to calculate  $\rho_E$  and  $\mathbf{j}_E$ .

Maxwell's equations then take densities  $\rho_q$  and current densities  $\mathbf{j}_q$  of electric charges as inputs, and calculate the energy density and energy flux density of the electromagnetic field. While the dimensions of charges are our choice, the dimensions of the output

characteristics of the electromagnetic field are solid – energy density and energy flux density.

Based on these speculations, it seems reasonable to use Energy for the dimension of the electromagnetic charge, and Momentum for electrostatic charge. Maxwell's equations will then use the characteristics of an electromagnetic field having the same dimensions.

With this generalization of MLE, the notion of charge takes on a new physical meaning. We will use the electron rest energy  $E_e$  as the unit of electromagnetic charge. Positron possesses the same amount of energy. Electron-positron annihilation is simply a release of their combined energy.

There is of course a difference between particle and anti-particle electromagnetic quants of energy. One way to explain this difference is the resonance effect of the ponderomotive forces between two pulsating spheres immersed in the Medium of the World.

Lord Kelvin and C.A. Bjerknes investigated this mechanism between 1870 and 1910. Bjerknes showed that when two spheres immersed in an incompressible fluid were pulsated, they exerted a mutual attraction which obeyed Newton's inverse square law if the pulsations are in phase. The spheres repelled when the phases differed by a half wave.<sup>24</sup>

We apply this 140 years old mechanism to electric charges interaction. Recall the first line of Maxwell's equations for electromagnetic field:

$$e = \frac{2\alpha h}{e^2} = \frac{h}{2e} \quad 3.4.1$$

Using the flexibility of the electromagnetic charge dimension we replace  $e$  with  $e_g = 4\pi\left(\frac{L_F}{2\pi}\right)^2$ . Magnetic parameter  $\mu_0$

$$\mu_0 = \frac{2\alpha h}{ce^2} \quad 3.4.2$$

transforms into  $\mu_{0g}$ :

$$\mu_{0g} = \frac{2\alpha h}{ce_g^2} = \frac{2\pi^2\alpha}{3} \frac{\rho_{cr}}{c^2} = \frac{\rho_p}{c^2} \quad 3.4.3$$

$\mu_{0g}$  precisely equals to the value of proton mass density in the Medium of the World (see equation 2.5.5).

It follows that we can treat the electromagnetic field with constant magnetic parameter  $\mu_0$  in the time varying gravitational Medium with the magnetic parameter  $\mu_{0g}$  (which is the Medium's partial proton mass density proportional to  $t^{-1}$ ) and the time varying electric charge  $e_g$  proportional to  $t^{\frac{1}{2}}$  as a sphere with the radius  $\frac{L_F}{2\pi}$  proportional to  $t^{\frac{1}{4}}$ .

---

<sup>24</sup> V. F. K. Bjerknes, Fields of Force, Columbia Press (1906).

Of course, we can return to the equation describing the electromagnetic field with the constant magnetic parameter using  $\frac{L_F}{2\pi} \rightarrow r_e$  transformation:

$$4\pi r_e^2 \frac{2\alpha h}{c(4\pi r_e^2)^2} = \frac{h}{8\pi r_e^2} \quad 3.4.4$$

The electron can then be viewed as a pulsating sphere with radius  $r_e$ .

Energy irradiated by the pulsating spheres is compensated by the energy flux through the closed surfaces of objects from the surrounding Medium of the World.

In our Model, energy of an object equals to the area of a closed surface multiplied by the surface enthalpy  $\sigma_0$ :

$$E_e = 4\pi r_e^2 \sigma_0 \quad 3.4.5$$

We use 3.4.5 to calculate the radius of an electron:

$$r_e = (\pi\alpha)^{\frac{1}{2}} a_0 = 0.15141105 a_0 \quad 3.4.6$$

Electron radius  $r_e$  is about 6.6 times smaller than the classical electron radius  $a_0$ .

It is interesting to proceed with the same approach for the gravitoelectromagnetic field. The first line of Maxwell's equations for gravitoelectromagnetism is:

$$m \frac{4\pi G}{c^2} c \phi_m \quad 3.4.7$$

Using the flexibility of the gravitoelectromagnetic charge dimension we replace  $m$  with  $m_g = \frac{a^3}{2L_{Cm}}$ , where  $L_{Cm}$  is a Compton length of mass  $m$ . Gravitomagnetic parameter  $\mu_g$

$$\mu_g = \frac{4\pi G}{c^2} \quad 3.4.8$$

transforms into  $\mu_{gg}$ :

$$\mu_{gg} = \frac{2}{3} \frac{\rho_{cr}}{c^2} = \frac{\rho_M}{c^2} \quad 3.4.9$$

$\mu_{gg}$  precisely equals to the value of the Medium energy density  $\rho_M$  over  $c^2$ . The impedance of gravitational field  $Z_{gg} = \mu_{gg} c$  is the energy current density of the Medium over  $c^2$ . These conclusions emphasize the physical meaning and significance of energy density  $\rho_{cr}$  as one of the main characteristic of the World.

Gravitomagnetic Flux  $\phi_m$  transforms to mass current  $I_m = \frac{I_E}{c^2}$ , where  $I_E$  is energy current, and gravitomagnetic flux density  $B_m$  becomes energy current density over  $c^2$ .

The same approach can be used for all particles in the Medium: protons, electrons, photons, neutrinos, and dark matter particles, whose energy densities were discussed in Sections 2.7, 2.9.

To summarize, electromagnetic and gravitoelectromagnetic charge can be expressed as Energy, or alternatively as Area multiplied by  $\sigma_0$ .

### 3.5. MODEL OF AN ELECTRON


---

The main idea of quantization of an electron electric charge  $e$  is connected to the existence of a magnetic monopole with an electric charge  $\mu$  which is located close enough to the electron. We transform this idea, and propose the existence of a magnetic dipole close to the electron.

A model of an electron in a semi-classical approach can then be pictured as follows: pulsating sphere with radius  $r_e$  and an electromagnetic energy  $E_e$  is rotating around a stationary magnetic dipole axis at a distance  $\frac{Lce}{4\pi}$  from the axis and  $\frac{Lce}{4\pi \sin \theta}$  from the dipole, where  $\theta$  is the angle between the dipole axis and the direction to the rotating electron that satisfies the following equation:

$$\cos \theta = \pm \sqrt{\frac{1}{3}} \tag{3.5.1}$$

The different signs ( $\pm$ ) correspond to an electron (+) and a positron (-) rotating on the different sides of the stationary magnetic dipole. It is a well-known result of electromagnetic theory for stable rotation of the point charge around the stationary point dipole.



In this “Cone” model of an electron, the orbital angular momentum  $L$  equals to spin  $s = \frac{h}{4\pi}$ , and the orbital magnetic dipole momentum equals to  $\frac{\mu_B}{2}$ . The dipole momentum

of the magnetic dipole also equals to  $\frac{\mu_B}{2}$  (3.2.13). The total magnetic momentum then equals to  $\mu_B$  – the Bohr magneton.

The classical model introduces  $g$ -factor with a value of 2 to explain the magnetic momentum of an electron. The Cone model avoids introduction of an additional arbitrary parameter.

Electrical potential  $U_d$  of a magnetic dipole is

$$U_d = \frac{\mu_B}{2} \frac{\cos \theta}{4\pi\epsilon_0 r^2} \quad 3.5.2$$

and interaction energy  $E_{ed}$  between the electron and the magnetic dipole is

$$E_{ed} = eU_d \quad 3.5.3$$

Taking  $b$  for the radius of an electron rotation, we obtain

$$E_{ed} = \frac{a \cos \theta (\sin \theta)^2}{16\pi^2 b^2} hc \quad 3.5.4$$

We proceed to calculate the rotation radius  $b$  of an electron with spin  $\frac{h}{4\pi}$  rotating with momentum  $m_e v$ :

$$b = \frac{h}{4\pi m_e v} \quad 3.5.5$$

and

$$E_{ed} = \alpha E_e \cos \theta (\sin \theta)^2 \frac{v^2}{c^2} \quad 3.5.6$$

When  $v = c$ :

$$E'_{ed} = \frac{2\sqrt{3}}{9} \alpha E_e \cong 1.4353 \text{ keV} \quad 3.5.7$$

When  $v = \alpha c$  (as it is in an atom):

$$E''_{ed} = \frac{2\sqrt{3}}{9} \alpha^3 E_e \cong 0.07643 \text{ eV} \cong 887 \text{ K} \quad 3.5.8$$

Note that according to 3.5.8, the electron binding energy is quite low, and the energy required for removal of an electron from a magnetic dipole can easily be supplied by an application of an electric field or temperature.

The above “Cone” model is rough, as it does not take the mass of the dipole  $m_0$  into account. Likewise, we have simplistically assumed that the dipole is stationary. It would be interesting to develop this model into a full-fledged theory, as the concept of “Cone” construction of charges around magnetic dipoles may come useful in other applications as well.

Magnetic Dipole – DIRAC – is a unique creature of the World. A DIRAC is comprised of two monopoles. The monopoles possess electrical charges of  $\mu^+$  and  $\mu^-$ , and masses of  $\frac{m_0}{2}$ . The charge-to-mass ratio of a monopole equals to that of an electron. DIRACs have a spin of 0, and consequently they are bosons.

Fermions  $\mu^+$  and  $\mu^-$  with masses  $\frac{m_0}{2}$  are the smallest building blocks of a cluster structure of protons, WIMPs, neutralinos, constituent masses of Up and Down quarks, mesons, pions, etc.

More than 60 years ago, Y. Nambu proposed an empirical mass spectrum of elementary particles with a mass unit close to one quarter of the mass of a pion (about  $\frac{m_0}{2} \cong 35 \text{ MeV}/c^2$ ).<sup>25</sup> He noticed that meson masses are even multiplies of a mass unit  $\frac{m_0}{2}$ , baryon (and also unstable lepton) masses are odd multiplies, and mass differences among similar particles are quantized by  $m_0 \cong 70 \text{ MeV}/c^2$ . During the last 40 years M. H. Mac Gregor studied this property extensively.<sup>26</sup>

---

### 3.6. FRACTAL STRUCTURE OF PARTICLES

---

*We are all agreed that your theory is crazy. The question that divides us is whether it is crazy enough to have a chance of being correct.*

Niels Bohr

In our Model, the masses of all particles of dark matter are proportional to basic mass  $m_0$ , and the coefficient of the proportionality is the fine-structure constant raised to different exponents. Each particle is  $\frac{1}{\alpha}$  times heavier than the previous one, and in our Model, is indeed built out of lighter particles. We will now take a closer look at this phenomenon.

In order for a larger particle to be  $\frac{1}{\alpha} \approx 137$  times heavier than the lighter one, it has to be composed of more than 137 lighter particles, taking binding energy into account. There is then nothing surprising about the value of the Fine Structure Constant not being an integer value.

The number of constituent particles is not presently known, but we do know that it must satisfy two criteria: it must be odd, since all particles under consideration are fermions, and

---

<sup>25</sup> Y. Nambu, Prog. Theor. Phys. 7, 131 (1952).

<sup>26</sup> M. H. Mac Gregor. The Power of Alpha, World Scientific, Singapore, 2007.



it must be divisible by 3, since in our Model, heavier particles often consists of 3 clusters of lighter ones. The smallest number to satisfy these conditions is  $141 = 3 \times 47$ , and we will use it in subsequent equations as an example; but keep in mind that  $147 = 3 \times 49$ ,  $153, 159$  etc. would fit our Model just as well.

In our Model, sterile neutrino is the lightest particle of Dark Matter. An electron is  $\frac{1}{\alpha}$  heavier and consists of 3 preons of 47 sterile neutrinos each. An ELOP is a dipole made up of two preons. Other Dark Matter particles are organized in a similar fashion.

Dirac's monopoles with mass  $\frac{m_0}{2} \cong 35 \text{ MeV}/c^2$  are the basic building blocks of fractal cluster structures of different hadron particles. Protons, WIMPs, neutralinos, mesons, pions, constituent masses of Up and Down quarks are all built from Dirac's monopoles. Additionally, Dirac's monopoles are the agent responsible for the strong nuclear interaction.

All "elementary" particles of the World are fermions and they possess masses. Bosons such as photons, X-rays, and gamma rays are composite particles and consist of an even numbers of fermions.

An axion is the boson possessing the lowest mass  $m_a$  (see Section 2.6). It consists of two interacting neutrinos (one of the possible super-weak interactions, see Section 3.7), for example electron and muon neutrinos:

$$m_a = \left(\frac{\alpha}{\beta}\right)^{\frac{1}{2}} m_0 \times Q^{-\frac{1}{2}} < \frac{m_{\nu_e} m_{\nu_\mu}}{m_0} = \frac{1}{24} m_0 \times Q^{-\frac{1}{2}} \quad 3.6.1$$

Gamma rays are usually distinguished from X-rays by their origin: *X-rays are emitted by electrons outside the nucleus, while gamma rays are emitted by the nucleus* [Wikipedia, Gamma ray]. A better way to distinguish the two, in our opinion, is the type of fermion composing the core of X-quant and gamma-quant.

Super soft X-rays [Wikipedia, Super soft X-ray source] have energies in the 0.09 to 2.5 keV range, whereas soft gamma rays have energies in the 10 to 5000 keV range. We assume that X-quant are composed of two interacting muonic and tauonic neutrinos and have rest mass of  $m_X \sim m_0 \times Q^{-\frac{1}{2}}$  which is decreasing with time similarly to an axion (3.6.1):  $m_X \propto t^{-\frac{1}{2}}$ .

Soft gamma-quant are composed of two sterile neutrinos (3.7 keV each). Hard and super-hard gamma-quant may be composed of two preons (0.17 MeV each), which are ELOPs in our Model, two Dirac's monopoles (35 MeV each) which are, in fact, DIRACs, two WIMPs (9.6 GeV each) or two neutralinos (1.3 TeV each). Rest masses of gamma-quant remain constant with time.

As a result of electron-positron annihilation in the low energy case, two or three gamma ray photons are created. At the first step the interaction of electron and positron stimulates

decomposition of their cluster structures into three preons each. The six preons combine into 3 ELOPs, which form the cores of gamma ray photons.

The reverse reaction, electron-positron creation, is a form of pair production from three ELOPs in the Medium of the World induced, for example, *by a super-strong intrinsic magnetic field and a circularly polarized electromagnetic wave propagating along the magnetic field lines in the magnetized pair plasma near the polar caps of the pulsar.* <sup>27</sup>

### 3.7. GRAND UNIFIED THEORY

---

*The Grand Unified Theory is a model in particle physics in which at high energy, the three gauge interactions of the Standard Model which define Weak, Electromagnetic, and Strong interactions, are merged into one Single interaction characterized by one Larger gauge symmetry and thus one Unified Coupling constant [Wikipedia, Grand Unified Theory].*

By definition: *a Coupling constant is a number that determines the strength of an interaction. Usually the Lagrangian or the Hamiltonian of a system can be separated into a kinetic part and an interaction part. The Coupling constant determines the strength of the interaction part with respect to the kinetic part, or between two sectors of the interaction part [Wikipedia, Coupling constant].*

For example, the gravitational coupling parameter  $\alpha_G$  can be defined as follows:

$$\alpha_G = \frac{2\pi G m_e^2}{hc} = \left(\frac{m_e}{M_P}\right)^2 \quad 3.7.1$$

and the electromagnetic coupling constant  $\alpha_{EM}$  as:

$$\alpha_{EM} = \frac{e^2}{2\varepsilon_0 hc} = \alpha \quad 3.7.2$$

$\alpha$  determines the strength of the electromagnetic force of electrons.

At an atomic scale, the strong interaction is about 100 times stronger than electromagnetic interaction, which in turn is about  $10^{10}$  times stronger than the weak force, and about  $10^{40}$  times stronger than the gravitational force, when forces are compared between particles interacting in more than one way.

All these definitions are based on **strength** of the force between a particular pair of particles, and depend on the choice of such particles. Clearly, the gravity between a pair of electrons will differ from that of a pair of protons.

---

<sup>27</sup> Y. Luo and P. Ji, Mon. Not. R. Astron. Soc. **420**, 1673 (2012).

A different way of comparing interactions is looking at their cross-sections. According to Wikipedia, *the concept of a Cross-section is used to express the likelihood of an interaction between particles* [Wikipedia, Cross section (physics)]. In our opinion, all fundamental interactions of the World should be described and compared using their **cross-sections**, which are the measure of the interaction likelihood, and don't depend on the choice of particles. The larger the cross-section, the faster an interaction occurs.

For example, an electromagnetically decaying neutral pion has a life of about  $10^{-16}$  seconds; a weakly decaying charged pion lives for about  $10^{-8}$  seconds, and a free neutron lives for about 15 minutes, making it the unstable subatomic particle with the longest known mean life.

Let's start with the gravitational interaction which is expressed by gravitational parameter  $G$ :

$$G = \frac{1}{4\pi\epsilon_g} \tag{3.7.3}$$

Recall from Section 3.1 that we are free to choose an arbitrary charge transformation parameter  $K$  without affecting the outcome of force, energy density, and energy flux density calculations.

Let's choose  $K = \frac{c}{h}$  and express mass  $m$  of an object in terms of Compton length  $L_{cm}$  by multiplying  $m$  by  $K$ :

$$mK = m \frac{c}{h} = \frac{1}{L_{cm}} \tag{3.7.4}$$

and divide the interaction parameter  $G = \frac{1}{4\pi\epsilon_g}$  by the same coefficient  $K$  squared:

$$G \left(\frac{h}{c}\right)^2 = P \times Q^{-1} \tag{3.7.5}$$

where parameter  $P = \frac{a^2 hc}{8\pi}$ .

By dividing the left side of 3.7.5 by  $P$  we obtain the dimensionless gravitational coupling parameter  $\alpha_G$ :

$$\alpha_G = Q^{-1} \tag{3.7.6}$$

Note that following this transformation, the dimension of the gravitoelectric field is "Energy," and the dimension of the gravitomagnetic flux density is "Momentum". Then the well-known fundamental invariant of the electromagnetic field

$$\left(\frac{E_q}{c}\right)^2 - \mathbf{B}_q^2 = Inv \tag{3.7.7}$$

transforms into the fundamental invariant of the gravitoelectromagnetic field:

$$\left(\frac{E_m}{c}\right)^2 - \mathbf{B}_m^2 = Inv \quad 3.7.8$$

and coincides with the fundamental invariant of the particles:

$$\left(\frac{E}{c}\right)^2 - \mathbf{p}^2 = Inv = (mc)^2 \quad 3.7.9$$

Let's use the same approach for electromagnetic interaction: divide the charge  $e$  by the magnetic dipole  $\frac{\mu\alpha_0}{2}$ :

$$\frac{e}{\mu\alpha_0/2} = \frac{8\pi\alpha}{a} \quad 3.7.10$$

and multiply the interaction parameter  $\frac{1}{4\pi\epsilon_0}$  by the magnetic dipole squared:

$$\frac{1}{4\pi\epsilon_0} \left(\frac{\mu\alpha_0}{2}\right)^2 = \frac{1}{16\pi^2\alpha} P \quad 3.7.11$$

We calculate the dimensionless electromagnetic coupling parameter  $\alpha_{EM}$ :

$$\alpha_{EM} = (16\pi^2\alpha)^{-1} \cong 0.8678 \quad 3.7.12$$

Neglecting the difference between  $\alpha_{EM}$  and value of 1, we can see that the ratio of the coupling parameters is

$$\frac{\alpha_G}{\alpha_{EM}} = Q^{-1} = 1.3156 \times 10^{-40} \quad 3.7.13$$

because the cross-section of the gravitational interaction is  $Q$  times smaller than the cross-section of the electromagnetic interaction.

*In particle physics, Fermi's interaction also known as Fermi coupling is an old explanation of the weak force, proposed by Enrico Fermi, in which four fermions directly interact with one another at one vertex. For example, this interaction explains beta decay of a neutron by direct coupling of a neutron with an electron, antineutrino, and a proton. The interaction could also explain muon decay via a coupling of a muon, electron-antineutrino, a muon- antineutrino and electron. The Feynman diagrams describe the interaction remarkably well [Wikipedia, Fermi's interaction].*

The strength of Fermi's interaction is given by the Fermi's coupling parameter  $G_F$ :

$$\frac{8\pi^3 G_F}{(hc)^3} = \frac{\sqrt{2} g^2}{8 M_W^2} = 1.16637 \times 10^{-5} GeV^{-2} \quad 3.7.14$$

Here  $g$  is the coupling parameter of the weak interaction, and  $M_W$  is the mass of the  $W$  boson.

In our Model, the following four fermions take part in beta decay of a neutron: a proton with mass  $m_p$ , an electron with mass  $m_e$ , a monopole with mass  $\frac{m_0}{2}$ , and an electron

antineutrino with mass  $\frac{m_F}{24}$ , as follows from the following expression for the Fermi coupling parameter:

$$\begin{aligned} \frac{8\pi^3 G_F}{(hc)^3} &= 16\pi^3 \frac{\sqrt{2} m_p \frac{m_F}{24}}{8 m_e \frac{m_0}{2}} \frac{P}{(hc)^3} = \\ &= 8\pi^3 \frac{\sqrt{2}}{8} \frac{\beta}{6\alpha} \frac{P}{(hc)^3} \times Q^{-\frac{1}{4}} = 1.16621 \times 10^{-5} \text{ GeV}^{-2} \end{aligned} \quad 3.7.15$$

which is very close to the presently adopted value of  $1.16637 \times 10^{-5} \text{ GeV}^{-2}$ .

The accuracy of the above calculations depends on the accuracy of measurement of the gravitational parameter  $G$ , that is not better than  $10^{-4}$  at present time.

The equality of  $G_{Fth}$  and  $G_{Fexp}$  yields an electron antineutrinos mass of  $\frac{m_F}{24}$ , that corresponds to its contribution to the energy density of the World's Medium (see Section 2.7).

We find the cross-section of the weak interaction by multiplying  $\frac{G_F}{(hc)^3}$  by  $(hc)^3 \frac{8}{\sqrt{2}} \frac{6\alpha}{\beta}$ :

$$\frac{8}{\sqrt{2}} \frac{6\alpha}{\beta} G_F (hc)^3 = G_F^* = P \times Q^{-1/4} \quad 3.7.16$$

with the dimensionless weak interaction coupling parameter  $\alpha_W$ :

$$\alpha_W = Q^{-1/4} \quad 3.7.17$$

and the ratio of  $\alpha_W$  to  $\alpha_{EM}$

$$\frac{\alpha_W}{\alpha_{EM}} = Q^{-1/4} = 1.0710 \times 10^{-10} \quad 3.7.18$$

As for the strong interaction, the dimensionless coupling parameter  $\alpha_S$  equals to the coupling parameter of the electromagnetic interaction  $\alpha_{EM}$ :

$$\alpha_S = \alpha_{EM} = 1 \quad 3.7.19$$

The difference in the strong and the electromagnetic interactions is not in the coupling parameters but in the **strength** of these interactions depending on the particles involved: electrons with charge  $e$  and monopoles with charge  $\mu$  in electromagnetic and strong interactions respectively.

At the very Beginning ( $Q = 1$ ) all extrapolated fundamental interactions of the World had the same cross-section  $\frac{\alpha^2}{4}$  and were characterized by the Unified coupling constant:

$$\alpha_U = \alpha_S = \alpha_{EM} = \alpha_W = \alpha_G = 1 \quad 3.7.20$$

At that time, the energy density of the World  $\rho_{cr0}$  and the equivalent mass density  $\frac{\rho_{cr0}}{c^2}$  were:

$$\rho_{cr0} = \frac{3hc}{a^4} = 6.0640 \times 10^{30} \frac{J}{m^3} \quad 3.7.21$$

$$\frac{\rho_{cr0}}{c^2} = \frac{3m_0}{a^3} = 6.7470 \times 10^{13} \frac{kg}{m^3} \quad 3.7.22$$

Note that the mass density at the Beginning is much smaller than the nuclear density  $\sim 10^{17} \frac{kg}{m^3}$ . The average energy and mass density of the World has since been decreasing, and their present values are given by

$$\rho_{cr} = \rho_{cr0} \times Q^{-1} = 7.9775 \times 10^{-10} \frac{J}{m^3} \quad 3.7.23$$

$$\frac{\rho_{cr}}{c^2} = \frac{\rho_{cr0}}{c^2} \times Q^{-1} = 8.8760 \times 10^{-27} \frac{kg}{m^3} \quad 3.7.24$$

The gravitational coupling parameter  $\alpha_G$  is similarly decreasing:

$$\alpha_G = Q^{-1} \propto t^{-1} \quad 3.7.25$$

The weak coupling parameter is decreasing as follows:

$$\alpha_W = Q^{-\frac{1}{4}} \propto t^{-\frac{1}{4}} \quad 3.7.26$$

The strong and electromagnetic parameters remain constant in time:

$$\alpha_S = \alpha_{EM} = 1 \quad 3.7.27$$

Our Model predicts two more types of interactions:

- Super weak with the coupling parameter  $\alpha_{SW}$ :

$$\alpha_{SW} = Q^{-\frac{1}{2}} \quad 3.7.28$$

- Extremely weak with the coupling parameter  $\alpha_{EW}$ :

$$\alpha_{EW} = Q^{-\frac{3}{4}} \quad 3.7.29$$

The proposed new interactions may be revealed in the near future. In this light, we should take another look at interactions of neutrinos with matter, photons with dark matter, photons with neutrinos, dark matter with matter, neutrino oscillations, etc.

## 4. THE WORLD

---

*The constitution of the ether, if it ever would be discovered, will be found to be quite different from any thing that we are in the habit of conceiving, though at the same time very simple and very beautiful. An elastic medium composed of points acting on each other in the way supposed by Poisson and others will not answer.*

James McCullagh

The World – Universe Model is developed around two fundamental parameters: Fine-Structure Constant  $\alpha$  and dimensionless quantity  $Q$ . While  $\alpha$  is a constant,  $Q$  increases with time, and in fact defines the size and the age of the World.

The Model is based on Maxwell's equations for the electromagnetism and gravitoelectromagnetism which contain a single constant – electrodynamic constant  $c$ ; two parameters of the Medium – magnetic parameter  $\mu_0$  and gravitomagnetic parameter  $\mu_g$ ; and two measurable characteristics: energy density and energy flux density. All other notions are used for calculations of these two measurable characteristics.

Throughout our discussion we have paid close attention to energy density. Sometimes we used the notion of mass density to facilitate understanding of the Model and correlations of its results with the existent theories and models, but the two concepts were shown to be interchangeable.

For all particles under consideration we used four-momentum, but the final result of the statistical analysis was energy density.

The Fundamental quantities of the World are as follows:

- The surface enthalpy of the World – Universe Front  $\sigma_0$ ;
- The electrodynamic and gravitoelectrodynamics constant  $c$ , which is the speed of the World – Universe Front;
- The radius of the World's Core  $a$ ;
- The dimensionless parameter  $Q$ , which is the size and age of the World measured in the basic units  $a$  and  $\frac{a}{c}$  respectively.

All Fundamental constants are connected to the physical characteristics of the World's Core – the Beginning of the World. All physical parameters of the World can then be expressed in terms of these Fundamental quantities:

$H_0 = \frac{c}{a} \times Q^{-1}$	Hubble parameter
$A_t = \frac{a}{c} \times Q$	Age of the World
$R = a \times Q$	Size of the World
$\rho_0 = \frac{\sigma_0}{a}$	Basic energy density
$\rho_{cr} = \frac{3\sigma_0}{a} \times Q^{-1}$	Critical energy density
$E_0 = \sigma_0 a^2$	Basic energy
$E_W = 4\pi\sigma_0 a^2 \times Q^2$	Energy of the World
$h = \frac{\sigma_0 a^3}{c}$	Plank constant
$G = \frac{c^4}{8\pi\sigma_0 a} \times Q^{-1}$	Gravitational parameter
$\frac{8\pi G}{c^4} = \frac{1}{\sigma_0 a} \times Q^{-1}$	Einstein's parameter

An alternative set of basic parameters fully describes the World as well:

- The Hubble parameter  $H_0$ ;
- The basic energy  $E_0$ ;
- The basic energy density  $\rho_0$ ;
- The dimensionless parameter  $Q$ .

All physical parameters of the World can be expressed through these basic parameters:

$A_t = H_0^{-1}$	Age of the World
$a = \left(\frac{E_0}{\rho_0}\right)^{\frac{1}{3}}$	Size of the World's Core at the Beginning
$\sigma_0 = (E_0 \rho_0^2)^{\frac{1}{3}}$	Surface enthalpy of the World-Universal Front
$c = H_0 \left(\frac{E_0}{\rho_0}\right)^{\frac{1}{3}} \times Q$	Electrodynamic constant
$R = \left(\frac{E_0}{\rho_0}\right)^{\frac{1}{3}} \times Q$	Size of the World
$\rho_{cr} = 3\rho_0 \times Q^{-1}$	Critical energy density



$$E_W = 4\pi E_0 \times Q^2 \quad \text{Energy of the World}$$

$$h = \frac{E_0}{H_0} \times Q^{-1} \quad \text{Plank constant}$$

$$G = \frac{c^4}{8\pi(\rho_0 E_0^2)^{\frac{1}{3}}} \times Q^{-1} \quad \text{Gravitational parameter}$$

$$\frac{8\pi G}{c^4} = (\rho_0 E_0^2)^{-\frac{1}{3}} \times Q^{-1} \quad \text{Einstein's parameter}$$

In our discussion we have often used well-known physical parameters, keeping in mind that all of them can be expressed through Fundamental quantities of the World.

We have developed the Fractal model of the World that describes the macroobjects possessing energies proportional to the total World's macroobjects energy

$E_{MO} = \frac{1}{3} E_W$  with varying coefficients:

- World: 1
- Galaxy clusters:  $Q^{-\frac{1}{8}}$
- Galaxies:  $Q^{-\frac{1}{4}}$
- Globular clusters:  $Q^{-\frac{3}{8}}$
- Extrasolar systems:  $Q^{-\frac{1}{2}}$

The World consists of the Medium and macroobjects. The World has a preferred frame defined by the Medium. Cosmic Microwave Background radiation (CMB) is a component of the Medium. Based on the analysis of the CMB radiation we can measure the speeds of the Milky Way and the Sun relative to CMB rest frame: 552 and 397 km/s respectively.

The World is not empty; instead, it must be treated as a Medium filled galaxies and stars. The Medium behaves as an ideal liquid with unique properties.

Long time ago it was realized that there are no longitudinal waves in the Medium, and hence the Medium could not be an elastic matter of an ordinary type. In 1839 James McCullagh proposed a theory of a rotationally elastic medium, i.e. a medium in which every particle resists absolute rotation.

The potential energy of deformation in such a medium depends only on the rotation of the volume elements and not on their compression or general distortion. This theory produces equations analogous to Maxwell's electromagnetic equations.

The World – Universe Model is based on Maxwell's equations, and McCullagh's theory is a good fit for description of the Medium.

In any homogeneous and isotropic cosmology including our Model with homogeneous and isotropic Medium, the Hubble parameter  $H_0$  and its inverse, the Hubble age of the World,

and also the Hubble length defined as  $\frac{c}{H_0}$ , are absolute and not relative quantities. The Center-of-mass frame of the Hubble sphere can be regarded as a preferred frame.

In light of the Hubble effect, we apply the following transformation to Maxwell's equations: multiply the left side parameters of the Table 4 by the parameter  $\frac{a^3 c}{2h}$ , divide the impedance of the Medium by the same parameter, and leave the right side parameters of the Table 4 alone:

$$q_g = \frac{a^3}{2L_{cm}} \quad Z_g = \frac{1}{R} \times c = H_0 \quad ac \quad 4.1$$

As a result of this transformation:

- All parameters of the gravitoelectromagnetic field have dimensions of length and time; "mass" dimension has disappeared;
- The gravitoelectromagnetic charge has a dimension of "area", which is equivalent to energy, with coefficient that equals to the surface enthalpy  $\sigma_0$ ;
- The impedance of the Medium equals to the Hubble's parameter for the whole World.

It follows that measuring the value of Hubble's parameter anywhere in the World and taking its inverse value allows us to calculate the absolute age of the World. The Hubble's parameter is then the most important characteristic of the World, as it defines the World's age.

The second important characteristic of the World is the gravitomagnetic parameter  $\mu_g$ :

$$\mu_g = \frac{1}{R} \quad 4.2$$

Taking its inverse value, we can define the absolute radius of the World.

We emphasize that the above two parameters are principally different characteristics of the Medium that are connected through the gravitoelectrodynamical constant  $c$ .

The World - Universe Model is the first unified model of the World that successfully describes all of its primary parameters and their relationships, ranging in scale from cosmological structures to elementary particles. The Model allows for precise calculation of values that were only measured experimentally earlier (age of the World, MBR temperature, etc.), and makes verifiable predictions. While the Model needs significant further elaboration, it can already serve as a basis for a new understanding of the World.

**Acknowledgements:** I am grateful to Felix Lev, my life-long friend, for our frequent discussions of history and philosophy of Physics. Special thanks to my son Ilya Netchitailo who questioned every aspect of the manuscript and helped shape it to its present form.