

Linkage of Classical (3 Dimensional) and QM geometry(2 Dimensional) via Hopf mapping and its implications for relic GW power production

A. Beckwith¹

Chongqing University *Department of Physics: Chongqing University,*
Chongqing , PRC , 400044

E-mail: abeckwith@uh.edu

Abstract. Hopf mapping from 2 dimensions (QM) to 3 dimensions (CM) are examined in terms of a formalism started by Feynman which has linkage to the (CM) equations of motion have linkage to the Serret - Frenet form (for Differential equations). We argue that in doing so we may then link QM representations of qubits to a solved version of the rotating rod problem. Furthermore since a rotating rod has linkage to GW generation, as given by Lightman et. al. it is a way to tie qubits (Quantum information) to GW generation. We then make observations as to what the results mean in terms of QM initial states and the power of GW production from early universe conditions.

1.Introduction

We examine the results of the Hopf mapping from 2 dimensions (QM) to 3 dimensions (CM) in terms of generalizations to a rigid rod rotation [1] which could generate GW (gravitational waves) . This paper was initiated by a question by Stephen Kauffman about black holes. I.e. the reality of black hole singularities. To this end, this formalism was chosen to be the basis of finding a solution to trying to investigate black hole physics. After this formalism is further developed, we will use the two to the three sphere mapping to ask if singularities have a presence in astrophysical problems. To start developing mathematics relevant to that future development, we solve on our own a set of equations (in 3 dimensions) pertinent to a non-symmetric object in rotation in early universe which is a way to generate GW and from there state some caveats as to the power of GW which may ensue. The final conclusion of our document is in linking a quantum qubit form with the power created by/ during GW generation which conceivably could be identified by a suitably designed detector. The document first examines what Feynman did with respect to 2 level QM systems, their generalization to Classical rigid rod rotation, and then we solve the resulting CM equation of motion. Feynman decomposed the solutions in x, y, and z in terms of the 2 level QM system [2], [3] a decomposition which we hold as still relevant and valid, and then, using the case of a uniform magnetic field 'down' the z axis, as a driver to the physical process leading to non-symmetric rigid body rotation. That the body is non-symmetric allows us to approximate the GW power generated as to the conventions outlined by Lightman, Press, Price, and Teukolsky [4] . We then conclude with a description of what our model says about QM generation of states relevant to GWs. In the early universe.

¹ To whom any correspondence should be addressed.

2. Outlining the Feynman development of a classical system from 2 level QM system.

We look at how Feynman [1], [2], [3] linked a 2 dim quantum system to a 3 dimensional rigid rod style classical mechanics system. In doing so, Feynman worked with a quantum system given as

$$i \cdot \frac{d}{dt} \begin{pmatrix} a \\ b \end{pmatrix} = -\frac{1}{2} \cdot \begin{pmatrix} H_z & H_x - i \cdot H_y \\ H_x + i \cdot H_y & -H_z \end{pmatrix} \cdot \begin{pmatrix} a \\ b \end{pmatrix} \quad (1)$$

In doing so via the transformation

$$\begin{aligned} x &= a \cdot b^* + b \cdot a^* \\ y &= i \cdot (a \cdot b^* - b \cdot a^*) \\ z &= a \cdot a^* - b \cdot b^* \end{aligned} \quad (2)$$

$$\frac{d}{dt} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = -\frac{1}{2} \cdot \begin{pmatrix} 0 & H_z & -H_y \\ -H_z & 0 & H_x \\ H_y & -H_x & 0 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad (3)$$

The simplest decomposition of this problem is to set $H_y = H_x = 0$ so then the situation is that we have

$$\begin{pmatrix} a(t) \\ b(t) \end{pmatrix} = \begin{pmatrix} a(0) \cdot \exp(i \cdot t \cdot H_z / 2) \\ b(0) \cdot \exp(-i \cdot t \cdot H_z / 2) \end{pmatrix} \quad (4)$$

And

$$\begin{pmatrix} x(t) \\ y(t) \\ z(t) \end{pmatrix} = \begin{pmatrix} x(0) \cdot \cos(H_z \cdot t) + y(0) \cdot \sin(H_z \cdot t) \\ y(0) \cdot \cos(H_z \cdot t) - x(0) \cdot \sin(H_z \cdot t) \\ z(0) \end{pmatrix} \quad (5)$$

As can be seen by Maggiorie [5] and also Lightman, Press, Price, and Teukolsky [4] since the solution as given by Eq.(5) is for a circular moment of a GW there would be a GW associated with it,

We also will be looking at a more complex three dimensional example of motion which is highly complex non withstanding we go to a more complete version of Eq.(1) to Eq.(3) with only $H_y = 0$. Then we get

$$i \cdot \frac{d}{dt} \begin{pmatrix} a \\ b \end{pmatrix} = -\frac{1}{2} \cdot \begin{pmatrix} H_z & H_x \\ H_x & -H_z \end{pmatrix} \cdot \begin{pmatrix} a \\ b \end{pmatrix} \quad (6)$$

$$\frac{d}{dt} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = -\frac{1}{2} \cdot \begin{pmatrix} 0 & H_z & 0 \\ -H_z & 0 & H_x \\ 0 & -H_x & 0 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad (7)$$

The above two equations have the setting of what is called the Serret – Frenet form and we will solve these two DE equation systems taking the approximation that H_z, H_x are constants in lieu of the first example. The next section solves these two equations with this in mind, leading to a non-symmetric rotation in 3 dimensional space which is needed for GW production.

3. Solving a simplified version of Eq. (6) and Eq.(7) to come up with a non- symmetric rigid body rotation sufficient to obtain GW.

To do this we look at Eq. (7) in such a way as to have

$$\begin{aligned}
 x(t) &= x_0 + x_1 \cdot \sin(H_z \cdot t) \\
 y(t) &= \frac{2}{H_x} \cdot \frac{dz(t)}{dt} = -2x_1 \cdot \frac{H_z}{H_x} \cos(H_z \cdot t) \\
 z(t) &= z_0 - \frac{x_1}{2} \cdot \sin(H_z \cdot t) \\
 \frac{dy(t)}{dt} &= \frac{H_z}{2} \cdot x(t) - \frac{H_x}{2} \cdot z(t)
 \end{aligned} \tag{8}$$

To which we add in the reconciliation of the variables equation result from the last part of Eq.(8), namely

$$2x_1 \frac{H_z}{H_x} \cdot \sin(H_z \cdot t) = \frac{1}{2} \cdot [x_0 + x_1 \cdot \sin(H_z \cdot t)] - \frac{1}{2} \cdot \left(\frac{H_x}{H_z} \right) \cdot \left(z_0 - \frac{x_1}{2} \cdot \sin(H_z \cdot t) \right) \tag{9}$$

Leading to

$$x_0 = z_0 \cdot \left(\frac{H_x}{H_z} \right) + x_1 \cdot \left(4 \cdot \frac{H_z}{H_x} - 1 - \frac{1}{2} \cdot \frac{H_x}{H_z} \right) \cdot \sin(H_z \cdot t) \tag{10}$$

Leading to

$$\begin{aligned}
 x(t) &= z_0 \cdot \frac{H_x}{H_z} + 4x_1 \cdot \left[4 \cdot \frac{H_z}{H_x} - \frac{1}{2} \cdot \frac{H_x}{H_z} \right] \sin(H_z \cdot t) \\
 y(t) &= -2x_1 \cdot \frac{H_z}{H_x} \cos(H_z \cdot t) \\
 z(t) &= z_0 - \frac{x_1}{2} \cdot \sin(H_z \cdot t)
 \end{aligned} \tag{11}$$

Combined once again with Eq.(2), and assuming that the quantity we roughly identify with the “magnetic field H_z ” is parallel to the z axis, so long as z_0 as an initial starting point for the z axis is non zero, then we have fulfilled the requirement for a non-uniform motion of a ‘rigid body’ which if related to the quantities in Eq.(2) and also Maggiore’s criteria for GW from a non-uniform non spherical generation of GW leads to the final part of the GW requirement of non-spherically symmetric motion which lends itself to GW generation. We will then make a comment as to how to link this to GW power using Lightman, Press, Price, and Teukolsky[3] to show how frequency from this example can lead to GW generation.

4. Conditions permitting GW power production using the inputs from Eq. (5)

The idea is that we need to calculate the following , i.e. a moment of inertia for a system, and also the frequency. As to Eq. (5) according to the following, we can come up with a generic Eqn. of motion, namely if we do averaging and set out a general time averaging. Fortunately for us the trig identities naturally vanish.

$$I = \alpha \cdot m \cdot r^2 = \alpha \cdot m \cdot \left\langle x^2(t) + y^2(t) + z^2(t) \right\rangle_{Eq.(5)} = \alpha \cdot m \cdot \left(x^2(0) + y^2(0) + z^2(0) \right) \tag{12}$$

We can, as an approximation use \mathbf{m} above to be the net mass of the assumed geometry and set. $\alpha \approx \frac{1}{12}$ I.e.

Then we look at the power loss according to a ‘rigid rod’ construction for GW power generation[4] ,[5]i.e.

$$\frac{d\varepsilon}{dt} = -\frac{32}{5}G \cdot I^2 \omega^6 \quad (13)$$

Note that we can approximate the frequency in this case as directly proportional to the input frequency of the magnetic field parallel to the z axis, i.e. looking to first approximation at $H_z \sim \omega$ according to the conventions as given by Kholodenko [1] on page 157. This means that up to a point, if one picks representative positions as given by $x^2(0)+y^2(0)+z^2(0)$ with each of these initial positions, squared, and a net mass m . Then we can calculate the net GW (gravitational wave) power loss of this system. We will in the end make a comment as to this Eq.(13) value, for the specified inputs into the equation and the Feynman quidbit (QM) results for while comparing them to what we can infer as to Eq. (4), and its up and down 2 dimensional QM states. I.e. this problem is comparatively easy to calculate. In this case the value of Eq.(13) if we are near the cosmological origin would have a value of about

$$\left\langle \dot{\varepsilon} \sim 10^{45} - 10^{50} \text{ Joule / sec} \right\rangle_{\text{relic-condit}} \Rightarrow \left\langle \dot{\varepsilon} / A \sim 10^2 \text{ Joule / sec} \right\rangle_{\text{Earth}} \quad (14)$$

Next we will look at what happens if we assume the input geometry as given by Eq. (11).

Both of these results will be then compared to as to the simple case of Eq.(4) as due to the first set of inputs into Eq. (13) if the spatial geometry of Eq.(5) is used, and then Eq. (4) will be guessed at if we use the geometry of Eq. (11). I.e. we will guess what Eq. (11) does to Eq. (4) and compare that with what Eq. (11) does to Eq. (4)

5. Conditions permitting GW power production using the inputs from Eq. (11)

This is a mess. I.e. what we have to do is to look at how to calculate the moment of inertia, and then going to Eq. (13), even if we assume the same mass which was used earlier to calculate Eq. (14) above for relic conditions.

To start this, look at, even if $\alpha \approx \frac{1}{12}$

$$I = \alpha \cdot m \cdot r^2 = \alpha \cdot m \cdot \left\langle x^2(t) + y^2(t) + z^2(t) \right\rangle_{\text{Eq.(8)}} \quad (15)$$

The problem starts immediately, in that the parenthesis of Eq. (15) above would have to be a time averaged quantity. I.e. we would be looking at $x^2(t)+y^2(t)+z^2(t)$ **with left over terms in this set analytical expression, should they exist to be time averaged, i.e. if**

$z_0 = 0$,

$$x^2(t) + y^2(t) + z^2(t) = x_1^2 \cdot \left(\frac{H_z}{H_x} \right)^2 \cdot \left[4 + \left(252 + 4 \cdot \left(\frac{H_x}{H_z} \right)^4 - \frac{655}{4} \cdot \left(\frac{H_x}{H_z} \right)^2 \right) \cdot \sin^2(H_z \cdot t) \right]_{\text{Eq.(11)}} \quad (16)$$

The term to be time averaged would be $\langle \sin^2(H_z \cdot t) \rangle_{\text{Time-averaged}} \sim 1/2$. So the above would be approximated by

$$\left\langle x^2(t) + y^2(t) + z^2(t) \right\rangle_{\text{Time-averaged}} = x_1^2 \cdot \left(\frac{H_z}{H_x} \right)^2 \cdot \left[4 + \left(\frac{252}{2} + 2 \cdot \left(\frac{H_x}{H_z} \right)^4 - \frac{655}{8} \cdot \left(\frac{H_x}{H_z} \right)^2 \right) \right]_{\text{Eq.(11)}} \quad (17)$$

Using a ratio, as give of $\left(\frac{H_x}{H_z} \right)^2 \sim 1/2$, the above then becomes approximately

$$\left\langle x^2(t) + y^2(t) + z^2(t) \right\rangle_{\text{Time-averaged}} = x_1^2 \cdot \left[\frac{1441}{16} \right] \sim 91 \cdot x_1^2 \quad (18)$$

Then the magnitude of the GW power would be, per second about 10,000 times bigger.

$$\left\langle \dot{\varepsilon} \sim 10^{49} - 10^{53} \text{ Joule / sec} \right\rangle_{\substack{\text{Eq.(11)} \\ \text{Time-Averaged}}}^{\text{relic-condit}} \Rightarrow \left\langle \dot{\varepsilon} / A \sim 10^6 \text{ Joule / sec} \right\rangle_{\substack{\text{Eq.(11)} \\ \text{Time-Averaged}}}^{\text{Earth}} \quad (19)$$

6. Comparison of Eq. (14) and Eq. (19) results in terms upon solving Eq.(1)

The value for the simple geometry (in terms of simple quibits) to understand working with both Eq. (1) and then Eq.(4) has , if a particle is in a constant magnetic field, then according to [4] it is a special case of working with quibits, according to [1], [5] , the values of $\begin{pmatrix} a \\ b \end{pmatrix}$ if only H_z is non zero, for the below equation

become very simple. The problem of solving for the functions of an applied non zero H_z in [1] , [2] , [3]

$$\varphi = a \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} + b \cdot \begin{pmatrix} 0 \\ 1 \end{pmatrix} \doteq a \cdot |1\rangle + b \cdot |0\rangle \quad (20)$$

is much simpler than when H_z and H_x are both non zero. Is in the case of Eq. (4) with only H_z not equal to zero, then looking at the terms for **a** and **b** in the case of Eq. (4) is extremely simple, for the situation for Eq.(14) as diagrammed out above. It is the same problem for Eq. (19) and the much larger GW power situation,

but due to the ‘noisy’ values for **a** and **b**, then $\begin{pmatrix} a \\ b \end{pmatrix}$ is the same as looking at highly non-linear inputs into

$\begin{pmatrix} a \\ b \end{pmatrix}$ QM values which are still then mapped into the 3 dimensional CM results. . Still the same rotating rigid

body problem approximated by a rod in spatial rotation, but the movements and more would become much more

complicated. And then we find that $\begin{pmatrix} a \\ b \end{pmatrix}$ the input values are MUCH harder to solve.

7. Conclusion. CM and QM correspondence remains, but turbulence, a.k.a. Duerrer and Beckwith results for early universe GW generation makes the QM connection very hard to mathematically identify. Simple logical process, MESSY algebra ahead. With a possible answer to the question if singularities are essential in Cosmology.

Looking at Eq. (13) , simpler and harder cases, still in the case of relic GW production has large number correspondence and scaling as mentioned by Valev [6], with his H, not a Hamiltonian, but instead

$$r(\text{radius} - \text{of} - \text{universe}) \sim cH^{-1} \quad (21)$$

Also, the mass of the Universe, as given by Valev [6] is

$$M = (\text{Mass} - \text{of} - \text{universe}) \sim c^3 \cdot 2^{-1} \cdot (G \cdot H)^{-1} \quad (22)$$

More or less there is , when we look at physics innate simplicity in the inter relationship, of the sort mentioned by Valev, in terms of space-time geometry. The inter relationship of CM and QM given by Eq. (1) and then Eq. (3) with the stunning interplay between **x**, **y**, **z** and **a,b** given by Eq.(2) is, we believe, obscured by how complex

the problem is of finding $\begin{pmatrix} a \\ b \end{pmatrix}$. However, there is a tremendous inter relationship between

the quibits given in Eq.(20)and the inputs into Eq. (2), depending upon the values of inputs into the complex systems as given by Eq.(11) for CM, or the much simpler space geometry represented by the simple Eq.(5) as seen in references[7],[8]. Finally the author proposes that the inter linkage of QM to classical conditions in three sphere topology, which is where turbulence relevant to GW development and generation is the perfect template to answering a question raised by Stephen Kenneth Kauffman as to the reality of black holes and cosmological singularities. As related by Dr. Kauffmann, quantum mechanics routinely prunes out non-physical solutions in regards to hydrogen atom problems and picks only solutions which are physically feasible, which Kauffman asserts is not done in black hole and early

universe singularity physics. As written by Dr. Kauffmann to Dr. Corda, Dr. Beckwith, Johnathan Dickau, and others, there is a conscious choice given in classical GR to accept nonsensical solutions. The author (Beckwith) in [8] that GW may be generated in the electro weak era, at the beginning and perhaps as early as the Planckian regime of space-time. Furthermore Beckwith asserts that turbulence as signified by [7] is the driver for relic GW and GW power. If so, further development of the Hopf mapping results from 2 dimensions (QM) to 3 dimensions (CM) may enable us to identify necessary conditions for finding $\begin{pmatrix} a \\ b \end{pmatrix}$ in initial QM states necessary for relic GW development. It is well known that spherically symmetric geometry will generally not generate GW. This puts a restriction upon finding $\begin{pmatrix} a \\ b \end{pmatrix}$ which in turn would answer if an initial space-time singularity is even feasible. While String theory and loop quantum gravity via different mechanisms [9],[10] purport to have solutions to the initial cosmological state, the author views development of mandatory restrictions upon acceptable $\begin{pmatrix} a \\ b \end{pmatrix}$ quantum initial states for GW development as essential for finding and investigating rigorously the question if singularities are indeed necessary and even allowed in astrophysical problems. The author, also has an accepted publication in the Journal of Applied mathematics, SCIRP in part on this very question [11] with the answer that this question is not open and shut. I.e. that further work needs to be done in this area. This present Hopf mapping from QM to CM with , as argued here, CM turbulence generating GW, may be a way to determine restrictions upon initial states in QM which the author views, as a way to answer Dr. Kauffmann's question in a rigorous role in the near future.

References

1. A. L. Kholodenko, *Applications of Contact Geometry and Topology in Physics*, World Scientific, 2013
2. R. Feynman, F. Verno, and R. Hellwarth, "Geometrical representation of the Schrodinger Equation for solving Master Equation", *J. Appl. Phys* **28**, 49-52, (1957)
3. R. Feynman, *The Feynman Lectures on Physics, Volume 3*(Adsdison-Wesley, Reading , 1966)
4. A. Lightman, W. Press, R. Price, and S. Teukolsky, *Problem book in Relativity and Gravitation*, Princeton University Press, Princeton, New Jersey, 1979
5. M. Maggiore, *Gravitational Waves, Volume 1 : Theory and Experiment*, Oxford Univ. Press(2008)
6. D. Valev, "Estimation of the total mass and energy of the Universe", <http://arxiv.org/pdf/1004.1035.pdf>
7. R. Durrer, <http://arxiv.org/pdf/1303.7121.pdf>; <http://arxiv.org/abs/0901.0650>; R. Durrer, [Massimiliano Rinaldi](http://arxiv.org/abs/0901.0650) , Phys.Rev.D79:063507,2009, <http://arxiv.org/abs/0901.0650>
8. A. Beckwith, <http://vixra.org/abs/1308.0110>; ICISE inaugural conference accepted conference proceedings, in Quy Nohn, Vietnam.
9. T. Padmanabhan, *Gravitation, Foundations and frontiers*, Cambridge University Press, Cambridge, UK, 2010
10. G. Ellis, R. Maartens, M. Mac Callum, *Relativistic Cosmology* Cambridge University Press, Cambridge, UK, 2012
11. A. Beckwith, <http://vixra.org/abs/1304.0156> (accepted for publication in Journal for Applied mathematics, SCIRP, September, 2013)