

# QUANTUM THEORY DEPENDING ON MAXWELL EQUATIONS

WU SHENG-PING

ABSTRACT. This article try to unified the four basic forces by Maxwell equations, the only experimental theory. Self-consistent Maxwell equation with the current from electromagnetic field is proposed. and is solved to four kinds of electrons and the structures of particles. The static properties and decay and scattering are reasoned, all meet experimental data. The momentum-energy tensor of the electromagnetic field coming to the equation of general relativity is discussed. In the end that the conformation elementarily between this theory and QED and weak theory is discussed compatible, except some bias in some analysis.

## CONTENTS

1. Unit Dimension of $sch$	2
2. Quantization	2
3. Self-consistent Electrical-magnetic Fields	3
4. Stable Particle	4
5. Radium Function	4
6. Solution	5
7. Electrons and Their Symmetries	6
8. Mechanic Feature	7
9. Propagation and Movement	8
10. Antiparticle	10
11. Conservation Law and Balance Formula	11
12. Muon	11
13. Pion Positive	12
14. Pion Neutral	12
15. tau	13
16. Proton	13
17. Scattering	14
18. The Great Unification	15
19. Conclusion	15
References	16

---

*Date:* Jan 14, 2013.

*Key words and phrases.* Maxwell equations, Decay, Antiparticle.

1. UNIT DIMENSION OF  $sch$ 

A rebuilding of units and physical dimensions is needed. Time  $s$  is fundamental. The velocity of light is set to 1

$$Velocity : c = 1$$

Hence the dimension of length is

$$L : c$$

The  $\hbar$  is set to 1

$$Energy : \hbar$$

In Maxwell equations the following is set

$$c\epsilon = 1, c\mu = 1$$

One can have

$$\epsilon : \frac{Q^2}{\varepsilon L};$$

$$\mu : \frac{\varepsilon L}{c^2 Q^2}$$

$$UnitiveElectricalCharge : \sigma = \sqrt{\hbar}$$

It's very strange that the charge is analyzed as space and mass. Charge  $Q$  is then defined as  $Q/\sigma$  here,

$$\sigma = 1.03 \times 10^{-17} C = 64e, e/\sigma = e/\sigma = 1/64 = 1.56 \times 10^{-2}$$

$$H : Q/(LT) : \sqrt{\hbar}/c$$

$$E : \varepsilon/(LQ) : \sqrt{\hbar}/c$$

If  $\hbar, c$  is taken as a number instead of unit, then all physical units is described as the powers of the second:  $s^n$ .

The unit of charge can be reset by *linear variation of charge-unit*

$$Q \rightarrow CQ, Q : \sigma/C$$

We will use it without detailed explanation.

## 2. QUANTIZATION

All discussion base on a explanation of quantization, or *real* probability explanation for quantum theory, which bases on a Transfer Probability Matrix (TPM)

$$P_i(x)M = P_f(x)$$

As a fact, that a particle appears in a point at rate 1 is independent with appearing at anther point at rate 1. There still another pairs of independent states

$$S_1 = e^{ipx}, S_2 = e^{ip'x}$$

because

$$\langle s_1, s_2 \rangle_4 = \int dV s_1 s_2^* = N \delta(p - p')$$

$\langle s_1, s_2 \rangle_4$  means make product integrated in time-space. Similarly the symbol

$$\langle s_1, s_2 \rangle$$

is the product integrated in space and *always means its branch of zero frequency*. In fact in the TPM formulation, it's been accepted for granted that the Hermitian

inner-product is the measure of the dependence of two states, and it is also implied by the formula

$$P_1 M P_2^*$$

Depending on this view point one can constructs a wave

$$e^{ipx}$$

and gifts it with the momentum explanation  $p$ , Then all quantum theory is set up.

### 3. SELF-CONSISTENT ELECTRICAL-MAGNETIC FIELDS

The Maxwell equations are

$$\frac{\partial H}{\partial t} + \nabla \times E = 0$$

$$\frac{\partial E}{\partial t} - \nabla \times H + \mathbf{j} = 0$$

Try equation for the free E-M field

$$(3.1) \quad A_{,j}^{i,j} - A_{,j}^{j,i} = iA_{\nu}^* \cdot \partial^i A^{\nu} / 2 + cc., \quad Q_e = 1$$

$$(3.2) \quad Q_e = \int dV (iA_{\nu}^* \cdot \partial^t A^{\nu} / 2 + cc.)$$

It's in units  $e$ , in which the coefficients are 1. It's deduced by using momentum to express current.

$$(A^i) := (V, \mathbf{A}), (J^i) = (\rho, J)$$

$$\partial := (\partial_i) := (\partial_t, \partial_{x_1}, \partial_{x_2}, \partial_{x_3})$$

$$\partial' := (\partial^i) := (\partial_t, -\partial_{x_1}, -\partial_{x_2}, -\partial_{x_3})$$

The equation 3.1 have symmetry

$$CPT, cc.PT$$

If the gauge is

$$\partial_{\mu} A^{\mu} = 0$$

the continuous charge current meets

$$\partial_{\mu} \cdot J^{\mu} = 0$$

The energy of field  $A$  is  $\varepsilon = \int dV (E^2 + H^2) / 2$

$$\varepsilon = \langle \partial A | \partial A \rangle$$

This quantity always means its constant part.

For decay process of a particle make

$$\varepsilon = K(t)\varepsilon_i + (1 - K(t))\varepsilon_f$$

then

$$A := K(t)^{1/2} A_i + (1 - K(t))^{1/2} A_f + BG, K(0) = 1, K(\infty) = 0$$

$BG$  is background stable like

$$\sum_k b_k \sin(a_k t), a_k \in \mathbf{R}$$

By the equation 3.1 and zero border codition

$$\langle A^{\nu}, \partial' \partial A_{\nu} \rangle = BG + C = \langle \partial' A^{\nu}, \partial A_{\nu} \rangle$$

Substitute the expression of  $A$

$$\begin{aligned} & K'(t) \langle A_i^\nu, \partial_t A_{i\nu} \rangle + K'(t) \langle A_f^\nu, \partial_t A_{f\nu} \rangle \\ &= K(t) \langle \partial' A_i^\nu, \partial A_{i\nu} \rangle + (1 - K(t)) \langle \partial' A_f^\nu, \partial A_{f\nu} \rangle = BG + C \\ & K'(t)k_i/m_i + K'(t)k_f/m_f = K(t)m_i - K(t)m_f \end{aligned}$$

The  $m$  static mass of harmonic wave. The field is normalized with the static mass. Then we have

$$K(x) = e^{-Kt}, K = \frac{m_i - m_f}{k_i/m_i + k_f/m_f}$$

#### 4. STABLE PARTICLE

All particles are elementarily E-M fields is presumed. It's trying to find stable solution of the Maxwell equations *in complex domain*. One can write down a function initially and correct it by re-substitution. Here is the initial state

$$V = V_i e^{-ikt}, A_i = V$$

Substituting into equation 3.1

$$\begin{aligned} \partial_\mu \partial^\mu A_i^\nu - \partial^\nu \partial_\mu A_i^\mu &= 2J_i \\ 2J_i &= -\partial^\nu \partial_\mu A_i^\mu = -\partial^\nu \partial_t V \end{aligned}$$

It has the properties

$$\partial \cdot J_i = 0$$

$J_i$  causes the initial fields  $V$ , so that it is the real seed of recursive algorithm.

We can calculates the solution by recursive re-substitution for the two sides of the equation. The static fields  $E_0, H_0$

$$(4.1) \quad \begin{aligned} \nabla \cdot E_2 &= ie_{/\sigma} A_{i\nu}^* \cdot \partial_t A_i^\nu / 2 + cc. \\ \nabla \times H_2 &= -ie_{/\sigma} A_{i\nu}^* \cdot \nabla A_i^\nu / 2 + cc. \end{aligned}$$

We calls the fields' correction  $A_n$  with  $n$  degrees of  $A_i$  the  $n$  degrees correction.

#### 5. RADIUM FUNCTION

Firstly

$$\nabla^2 A = -k^2 A$$

is solved. Exactly, it's solved in spherical coordinate


$$0 = r^2 \nabla^2 f + k^2 f = (r^2 f_r)_r + k^2 r^2 f + \frac{1}{\sin \theta} (\sin \theta f_\theta)_\theta + \frac{1}{\sin^2 \theta} (f_\phi)_\phi$$

Its solution is

$$\begin{aligned} f &= R\Theta\Phi = R_l Y_{lm} \\ \Theta &= P_l^m(\cos \theta), \Phi = \cos(\alpha + m\phi) \\ R_l &= N \eta_l(kr), \eta_l(r) = r^l \int_0^\infty \frac{(1-\lambda)^l}{(1+\lambda)^{l+2}} \cos(\lambda r) d\lambda \\ &\int_0^\infty dr \cdot r^2 R^2 = 1 \end{aligned}$$

$R$  is solved like

$$\begin{aligned} (r^2 R_r)_r &= -k^2 r^2 R + l(l+1)R, l \geq 0 \\ R &\rightarrow rR' \\ (r^2 R')_{rr} &= -k^2 r^2 R' + l(l+1)R' \end{aligned}$$


 FIGURE 1. the shape of radium function  $R_1$  by DFT

$$R' \rightarrow r^{l-1}R'$$

$$rR'_{rr} + 2(l+1)R'_r + k^2rR' = 0$$

$$r \rightarrow r/k$$

$$(s^2F)' + 2(l+1)F + F' = 0, F = F(R')$$

$F()$  is the Fourier transform

$$R' = \int_0^\infty \frac{(1-\lambda)^l}{(1+\lambda)^{l+2}} \cos(\lambda r) d\lambda$$

The function  $R_1$  has zero derivative at  $r = 0$  and is zero as  $r \rightarrow \infty$ .

## 6. SOLUTION

The derivatives of the function of electron has a strange breaking point in coordinate origin hence without normal convenience of Fourier transform. The following are some proximation of the first rank. The solution of  $l = 1, m = 1, Q = e/\sigma$  is calculated or tested for electron.

$$A_1 = NR_1(kr)Y_{1,1},$$

The curve of  $R_1$  is like the one in the figure 1.

The magnetic dipole moment  $\mu_z$  is calculated as the first rank of proximation

$$\mu_z = \langle A_\nu | -i\partial_\phi | A^\nu \rangle / 2$$

$$= 1/2, k_e = 1, Q_e = 1$$

The power of unit of charge is not equal, but it's valid for unit  $Q = e$ .

$$\frac{1}{2} = \mu_B, k_e = 1, Q_e = 1$$

## 7. ELECTRONS AND THEIR SYMMETRIES

Some states of electrical field  $A$  are defined as the core of the electron, it's the initial function  $A_1 = V$  for the re-substitution to get the whole electron function.

$$\begin{aligned} e_r^+ &: NR_1(kr)Y_{1,1}e^{-ikt} \\ e_l^+ &: NR_1(kr)Y_{1,-1}e^{-ikt} \\ e_r^- &: -NR_1(kr)Y_{1,1}e^{ikt} \\ e_l^- &: -NR_1(kr)Y_{1,-1}e^{ikt} \\ k &= m_e \end{aligned}$$

$r, l$  is the direction of the spin. We use these symbols  $e$  to express the complete potential field  $A$  or the abstract particle.

Energy of static E-field crossing is discussed. In the zero rank of correction ie. the static field is

$$e^*(i\partial')e = J_e, Q_e = 1$$

then the charge is normalized

$$Q = \langle e_\mu | i\partial_t | e^\mu \rangle = 1, Q_e = 1$$

Hence

$$\langle \partial e | \partial e \rangle / 2 = k_e / e_{/\sigma}, k_e = m_e$$

The static energy of electric field between  $A_2$  is

$$\varepsilon_q = -e_{/\sigma}^3 k_e / 2 = -\frac{1}{6.7 \times 10^{-16} s}$$

Energy of the static M-field crossing

$$\varepsilon'_m = \varepsilon_e$$

Hence the gross energy is

$$\varepsilon_e = 2\varepsilon_q = -\frac{1}{3.355 \times 10^{-16} s}$$

The value of crossing term generated by static fields between electrons are

$\varepsilon_e$	$e_r^+$	$e_r^-$	$e_l^+$	$e_l^-$
$e_r^+$	+	-	0	0
$e_r^-$	-	+	0	0
$e_l^+$	0	0	+	-
$e_l^-$	0	0	-	+

The field of four kinds of electrons has symmetries

$$\begin{aligned} e_l^+ &: E_n + M_n \\ e_r^+ &: E_n - M_n \\ e_l^- &: (-1)^{n-1} E_n + (-1)^{n/2} M_n \\ e_r^- &: (-1)^{n-1} E_n - (-1)^{n/2} M_n \end{aligned}$$

$E : E(A_i)$  is the abstract electrical field,  $M : M(A_i)$  is the abstract magnetic field, the derivatives are solved.  $n$  is the degree of the correction. The non zero crossing in re-substitution is the crossing with  $A_i$ . The higher absolute frequency than  $k_e$  is also zero.

Calculating the crossing part between  $e_r^+, e_l^-$ . the non zero results of crossing is between  $A_2$  and  $A_6$  or between  $A_4$ .

$$\varepsilon_x \approx -e_{/\sigma}^7 k_e / 4 \approx -\frac{1}{2.18 \times 10^{-8} s}$$

The theorem 9.6 is used. The value of this crossing term generated between electrons are

$\varepsilon_x$	$e_r^+$	$e_r^-$	$e_l^+$	$e_l^-$
$e_r^+$	+	0	0	-
$e_r^-$	0	+	-	0
$e_l^+$	0	-	+	0
$e_l^-$	-	0	0	+

## 8. MECHANIC FEATURE

If the equation that connects space and E-M fields is written down for cosmos of electrons, it's the following:

$$(8.1) \quad R_{ij} - \frac{1}{2} R g_{ij} = 8\pi G T_{ij} / c^4$$

$$T_{ij} = F_i^{k*} F_{kj} - g_{ij} F_{\mu\nu} F^{\mu\nu*} / 4$$

$F$  is the electromagnetic tensor. This equation give mass because the space is decided by E-M fields instantly.

Because fields  $F$  is additive, the group of electrons are express by:

$$F = \sum_i f_i * \partial e_i, \langle f_i | f_i \rangle = 1$$

*The convolution is made only in space:*

$$f * g = \int dV f(t, y - x) g(t, x)$$

It's called *propagation*. Each  $f_i$  is normalized to 1. We always use

$$\sum_i f_i * e_i, \sum_i f_i * \partial e_i$$

to express its abstract construction and the field.

When the mechanical physical is discussed, observing the Energy-Momentum tensor  $T$  we have the momentum is

$$(8.2) \quad p_{,0}^\mu = T_0^\mu$$

$$\varepsilon = \langle \partial A | \partial A \rangle, \mathbf{p}_{,t} = E \times B$$

So that the spin of electron is calculated as

$$S_e = \langle A | \partial_\phi \cdot \partial_t | A \rangle / 2 = 1/2$$

The MDM (magnetic Dipole moment) of electron is calculated by the equation 3.1

$$\mu_e = \frac{1}{2} \langle A | -i \partial_\phi | A \rangle = 1 / (2k_e), Q_e = 1$$

## 9. PROPAGATION AND MOVEMENT

Define symbols for particle  $x$

$$\begin{aligned} e_{xr}^+ &:= N \cdot R_1(k_x r) Y(1, 1) e^{-ik_x t}, \\ e_{xx}^+ &:= (e_{xl}^+ + e_{xr}^+) / \sqrt{2} \\ &< e_x | e_x \rangle = 1 \end{aligned}$$

The propagations is the  $f(x)$  in

$$f(x) * e$$

The following are stable propagation:

<i>particle</i>	<i>electron</i>	<i>photon</i>	<i>neutino</i>
<i>notation</i>	$e_r^+$	$\gamma_r$	$\nu_r$
<i>structure</i>	$e_r^+$	$(e_r^+ + e_r^-)$	$(e_r^+ + e_l^-)$

By mathematic

$$\varsigma_{k,l,m}(x) := R_l(kr) Y_{l,m}, \varsigma_k(x) := \varsigma_{k,1,\pm 1}(x)$$

meets the following results

**Theorem 9.1.**  $C_A$  is a global area with its center in  $A$  and its diameter is  $r_A$

$$\lim_{r_o=r_y \rightarrow 0} \int_{I-\sum C_i} dV \varsigma_k(x) \varsigma_k^*(x-y) = 0, y \neq O$$

For this function, it's strange in grid origin.

*Proof.* Use the limit

$$\lim_{k' \rightarrow k} \lim_{r_o=r_y \rightarrow 0} \left( \int_{I-\sum C_i} dV \varsigma_k(x) \varsigma_{k'}^*(x-y) \right)$$

□

We have of course:

$$\varsigma_k(\mathbf{r}) = \int d\mathbf{p} C_p e^{i\mathbf{p}\mathbf{r}}, \mathbf{p}^2 = k^2$$

**Theorem 9.2.**

$$\langle f(x) * \varsigma_k | g(x) * \varsigma_k \rangle = \langle f(x) | g(x) \rangle$$

$|f|^2, |g|^2$  is integrable.

It's proved by

$$\langle \sum_i a_i \varsigma_k(x-x_i) | \sum_i b_i \varsigma_k(x-x_i) \rangle = \sum_i a_i^* b_i$$

**Theorem 9.3.** if  $e^{i\mathbf{p}\mathbf{r}}, \varsigma_k$  is normalized to 1,

$$e^{i\mathbf{p}\mathbf{r}} * \varsigma_k = \omega e^{i\mathbf{p}\mathbf{r}}, |\omega| = 1$$

**Theorem 9.4.**

$$\nabla(\varsigma_k * \varsigma_{k'}) = (\nabla \varsigma_k) * \varsigma_{k'} + \varsigma_k * \nabla(\varsigma_{k'})$$



$$\begin{aligned}
 & -\partial_y \int dV_x I(y-x) \varsigma_k(x-y) \varsigma_{k'}(x) \\
 &= -\int dV_x I'(y-x) \varsigma_k(x) \varsigma_{k'}(x) \\
 &= \int dV_x I(x-y) (\varsigma_k(x-y) \varsigma_{k'}(x))_x \\
 &= \int dV_z I(z) (\varsigma_k(z) \varsigma_{k'}(z+y))_{z, z=x-y} \\
 &= \int dV_z (\varsigma'_k(-z) \varsigma_{k'}(z+y) + \varsigma_k(-z) \varsigma'_{k'}(z+y)) \\
 &= \int dV_z (\varsigma'_k(z) \varsigma_{k'}(-z+y) + \varsigma_k(z) \varsigma'_{k'}(-z+y)) \\
 & I(y-x) := \begin{cases} 0, & x \neq O \\ 1, & x = O \end{cases}
 \end{aligned}$$

**Theorem 9.5.**

$$\langle (\nabla \varsigma_k) * \varsigma_1 | \varsigma_k * \varsigma_1 \rangle = \langle k \varsigma_k * \nabla \varsigma_1 | \varsigma_k * \varsigma_1 \rangle$$

**Theorem 9.6.**

$$\varsigma_1 * \frac{1}{r} = \varsigma_1$$

$C$  is relative to the measure of sampling dense of integration

$$\begin{aligned}
 & \int dV \varsigma_1 \cdot \varsigma_1^* \cdot \varsigma_1 \cdot \varsigma_1^* \\
 &= C \int dP \cdot F(\varsigma_1) * F(\varsigma_1^*) * F(\varsigma_1) * F(\varsigma_1^*) \\
 &= \int dP \omega * \omega^* * \omega * \omega^* = 1
 \end{aligned}$$

**Theorem 9.7.**

$$\begin{aligned}
 & \langle (\nabla \varsigma_k) * \varsigma_1 | (\nabla \varsigma_k) * \varsigma_1 \rangle = \langle (\nabla \varsigma_k) * \varsigma_1 | \varsigma_k * (\nabla \varsigma_1) \rangle \\
 & \approx k \langle \varsigma_k * \varsigma_1 | \varsigma_k * \varsigma_1 \rangle
 \end{aligned}$$

The figure 2 is the shape of distribution of momenta of electron function  $e_x$ .

The movement of the propagation is called *Movement*, ie. the third level wave, harmonic wave. The mechanical movements  $p$  of particle  $e_x * \sum e$  by relative theory is

$$e^{i\mathbf{P}\mathbf{X}-ikt} * e_{\mathbf{x}} e^{-ik_x t} * \sum e$$

the dual mechanical movement is

$$e^{i\mathbf{P}\mathbf{X}+ikt} * e_{\mathbf{x}} e^{-ik_x t} * \sum e$$

The measure of the movement of the field  $e_{\mathbf{x}}$  meets

$$-p^\mu p_\mu + k_x^2$$

Because the self coordinate measure of the particle  $e_x$  is  $k_x^2$ , so that

$$p^\mu p_\mu = 0, \mathbf{p}^2 - (k^2 + k_x^2) = -k_x^2$$




FIGURE 2. the shape of distribution of radioactive momenta of electron fields in one direction:  $k/(1+k)^4 - 4k/(1+k)^5$ , calculated through spherical Bessel functions

The static MDM (magnetic dipole moment) is approximately by the condition 8.2

$$\mu = \sum_i \langle \int dx_i \cdot e_x * \partial e_i(x_i) | -i\mathbf{r} \times \nabla | \int dx_i \cdot e_x * \partial e_i(x_i) \rangle / 2., Q_e = 1$$

$$\mu_z = \sum_i \langle e_x * e_i(x_i) | e_x * (-i\partial_\phi e_i(x_i)) \rangle > \frac{k_e}{2k_x}$$

Its spin is approximately

$$S_z = \sum_i \langle \int dx \cdot e_x * \partial e_i(x_i) | -\partial_\phi \partial_t | \int dx \cdot e_x * \partial e_i(x_i) \rangle / 2., Q_e = 1$$

$$= \sum_i \langle e_x * e_i(x_i) | e_x * (-i\partial_\phi e_i(x_i)) \rangle > k_{ei}/2$$

Mechanical spin decouples between electrons.

Calculating the following coupling system

$$F = e_x * \sum_i \partial e_i$$

$$\partial \cdot \partial' e_x = 0$$

$$(9.1) \quad e_x = e^{-iNt} \zeta_N, N \approx \langle \partial A | \partial A \rangle / 2, Q_e = 1$$

## 10. ANTIPARTICLE

Antimatter is the positive matter reverse world-line (PT), so that it meets

$$\partial_\nu \partial^\nu A^i = -ie_{/\sigma} A_\nu^* \cdot \partial^i A_\nu / 2 + cc.$$

From

$$\partial_\nu \partial^\nu (A^i(x) + B) = ie_{/\sigma} (-A_\nu(x) + B)^* \cdot \partial^i (A_\nu(x) + B) / 2 + cc.$$

We have

$$\partial_\nu \partial^\nu (A^i(-x) + B) = ie_{/\sigma} (A_\nu(-x) + B)^* \cdot \partial^i (A_\nu(-x) + B) / 2 + cc.$$

$B$  is outer field the particle is in. If  $A(x)$  describes positive matter,  $A(-x)$  is describes antimatter, we define

$$\overline{A(x)} := A(-x)$$

We have the reaction in four-dimension map

$$p \rightarrow, A(x) \rightarrow \bullet \rightarrow p'$$

equivalent to

$$p \rightarrow \bullet \rightarrow A(-x), \rightarrow p'$$

and

$$\overline{e_r^+} \approx e_l^-$$

### 11. CONSERVATION LAW AND BALANCE FORMULA

No matter in E-M fields level or in movement (the third) level, the conservation law is *conservation of momentum and conservation of angular momentum*. A *balance formula* for a reaction is the equivalent formula in positive matter, ie. after all anti-matter is shifted to the other side of the reaction formula. Balance formula is suitable for the analysis of the energy transition of E-M fields in the reaction. The *invariance of electron itself* in reaction is also a conservation law.

### 12. MUON

Generally, there are kinds of energy increments.

Weak coupling

$$W : \langle e_r^+ | e_l^- \rangle$$

Light coupling

$$L : \langle e_r^+ | e_r^- \rangle$$

Weak side coupling

$$Ws : \langle e_r^+ | e_l^- \rangle - \langle e_x * e_r^+ | e^{ipx} * e_l^- \rangle$$

Light side coupling

$$Ls : \langle e_r^+ | e_r^- \rangle - \langle e_x * e_r^+ | e^{ipx} * e_r^- \rangle$$

Strong coupling

$$S : \langle e_r^+ | e_r^+ \rangle$$

Because of the degrees of the derivatives the anti-matter couplings are the same except the strong coupling.

$\mu$  is composed of

$$\mu_r^+ : e_\mu * (e_r^+ + \overline{\nu}_r)$$

From the equation 8.1 and 9.1 and the other deductive,  $\mu$  is with mass  $3m_e/e_\sigma^2 = 3 \times 64k_e$ , spin 1/2, MDM  $\mu_B k_e/k_\mu$ .

The main channel of decay

$$\begin{aligned} \mu_r^+ &\rightarrow M_l^+ + \overline{\nu}_l \\ M_r^+ &= e_M * (\overline{e}_l^- + \nu_l) \end{aligned}$$

$$e_\mu * e_r^+ + e^{-ip_1x} * e_M^* * e_l^- + \overline{e^{ip_2x}} * \nu_l \rightarrow \overline{e_\mu} * \nu_r + \overline{e^{ip_1x}} * \overline{e_M} \nu_l$$

The outer waves  $e_\mu$  and  $e^{-ip_1x} * e_M^*$ ,  $e_\mu$  and  $e^{-ip_2x}$  are coupling. The energy difference is kind of  $Ws$ , the interacting field is  $E$  that between  $A_2, A_6$ .

$$2 < \overline{e_\mu} * \partial e_r^+ | \overline{e_\mu} * \partial e_l^- > - 2 < \overline{e_\mu} * \partial e_r^+ | e^{i\mathbf{p}_1\mathbf{r} + ik_\mu t} * \partial e_l^- >$$

$$p_i^\mu p_{i\mu} = 0$$

$$= 2(1 - \frac{k_\mu}{k_\mu + k_e}) < \overline{e_\mu} * \partial e_l^+ | e^{i\mathbf{p}_1\mathbf{r} + ik_\mu t} * \partial e_r^- >$$

$$= 2(1 - \frac{k_\mu}{k_\mu + k_e}) < \overline{e_\mu} * e_l^+ | e^{i\mathbf{p}_1\mathbf{r} + ik_\mu t} * \nabla^2 e_r^- >$$

sum up in spectrum of  $p_1 : p_1^\mu p_{1\mu} = 0$

$$= \frac{2k_e \varepsilon_x}{k_\mu}$$

The emission of decay is

$$= -\frac{1}{2.1 \times 10^{-6}s} [2.1970 \times 10^{-6}s][1]$$

The data in square bracket is experimental data of the full width. The decay of particle  $M$  is like a scattering with no energy emission

$$M_r^+ \rightarrow \overline{e_l^-} + \nu_l$$

### 13. PION POSITIVE

Pion positive is

$$\pi_l^- : e_\pi * (\overline{e_r^+} + e_l^-) + e_\pi^* * e_r^+$$

It's with mass  $5 \times 64m_e$ , spin 1/2 and MDM  $\mu_B k_e / k_\pi^+$ .

Decay Channels:

$$\pi_l^- \rightarrow \mu_l^- + \nu_r$$

It's with balance formula

$$e_\pi^* * e_r^+ + e_\pi * e_l^- + \overline{e^{ip_1x}} * \overline{e_\mu} * \nu_r \rightarrow \overline{e_\pi} * e_r^+ + e^{ip_1x} * e_\mu * e_l^- + e^{ip_2x} * \nu_r$$

The emission of energy is kind of  $W$

$$\varepsilon_x = -\frac{1}{2.18 \times 10^{-8}s} [(2.603 \times 10^{-8}s)[1]$$

The referenced data is the full width.

### 14. PION NEUTRAL

Pion neutral is atom-like particle

$$\pi^0 : e_{\pi^0} * \nu_r + e_{\pi^0}^* * \nu_l$$

It has mass  $4 \times 64m_e$ , zero spin and zero MDM. Its decay modes are

$$\pi^0 \rightarrow \gamma_r + \gamma_l$$

The loss of energy is kind of  $L$

$$4\varepsilon_e = -\frac{1}{8.39 \times 10^{-17}s} [8.4 \times 10^{-17}s][1]$$

## 15. TAU

$\tau$  maybe that

$$\tau_l^+ : e_\tau * (5e_r^+ + \overline{5e_r^+ + e_r^-})$$

Its mass  $51 \times 64m_e$ , spin  $1/2$ , MDM  $\mu_B k_e / k_\mu$ . It has decay mode

$$\tau_l^+ \rightarrow \mu_l^+ + \nu_l + \bar{\nu}_l$$

$$e_\tau * 5e_r^+ + \overline{e^{ip_1x}} * \overline{e_\mu} * \nu_l + \overline{e^{ip_2x}} * \nu_l \rightarrow \overline{e_\tau} * 5e_r^+ + \overline{e_\tau} * e_r^- + e^{ip_1x} * e_\mu * e_l^+ + e^{ip_3x} * \nu_l$$

The energy gap is kind of  $LS$ ,  $E = A_2$

$$\begin{aligned} & 5 < \overline{e_\tau} * \partial E_r^+ | \overline{e_\tau} * \partial E_r^- > \\ & -5 < \overline{e_\tau} * \partial E_r^+ | e^{i\mathbf{p}_1 \mathbf{r} - ik_\tau t} * \partial E_r^- > \\ & p_i^\mu p_{i\mu} = 0 \\ & = 5 \left(1 - \frac{k_\tau}{k_\tau + k_e}\right) < \overline{e_\tau} * E_r^+ | e^{i\mathbf{p}_1 \mathbf{r} - ik_\tau t - ik_e t} * \nabla^2 E_r^- > \\ & = \frac{5\varepsilon_e}{k_\tau / k_e} \\ & = -\frac{1}{6 \times 10^{-14} \text{s}} [2.9 \times 10^{-13} \text{s}, BR.0.17][1] \end{aligned}$$

Depending on this kinds of particle including

$$q_r^{n+} := n(e_r^+, \overline{e_r^+})$$

we can construct particles of great mass decaying without strong emission (light radiative), for example

$$e_L * (q_r^{n+}, \overline{e_l^-})$$

This series of particle has included  $\mu, \tau$  and in fact almost all light radiative particles are of this kind,  $W, Z$  and Higgs particle is reasonably are of this kind, they are created in colliding. Another condition is possibly that, in the collision, the created light radioactive particle ( $q_r^{n+}, \overline{e_l^-}$ ) with different  $n$  is mixed to some rates as to the detector can't distinguish them.

Because the channel width decides the channel branch rates, obviously the most experimental data violate this rule. So that the channels listing after the same name of a particle in fact belong to different particles.

The particle  $K^+$  possible is

$$K^+ = (q_r^{3+}, \overline{\nu_l + e_l^-}) \rightarrow \mu_r^+ + \bar{\nu}_l$$

It has emission of  $LS$ .

## 16. PROTON

Proton may be like

$$p_r^- : e_p * (e_r^+ + e_r^- + e_l^+ + e_l^- + \overline{e_l^+ + 2e_r^+}) + e_p^* * (e_r^+ + e_r^- + e_l^+ + e_l^- + \overline{2e_l^-})$$

The mass is  $27 \times 64m_e$  that's very close to the real mass. The MDM is calculated as  $3\mu_N$ , spin is  $1/2$ . The proton thus designed is eternal because if it decay even to the finest blocks the energy of emission is negative.

We define an unit: Mass-number Unit

$$m = m_e \sigma / e \approx 64m_e$$

## 17. SCATTERING

The scattering can be calculated as dynamic electromagnetic mechanical theory, ie. the magnitude scattered is

$$-ie \int dV_4 \hat{j}^\mu \hat{A}_\mu$$

From the equation 3.1 the operator of current is

$$2j^\mu = ie_{/\sigma} A_\mu^* \partial A^\mu - ie_{/\sigma} A^\mu \partial A_\mu^*, Q_e = 1$$

The reaction is like

$$\sum_i f_i * e_i \rightarrow \sum_i f'_i * e_i$$

$e_i$  are all positive matters. The interaction between electrons

$$I(e_i, e_j) = I(J(e_i), J(e_j))$$

is the *cross interaction*. The interaction is

$$I(j_1, j_2) = \int dV_4 A_1 J_2,$$

$$2J = ie_{/\sigma} A_i^* \partial A_f - ie_{/\sigma} A_f \partial A_i^*, Q_e = 1$$

For example the scattering

$$e^{ip_1x} * e_r^+ + e^{ip_2x} * e_r^- \rightarrow e^{ip_3x} * e_r^+ + e^{ip_4x} * e_r^-$$

The transfer is

$$i\mu \approx \frac{C(p'_1 + p'_3)^\nu (p'_2 + p'_4)_\nu}{(p'_1 - p'_3)^2}$$

The  $p'_i$  is the *cap momentum* relative to  $p_i$ . And we have

$$\int dV_4 I(J(e_i), J(e_j)) = \varepsilon_e$$

this calculation conforms to classical theory

$$iC = e^2 = \frac{\varepsilon_e}{k_e e}$$

The interaction is between  $A_0$ . In the mean effect rate of transfer for the scattering of one to one particles is

$$\frac{|\mu|^2}{2k_1 \cdot 2k_2 \cdot 2k_3 \cdot 2k_4}$$

The energy gap in fact is part of interaction for example

$$e^{p_1x} * e + e^{p_2x} * e \rightarrow e^{p_3x} * e + e^{p_4x} * e$$

$e$  is the same electron of the four kind and of the same polarization. In fact the final state includes

$$s := e_s * (e + e)$$

The interaction in the reaction is

$$I = I(j_{13}, j_{24}) = \int dV_4 A_{13} J_{24}^* \delta(x - G)$$

The domain  $G$  is the domain meeting Clain-Golden equation and momentum conservations, for all emitted matter. Taking the mass center system, the part generating  $s$  is

$$e + e \rightarrow s = e^{-ikt} e^{-ik_s t} \zeta_{k_s} * (e + e)$$

$$k + k_s = k_1$$

Using

$$\varsigma_{k_s} = \int d\mathbf{p} C_p e^{i\mathbf{p}\mathbf{r}}, \mathbf{p}^2 = k_s^2$$

the invariable magnitude is

$$i\mu \approx C \int d\mathbf{p} C_p \frac{-(\mathbf{p} + \mathbf{p}_1)^2 + 4k_1^2}{(\mathbf{p} - \mathbf{p}_1)^2}, \mathbf{p}^2 = k_s$$

For a decay with two particles emission, the first order term of the scattering effect is identity of the calculation of energy difference between the initial and final state, which a easy analysis can prove.

### 18. THE GREAT UNIFICATION

Firstly we redefine the unit second to simplify the equation 8.1

$$1 = 8\pi G/c^4$$

$$(8\pi G)_{T=s} C T^2 / \hbar = 1, c = 1, \hbar = 1$$

The general relative equation is

$$T_{ij} = R_{ij} - g_{ij} R/2$$

the EM equation is

$$T_{ij} = F_{ik}^* F_j^k - g_{ij} F_{\mu\nu}^* F^{\mu\nu} / 4$$

We observe that the co-invariant curvature is

$$R_{ij} = F_{ik}^* F_j^k + g_{ij} F_{\mu\nu}^* F^{\mu\nu} / 8$$

### 19. CONCLUSION

The relative theory is applied to electromagnetic wave to give the mechanism meaning of the fields, by energy-momentum tensor. In my view point the sum-up of the grains (as electrons) of electromagnetic field is expression of mechanic movement. Fortunately this model explained all the effects in the known world: strong, weak and electromagnetic effects, and even subclassify them further if not add new ones. In this model the only field is electromagnetic field, and this stands for the philosophical with the point of that unified world is from an unique source. All depend on a simple fact: the current of matter in a system is time-invariantly zero in mass-center frame, and we can devise current of matter to analysis the E-M current. So that all effects is explained with diffusion process.

Except electron function my description of particles in fact has the same form with Quantum Electromagnetic Mechanics, and they two should reach the same result except for some little bias. But my theory isn't compatible to the theory of quarks, the upper part of standard model, if not it is calculated in the style of Quantum Electromagnetic Mechanics. In fact, The electron function is a good promotion for the experimental model of proton that went up very early.

Underline my calculations a fact is that the electron has the same phase (electron resonance), which the Cosmos Explosion will explain, all electrons are generated in the same time an place, the same source.

## REFERENCES

- [1] K. Nakamura et al. (Particle Data Group), JPG 37, 075021 (2010) (URL: <http://pdg.lbl.gov>)  
*E-mail address: hiyaho@126.com*

WUHAN UNIVERSITY, WUHAN, HUBEI PROVINCE, THE PEOPLE'S REPUBLIC OF CHINA. POST-CODE: 430074