

Connections between Extenics and Refined Neutrosophic Logic

Florentin Smarandache, Ph. D.
University of New Mexico
Math & Science Division
705 Gurley Ave.
Gallup, NM 87301, USA
E-mail: smarand@unm.edu

Abstract.

The aim of this presentation is to connect Extenics with new fields of research, i.e. fuzzy logic and neutrosophic logic.

We show herein:

- How Extenics is connected to the 3-Valued Neutrosophic Logic,
- How Extenics is connected to the 4-Valued Neutrosophic Logic,
- How Extenics is connected to the n -Valued Neutrosophic Logic,

when contradictions occurs.

Introduction.

In this paper we present a short history of logics: from particular cases of 2-symbol or numerical valued logic to the general case of n -symbol or numerical valued logic, and the way they are connected to Prof. Cai Wen's Extenics Theory (1983). We show generalizations of 2-valued Boolean logic to fuzzy logic, also from the Kleene's and Lukasiewicz' 3-symbol valued logics or Belnap's 4-symbol valued logic to the most general *n -symbol or numerical valued refined neutrosophic logic*. Two classes of neutrosophic norm (*n -norm*) and neutrosophic conorm (*n -conorm*) are defined. Examples of applications of neutrosophic logic to physics are listed in the last section.

Similar generalizations can be done for *n -Valued Refined Neutrosophic Set*, and respectively *n -Valued Refined Neutrosophic Probability* in connections with Extenics.

1. Two-Valued Logic

a) The Two Symbol-Valued Logic.

It is the Chinese philosophy: *Yin* and *Yang* (or Femininity and Masculinity) as contraries:



Fig 1. Ying and Yang

It is also the Classical or *Boolean Logic*, which has two symbol-values: truth T and falsity F .

b) The Two Numerical-Valued Logic.

It is also the Classical or *Boolean Logic*, which has two numerical-values: truth 1 and falsity 0 .

More general it is the *Fuzzy Logic*, where the truth (T) and the falsity (F) can be any numbers in $[0, 1]$ such that $T + F = 1$.

Even more general, T and F can be subsets of $[0, 1]$.

2. Three-Valued Logic

a) The Three Symbol-Valued Logics:

i) *Lukasiewicz's Logic*: True, False, and Possible.

ii) *Kleene's Logic*: True, False, Unknown (or Undefined).

iii) Chinese philosophy extended to: *Yin*, *Yang*, and *Neuter* (or Femininity, Masculinity, and Neutrality) – as in Neutrosophy.

Neutrosophy philosophy was born from neutrality between various philosophies. Connected with Extenics (Prof. Cai Wen, 1983), and Paradoxism (F. Smarandache, 1980).

Neutrosophy is a new branch of philosophy that studies the origin, nature, and scope of neutralities, as well as their interactions with different ideational spectra.

This theory considers every notion or idea $\langle A \rangle$ together with its opposite or negation $\langle \text{anti}A \rangle$ and with their spectrum of neutralities $\langle \text{neut}A \rangle$ in between them (i.e. notions or ideas supporting neither $\langle A \rangle$ nor $\langle \text{anti}A \rangle$).

The $\langle \text{neut}A \rangle$ and $\langle \text{anti}A \rangle$ ideas together are referred to as $\langle \text{non}A \rangle$.

Neutrosophy is a generalization of Hegel's dialectics (the last one is based on $\langle A \rangle$ and $\langle \text{anti}A \rangle$ only).

According to this theory every idea $\langle A \rangle$ tends to be neutralized and balanced by $\langle \text{anti}A \rangle$ and $\langle \text{non}A \rangle$ ideas - as a state of equilibrium.

In a classical way $\langle A \rangle$, $\langle \text{neut}A \rangle$, $\langle \text{anti}A \rangle$ are disjoint two by two. But, since in many cases the borders between notions are vague, imprecise, Sorites, it is possible that $\langle A \rangle$, $\langle \text{neut}A \rangle$, $\langle \text{anti}A \rangle$ (and $\langle \text{non}A \rangle$ of course) have common parts two by two, or even all three of them as well. Such contradictions involves Extenics.

Neutrosophy is the base of all neutrosophics and it is used in engineering applications (especially for software and information fusion), medicine, military, airspace, cybernetics, physics.

b) The Three Numerical-Valued Logic:

i) *Kleene's Logic*: True (1), False (0), Unknown (or Undefined) ($1/2$), and uses “min” for \wedge , “max” for \vee , and “1-” for negation.

ii) More general is the *Neutrosophic Logic* [Smarandache, 1995], where the truth (T) and the falsity (F) and the indeterminacy (I) can be any numbers in $[0, 1]$, then $0 \leq T + I + F \leq 3$.
 More general: Truth (T), Falsity (F), and Indeterminacy (I) are standard or nonstandard subsets of the nonstandard interval $] -0, 1 + [$.

When $t + f > 1$ we have conflict, hence Extenics.

3. Four-Valued Logic

a) The Four Symbol-Valued Logic

i) It is *Belnap's Logic*: True (T), False (F), Unknown (U), and Contradiction (C), where T, F, U, C are symbols, not numbers.

Now we have Extenics, thanks to $C =$ contradiction.

Below is the Belnap's conjunction operator table:

\cap	F	U	C	T
F	F	F	F	F
U	F	U	F	U
C	F	F	C	C
T	F	U	C	T

Table 1.

Restricted to T, F, U , and to T, F, C , the Belnap connectives coincide with the connectives in Kleene's logic.

ii) Let $G =$ Ignorance. We can also propose the following two 4-Symbol Valued Logics: (T, F, U, G), and (T, F, C, G).

iii) *Absolute-Relative 2-, 3-, 4-, 5-, or 6-Symbol Valued Logics* [Smarandache, 1995].

Let T_A be truth in all possible worlds (according to Leibniz's definition);

T_R be truth in at last one world but not in all worlds;

and similarly let I_A be indeterminacy in all possible worlds;

I_R be indeterminacy in at last one world but not in all worlds;

also let F_A be falsity in all possible worlds;

F_R be falsity in at last one world but not in all worlds;

Then we can form several Absolute-Relative 2-, 3-, 4-, 5-, or 6-Symbol Valued Logics just taking combinations of the symbols $T_A, T_R, I_A, I_R, F_A,$ and F_R .

As particular cases, very interesting would be to study the Absolute-Relative 4-Symbol Valued Logic (T_A, T_R, F_A, F_R), as well as the Absolute-Relative 6-Symbol Valued Logic ($T_A, T_R, I_A, I_R, F_A, F_R$).

b) *Four Numerical-Valued Neutrosophic Logic*: Indeterminacy I is refined (split) as $U =$ Unknown, and $C =$ contradiction.

T, F, U, C are subsets of $[0, 1]$, instead of symbols;

This logic generalizes Belnap's logic since one gets a degree of truth, a degree of falsity,

a degree of unknown, and a degree of contradiction.
 Since $C = T \wedge F$, this logic involves the Extenics.

4. Five-Valued Logic

a) *Five Symbol-Valued Neutrosophic Logic* [Smarandache, 1995]:

Indeterminacy I is refined (split) as $U = \text{Unknown}$, $C = \text{contradiction}$, and $G = \text{ignorance}$;
 where the symbols represent:

$T = \text{truth}$;
 $F = \text{falsity}$;
 $U = \text{neither } T \text{ nor } F \text{ (undefined)}$;
 $C = T \wedge F$, which involves the Extenics;
 $G = T \vee F$.

b) If T, F, U, C, G are subsets of $[0, 1]$ then we get: a *Five Numerical-Valued Neutrosophic Logic*.

5. Seven-Valued Logic

a) *Seven Symbol-Valued Neutrosophic Logic* [Smarandache, 1995]:

I is refined (split) as U, C, G , but T also is refined as $T_A = \text{absolute truth}$ and $T_R = \text{relative truth}$, and F is refined as $F_A = \text{absolute falsity}$ and $F_R = \text{relative falsity}$. Where:

$U = \text{neither } (T_A \text{ or } T_R) \text{ nor } (F_A \text{ or } F_R) \text{ (i.e. undefined)}$;
 $C = (T_A \text{ or } T_R) \wedge (F_A \text{ or } F_R) \text{ (i.e. Contradiction)}$, which involves the Extenics;
 $G = (T_A \text{ or } T_R) \vee (F_A \text{ or } F_R) \text{ (i.e. Ignorance)}$.
 All are symbols.

b) But if $T_A, T_R, F_A, F_R, U, C, G$ are subsets of $[0, 1]$, then we get a *Seven Numerical-Valued Neutrosophic Logic*.

6. n-Valued Logic

a) *The n-Symbol-Valued Refined Neutrosophic Logic* [Smarandache, 1995].

In general:

T can be split into many types of truths: T_1, T_2, \dots, T_p , and I into many types of indeterminacies: I_1, I_2, \dots, I_r , and F into many types of falsities: F_1, F_2, \dots, F_s , where all $p, r, s \geq 1$ are integers, and $p + r + s = n$.

All subcomponents T_j, I_k, F_l are symbols for $j \in \{1, 2, \dots, p\}$, $k \in \{1, 2, \dots, r\}$, and $l \in \{1, 2, \dots, s\}$.
 If at least one $I_k = T_j \wedge F_l = \text{contradiction}$, we get again the Extenics.

b) *The n-Numerical-Valued Refined Neutrosophic Logic*.

In the same way, but all subcomponents T_j, I_k, F_l are not symbols, but subsets of $[0, 1]$, for all

$j \in \{1, 2, \dots, p\}$, all $k \in \{1, 2, \dots, r\}$, and all $l \in \{1, 2, \dots, s\}$.

If all sources of information that separately provide neutrosophic values for a specific subcomponent are independent sources, then in the general case we consider that each of the subcomponents T_j, I_k, F_l is independent with respect to the others and it is in the non-standard set $]0, I^+[$. Therefore per total we have for crisp neutrosophic value subcomponents T_j, I_k, F_l that:

$$]0 \leq \sum_{j=1}^p T_j + \sum_{k=1}^r I_k + \sum_{l=1}^s F_l \leq n^+ \quad (1)$$

where of course $n = p + r + s$ as above.

If there are some dependent sources (or respectively some dependent subcomponents), we can treat those dependent subcomponents together. For example, if T_2 and I_3 are dependent, we put them together as $]0 \leq T_2 + I_3 \leq I^+$.

The non-standard unit interval $]0, I^+[$, used to make a distinction between *absolute* and *relative* truth/indeterminacy/falsehood in philosophical applications, is replaced for simplicity with the standard (classical) unit interval $[0, I]$ for technical applications.

For at least one $I_k = T_j \wedge F_l = \text{contradiction}$, we get again the Extenics.

7. Neutrosophic Cube and its Extenics Part

The most important distinction between IFS and NS is showed in the below **Neutrosophic Cube** A'B'C'D'E'F'G'H' introduced by J. Dezert in 2002.

Because in technical applications only the classical interval is used as range for the neutrosophic parameters, we call the cube the **technical neutrosophic cube** and its extension the **neutrosophic cube** (or **absolute neutrosophic cube**), used in the fields where we need to differentiate between *absolute* and *relative* (as in philosophy) notions.

Another source which is capable to find only the degree of non-membership of an element;

Or a source which only computes the indeterminacy.

Thus, when we put the results together of these sources, it is possible that their sum is not 1, but smaller (Incomplete) or greater (Extenics).

8. Example of Extenics in 3-Valued Neutrosophic Logic

About a proposition P, the first source of information provides the truth-value $T=0.8$.

Second source of information provides the false-value $F=0.7$.

Third source of information provides the indeterminacy-value $I=0.2$.

Hence $NL_3(P) = (0.8, 0.2, 0.7)$.

Got Extenics, since Contradiction: $T + F = 0.8 + 0.7 > 1$.

Can remove Contradiction by normalization:

$NL(P) = (0.47, 0.12, 0.41)$; now $T+F \leq 1$.

9. Example of Extenics in 4-Valued Neutrosophic Logic

About a proposition Q, the first source of information provides the truth-value $T=0.4$, second source provides the false-value $F=0.3$, third source provides the undefined-value $U=0.1$, fourth source provides the contradiction-value $C=0.2$.

Hence $NL_4(Q) = (0.4, 0.1, 0.2, 0.3)$.

Got Extenics, since Contradiction $C = 0.2 > 0$.

Since $C = T \wedge F$, we can remove it by transferring its value 0.2 to T and F (since T and F were involved in the conflict) proportionally w.r.t. their values 0.4, 0.3:

$xT/0.4 = yF/0.3 = 0.2/(0.4+0.3)$, whence $xT=0.11$, $yF=0.09$.

Thus $T=0.4+0.11=0.51$, $F=0.3+0.09=0.39$, $U=0.1$, $C=0$.

Conclusion

Many types of logics have been presented above related with Extenics. Examples of Neutrosophic Cube and its Extenics part, and Extenics in 3-Valued and 4-Valued Neutrosophic Logics are given.

Similar generalizations are done for **n-Valued Refined Neutrosophic Set**, and respectively **n-Valued Refined Neutrosophic Probability** in connections with Extensions.

References

Cai Wen. *Extension Set and Non-Compatible Problems* [J]. Journal of Scientific Exploration, 1983, (1): 83-97.

Yang Chunyan, Cai Wen. *Extension Engineering* [M]. Beijing: Science Press, 2007.

Didier Dubois, Uncertainty Theories, Degrees of Truth and Epistemic States, http://www.icaart.org/Documents/Previous_Invited_Speakers/2011/ICAART2011_Dubois.pdf

Florentin Smarandache, editor, Proceedings of the Introduction to Neutrosophic Physics: Unmatter & Unparticle - International Conference, Zip Publ., Columbus, 2011.

Dmitri Rabounski, Florentin Smarandache, Larissa Borisova, Neutrosophic Methods in General Relativity. Neutrosophic Book Series, 10. *Hexis, Phoenix, AZ*, 2005. 78 pp.

Dmirii Rabunskii, Florentin Smarandache, Larissa Borisova, \cyr Neitrososfskie metody v obshchei teorii otноситel'nosti. (Russian) [Neutrosophic Methods in the General Theory of Relativity] Translated from the 2005 English edition [Hexis, Phoenix, AZ]. *Hexis, Phoenix, AZ*, 2005. 105 pp.

F. Smarandache, Neutrosophic Logic and Set, mss., <http://fs.gallup.unm.edu/neutrosophy.htm>, 1995.

Florentin Smarandache, *A Unifying Field in Logics: Neutrosophic Field*, Multiple-Valued Logic / An International Journal, Vol. 8, No. 3, 385-438, June 2002.

Umberto Rivieccio, *Neutrosophic logics: Prospects and problems*, Fuzzy Sets and Systems, Vol. 159, Issue 14, pp. 1860 – 1868, 2008.

Florentin Smarandache, *An Introduction to the Neutrosophic Probability Applied in Quantum Statistics*, <Bulletin of Pure and Applied Sciences>, Physics, 13-25, Vol. 22D, No. 1, 2003.

F. Smarandache, Neutrosophic Set — A Generalization of the Intuitionistic Fuzzy Set. *Int. J. Pure Appl. Math.* 24 (2005), No. 3, 287-297.

Florentin Smarandache, *A Unifying Field in Logics: Neutrosophic Logic*, <Multiple Valued Logic / An International Journal>, USA, ISSN 1023-6627, Vol. 8, No. 3, pp. 385-438, 2002. The whole issue of this journal is dedicated to Neutrosophy and Neutrosophic Logic.

J. Dezert, Open questions on neutrosophic inference. Neutrosophy and neutrosophic logic. *Mult.-Valued Log.* 8 (2002), no. 3, 439--472.

Webster's Online Dictionary, *Paraconsistent probability* {neutrosophic probability},

<http://www.websters-online-dictionary.org/definitions/Paraconsistent?cx=partner-pub-0939450753529744%3Av0qd01-tdlq&cof=FORID>