

## The continuum hypothesis

Daniel Cordero Grau  
dcgrau01@yahoo.co.uk

In this paper we prove of the continuum hypothesis, by proving that the theory of initial ordinals and the theory of cardinals are equivalent. To prove that the theorems of the theory of cardinals are theorems of the theory of initial ordinals, and that, conversely, the theorems of the theory of initial ordinals are theorems of the theory of cardinals, we use the definition of an isomorphism of theories in mathematical logic, in its equivalent form, the definition of an isomorphism of categories from the theory of categories, and also, we use the definition of a functor, the definition of a category, the axioms of mathematical logic and the axioms of the theory of categories, which include the Gödel-Bernays-von Neumann axioms for classes and sets, and so, applying both the theorem of comparability of ordinals to the theory of cardinals, and the fundamental theorem of cardinal arithmetic to the theory of ordinals, we prove the theorem.

Theorem "generalized continuum hypothesis": For every transfinite cardinal number  $\alpha$ , there is no cardinal number between  $\alpha$  and  $2^\alpha$ .

Proof: Let **Card** be the class of cardinals and let **Ord** be class of initial ordinals. Since, according to the definition of a category, every well order is a category, and the classes **Card** and **Ord** are well orders, **Card** and **Ord** are categories. We prove that **Card** and **Ord** are isomorphic, proving that there is a full and faithful functor  $T: \mathbf{Card} \rightarrow \mathbf{Ord}$  such that each initial ordinal  $\beta$  is isomorphic to an initial ordinal  $T\alpha$  for some cardinal  $\alpha$ .

Let  $T: \mathbf{Card} \rightarrow \mathbf{Ord}$  be the function of categories which assigns to every cardinal  $\alpha$  the initial ordinal  $T\alpha$  of its equipotence class,  $\alpha \mapsto T\alpha$ , and to every arrow  $f: \alpha \rightarrow \alpha'$  in **Card** the arrow  $Tf: T\alpha \rightarrow T\alpha'$  in **Ord**,  $f \mapsto Tf$ , for each pair of cardinals  $\alpha$  and  $\alpha'$ . The function  $T$  of categories is well-defined because each cardinal  $\alpha$  lies in a unique equipotence class defined by  $\alpha$ , and so, it defines uniquely  $T\alpha$ , and because there is only one arrow  $Tf$  in **Ord** for every arrow  $f$  in **Card**, since each arrow  $f$  in a category  $C$  is a pair of objects  $\alpha$  and  $\alpha'$  for which  $f: \alpha \rightarrow \alpha'$  is an arrow in  $C$ , to each pair of cardinals  $\alpha$  and  $\alpha'$  there is a unique pair of initial ordinals  $T\alpha$  and  $T\alpha'$ , which are isomorphic to  $\alpha$  and  $\alpha'$ , respectively, by definition of  $T$ , and so, by the orderings in **Card** and **Ord**, for which the arrow  $f: \alpha \rightarrow \alpha'$  is in **Card** if, and only if, the arrow  $g = Tf: T\alpha \rightarrow T\alpha'$  is in **Ord**, which is unique, for **Card** and **Ord** are preorders. The latter condition on  $T$  means that  $T$  is an order-preserving function of the linear order **Card** to the linear order **Ord**.

The function  $T$  of categories is a functor, because a functor is a function of categories preserving monoids, that is, preserving identities and composable pair of arrows,  $T1_\alpha = 1_{T\alpha}$  and  $T(f \circ g) = Tf \circ Tg$ , for every identity  $1_\alpha$  and every composable pair of arrows  $f$  and  $g$  in the function domain category. For, each identity  $1_\alpha$  in **Card** is a cardinal  $\alpha$ , the initial ordinal  $T\alpha$  of the equipotence class of each cardinal  $\alpha$  is the identity  $1_{T\alpha}$  in **Ord**, every category has all its identities, there is an identity  $1_\alpha$  for each object  $\alpha$ , and **Ord** is a category. And for, each arrow  $f \circ g$  is a composable pair of arrows  $f$  and  $g$ , each composable pair of arrows  $f$  and  $g$  is a triad of objects  $\alpha$ ,  $\alpha'$  and  $\alpha''$  such that  $f: \alpha \rightarrow \alpha'$ ,  $g: \alpha' \rightarrow \alpha''$  and  $f \circ g: \alpha \rightarrow \alpha''$  are arrows,  $Tf: T\alpha \rightarrow T\alpha'$ ,  $Tg: T\alpha' \rightarrow T\alpha''$  and  $T(f \circ g): T\alpha \rightarrow T\alpha''$  are arrows in **Ord**, by definition of  $T$ ,  $Tf$  and  $Tg$  is a composable pair of arrows in **Ord**, every category has all its composable pairs of arrows, the arrows in a preorder are unique, and **Ord** is a preorder.

The functor  $T$  is full, because to every pair of cardinals  $\alpha$  and  $\alpha'$  and to every arrow  $g: T\alpha \rightarrow T\alpha'$  in **Ord** there is an arrow  $f: \alpha \rightarrow \alpha'$  in **Card** such that  $Tf = g$ , because **Ord** is a preorder and  $T$  satisfies the condition above: the arrow  $f: \alpha \rightarrow \alpha'$  is in **Card** if, and only if, the arrow  $Tf: T\alpha \rightarrow T\alpha'$  is in **Ord**. The functor  $T$  is faithful, because to every pair of cardinals  $\alpha$  and  $\alpha'$  and to every pair of arrows  $f_1, f_2: \alpha \rightarrow \alpha'$  in **Card** the equality  $Tf_1 = Tf_2$  implies  $f_1 = f_2$ , because **Card** is a preorder, and by definition of  $T$ . Finally, since every cardinal  $|\beta|$  is isomorphic to the initial ordinal  $T|\beta|$  of its equipotence class, and since every initial ordinal  $\beta$  is isomorphic to the cardinal  $|\beta|$  of its equipotence class by definition of initial ordinal, cardinal and  $T$ , to each initial ordinal  $\beta$  there is an order-preserving isomorphism between  $\beta$  and the initial ordinal  $T|\beta|$  of the equipotence class of  $\beta$ , that is,  $\beta \cong T|\beta|$ , therefore, **Card**  $\cong$  **Ord**.

Thus, for isomorphic categories are isomorphic theories, by the definition of an isomorphism of theories, by the definition of an isomorphism of categories, by the axioms of mathematical logic, and by the axioms of the theory of categories, which include the Gödel-Bernays-von Neumann axioms, and both since there is no initial ordinal between  $\omega$  and  $\omega^\omega$ , by the theorem on the comparability of ordinals, and since  $\aleph_0^{\aleph_0} = 2^{\aleph_0}$ , by the fundamental theorem of cardinal arithmetic, the isomorphism between categories **Card** and **Ord** proves that there is no cardinal number between the transfinite cardinal numbers  $\aleph_0$  and  $2^{\aleph_0}$ , and that, in general, there is no cardinal number between any transfinite cardinal number  $\alpha$  and  $2^\alpha$ . As a consequence, there exist no inaccessible cardinals. In fact, the class of transfinite cardinals is isomorphic to  $\omega$ , because the order-preserving function  $f$  of  $\omega$  to it, which assigns to each finite ordinal  $\alpha$  the  $\alpha$ -th transfinite cardinal is an isomorphism, which is unique, by transfinite construction.

### The theorem in universal algebra

Thus, not only does the theorem prove that the class of cardinals is an infinite countable nondiscrete large category which is a closed complete and cocomplete semiring, with arrows, the polynomial maps and the exponential maps, that is an algebra by the action of the covariant exponential functor semiring  $e$ , itself, a functor algebra, but also, that the closed complete and cocomplete algebra of initial ordinals **Ord** is isomorphic to the closed complete and cocomplete algebra of cardinals **Card**.

### The theorem in categorical logic

In categorical logic, as all first order theories are infinite well orders isomorphic to  $\omega$  and have thereby cardinal smaller than the cardinal of the continuum, not only does the theorem proves that the categorical theories **Card** and **Ord** are equivalent, but also, that all higher order theories are continuums, for they are partial orders isomorphic to infinite countable products of first order theories.

### The theorem in topos theory

In topos theory, not only does the theorem prove that the category **Card** of cardinals is a topos, but also, that its topos of sheaves is the category **Sets**<sup>**Card**\*</sup>, denoting the dual category of the category **Card** by **Card**<sup>\*</sup>, of the set-valued contravariant functors on **Card** to **Sets** which assign to every cardinal number  $\beta$  its set of cardinal functions on  $\beta$  all of which turn out to be the cardinal continuous functions on the topology of the cardinals.

### Bibliography

- Alexandroff, Hopf, *Topologie*, Berlin: Springer 2013
- Cohen, *Set theory and the continuum hypothesis*, New York: Dover 2008
- Cohn, *Universal algebra*, Berlin: Springer 1981
- Dugundji, *Topology*, Iowa: William C Brown 1989
- Ebbinghaus, Flum, *Mathematical logic*, Berlin: Springer 1996
- Eilenberg, Steenrod, *Foundations of algebraic topology*, St Louis: Nabu 2011
- Gödel, *The consistency of the continuum hypothesis*, Tokyo: Ishi 2009
- Grätzer, *Universal algebra*, Berlin: Springer 2008
- Jänich, *Topologie*, Berlin: Springer 2008
- Johnstone, *Topos theory*, New York: Dover 2014
- Kelley, *General topology*, Berlin: Springer 2008
- Lambeck, Scott, *Introduction to higher order categorical logic*, Cambridge: Cambridge University Press 1986
- Mac Lane, *Categories for the working mathematician*, Berlin: Springer 1998