

# Generalized Neutrosophic Set and Generalized Neutrosophic Topological Spaces

<sup>1</sup>A. A. Salama, <sup>2</sup>S. A. Alblowi

<sup>1</sup>Egypt, Port Said University, Faculty of Sciences Department of Mathematics and Computer Science  
<sup>2</sup>Department of Mathematics, King Abdulaziz University, Gedh, Saudi Arabia,

**Abstract:** In this paper we introduce definitions of generalized neutrosophic sets. After given the fundamental definitions of generalized neutrosophic set operations, we obtain several properties, and discussed the relationship between generalized neutrosophic sets and others. Finally, we extend the concepts of neutrosophic topological space [9], intuitionistic fuzzy topological space [5, 6], and fuzzy topological space [4] to the case of generalized neutrosophic sets. Possible application to GIS topology rules are touched upon.

**Keywords:** Neutrosophic Set; Generalized Neutrosophic Set,; Neutrosophic Topology

## 1. Introduction

Neutrosophy has laid the foundation for a whole family of new mathematical theories generalizing both their classical and fuzzy counterparts, such as a neutrosophic set theory. The fuzzy set was introduced by Zadeh [10] in 1965, where each element had a degree of membership. The intuitionistic fuzzy set (Ifs for short) on a universe  $X$  was introduced by K. Atanassov [1, 2, 3] in 1983 as a generalization of fuzzy set, where besides the degree of membership and the degree of non-membership of each element. After the introduction of the neutrosophic set concept [7, 8, 9]. In this paper we introduce definitions of generalized neutrosophic sets. After given the fundamental definitions of generalized neutrosophic set operations, we obtain several properties, and discussed the relationship between generalized neutrosophic sets and others. Finally, we extend the concepts of neutrosophic topological space [9].

## 2. Terminologies

We recollect some relevant basic preliminaries, and in particular, the work of Smarandache in [7, 8], Atanassov in [1, 2, 3] and Salama [9]. Smarandache introduced the neutrosophic components  $T, I, F$  which represent the membership, indeterminacy, and non-membership values respectively, where  $]0^-, 1^+[$  is nonstandard unit interval.

**Definition.[7, 8]**

Let  $T, I, F$  be real standard or nonstandard subsets of  $]0^-, 1^+[$ , with

$$\begin{aligned} \text{Sup}_T &= t_{\text{sup}}, \text{inf}_T = t_{\text{inf}} \\ \text{Sup}_I &= i_{\text{sup}}, \text{inf}_I = i_{\text{inf}} \\ \text{Sup}_F &= f_{\text{sup}}, \text{inf}_F = f_{\text{inf}} \\ n\text{-sup} &= t_{\text{sup}} + i_{\text{sup}} + f_{\text{sup}} \\ n\text{-inf} &= t_{\text{inf}} + i_{\text{inf}} + f_{\text{inf}}, \end{aligned}$$

$T, I, F$  are called neutrosophic components

**Definition [9]**

Let  $X$  be a non-empty fixed set. A neutrosophic set ( $NS$  for short)  $A$  is an object having the form

$$A = \{ \langle x, \mu_A(x), \sigma_A(x), \gamma_A(x) \rangle : x \in X \}$$

Where  $\mu_A(x), \sigma_A(x)$  and  $\gamma_A(x)$  which represent the degree of membership function (namely  $\mu_A(x)$ ), the degree of indeterminacy (namely  $\sigma_A(x)$ ), and the degree of non-membership (namely  $\gamma_A(x)$ ) respectively of each element  $x \in X$  to the set  $A$ .

**Definition [9].** The NSS  $0_N$  and  $1_N$  in  $X$  as follows:

$0_N$  may be defined as:

$$\begin{aligned} (0_1) \quad 0_N &= \{ \langle x, 0, 0, 1 \rangle : x \in X \} \\ (0_2) \quad 0_N &= \{ \langle x, 0, 1, 1 \rangle : x \in X \} \\ (0_3) \quad 0_N &= \{ \langle x, 0, 1, 0 \rangle : x \in X \} \\ (0_4) \quad 0_N &= \{ \langle x, 0, 0, 0 \rangle : x \in X \} \end{aligned}$$

$1_N$  may be defined as:

$$\begin{aligned} (1_1) \quad 1_N &= \{ \langle x, 1, 0, 0 \rangle : x \in X \} \\ (1_2) \quad 1_N &= \{ \langle x, 1, 0, 1 \rangle : x \in X \} \end{aligned}$$

\* Corresponding author:  
 drsalama44@gmail.com (A. A. Salama)

Published online at <http://journal.sapub.org/computer>

Copyright © 2012 Scientific & Academic Publishing. All Rights Reserved

$$(1_3) \quad 1_N = \{ \langle x, 1, 1, 0 \rangle : x \in X \}$$

$$(1_4) \quad 1_N = \{ \langle x, 1, 1, 1 \rangle : x \in X \}$$

### 3. Generalized Neutrosophic Sets

We shall now consider some possible definitions for basic concepts of the generalized neutrosophic set.

#### Definition

Let  $X$  be a non-empty fixed set. A generalized neutrosophic set (GNS for short)  $A$  is an object having the form  $A = \{ \langle x, \mu_A(x), \sigma_A(x), \gamma_A(x) \rangle : x \in X \}$  Where  $\mu_A(x)$ ,  $\sigma_A(x)$  and  $\gamma_A(x)$  which represent the degree of membership function (namely  $\mu_A(x)$ ), the degree of indeterminacy (namely  $\sigma_A(x)$ ), and the degree of non-membership (namely  $\gamma_A(x)$ ) respectively of each element  $x \in X$  to the set  $A$  where the functions satisfy the condition  $\mu_A(x) \wedge \sigma_A(x) \wedge \nu_A(x) \leq 0.5$ .

#### Remark

A generalized neutrosophic  $A = \{ \langle x, \mu_A(x), \sigma_A(x), \gamma_A(x) \rangle : x \in X \}$  can be identified to an ordered triple  $\langle \mu_A, \sigma_A, \gamma_A \rangle$  in  $]0, 1[$  on  $X$ , where the triple functions satisfy the condition  $\mu_A(x) \wedge \sigma_A(x) \wedge \nu_A(x) \leq 0.5$

#### Remark

For the sake of simplicity, we shall use the symbol  $A = \langle x, \mu_A, \sigma_A, \gamma_A \rangle$  for the

$$GNS \quad A = \{ \langle x, \mu_A(x), \sigma_A(x), \gamma_A(x) \rangle : x \in X \}$$

#### Example

Every GIFS  $A$  a non-empty set  $X$  is obviously on GNS having the form

$$A = \{ \langle x, \mu_A(x), 1 - (\mu_A(x) + \gamma_A(x)), \gamma_A(x) \rangle : x \in X \}$$

#### Definition

Let  $A = \langle \mu_A, \sigma_A, \gamma_A \rangle$  a GNSS on  $X$ , then the complement of the set  $A$  ( $C(A)$ , for short) maybe defined as three kinds of complements

$$(C_1) \quad C(A) = \{ \langle x, 1 - \mu_A(x), \sigma_A(x), 1 - \nu_A(x) \rangle : x \in X \},$$

$$(C_2) \quad C(A) = \{ \langle x, \gamma_A, \sigma_A(x), \mu_A(x) \rangle : x \in X \}$$

$$(C_3) \quad C(A) = \{ \langle x, \gamma_A, 1 - \sigma_A(x), \mu_A(x) \rangle : x \in X \}$$

One can define several relations and operations between GNSS as follows:

#### Definition

Let  $X$  be a non-empty set, and GNSS  $A$  and  $B$  in the form  $A = \langle x, \mu_A(x), \sigma_A(x), \gamma_A(x) \rangle$ ,

$B = \langle x, \mu_B(x), \sigma_B(x), \gamma_B(x) \rangle$ , then we may consider two possible definitions for subsets ( $A \subseteq B$ )

( $A \subseteq B$ ) may be defined as

$$A \subseteq B \Leftrightarrow \mu_A(x) \leq \mu_B(x), \gamma_A(x) \geq \gamma_B(x) \text{ and } \sigma_A(x) \leq \sigma_B(x)$$

$\forall x \in X$

$$A \subseteq B \Leftrightarrow \mu_A(x) \leq \mu_B(x), \gamma_A(x) \geq \gamma_B(x) \text{ and } \sigma_A(x) \geq \sigma_B(x)$$

#### Proposition

For any generalized neutrosophic set  $A$  the following are holds

$$0_N \subseteq A, \quad 0_N \subseteq 0_N$$

$$A \subseteq 1_N, \quad 1_N \subseteq 1_N$$

#### Definition

Let  $X$  be a non-empty set, and  $A = \langle x, \mu_A(x), \gamma_A(x), \sigma_A(x) \rangle$ ,

$B = \langle x, \mu_B(x), \sigma_B(x), \gamma_B(x) \rangle$  are GNSS Then

$A \cap B$  maybe defined as:

$$(I_1) \quad A \cap B = \langle x, \mu_A(x), \mu_B(x), \sigma_A(x), \sigma_B(x), \gamma_A(x), \gamma_B(x) \rangle$$

$$(I_2) \quad A \cap B = \langle x, \mu_A(x) \wedge \mu_B(x), \sigma_A(x) \wedge \sigma_B(x), \gamma_A(x) \vee \gamma_B(x) \rangle$$

$$(I_3) \quad A \cap B = \langle x, \mu_A(x) \wedge \mu_B(x), \sigma_A(x) \vee \sigma_B(x), \gamma_A(x) \vee \gamma_B(x) \rangle$$

$A \cup B$  may be defined as:

$$(U_1) \quad A \cup B = \langle x, \mu_A(x) \vee \mu_B(x), \sigma_A(x) \vee \sigma_B(x), \gamma_A(x) \wedge \gamma_B(x) \rangle$$

$$(U_2) \quad A \cup B = \langle x, \mu_A(x) \vee \mu_B(x), \sigma_A(x) \wedge \sigma_B(x), \gamma_A(x) \wedge \gamma_B(x) \rangle$$

$$[ ]A = \langle x, \mu_A(x), \sigma_A(x), 1 - \mu_A(x) \rangle$$

$$\langle \rangle A = \langle x, 1 - \gamma_A(x), \sigma_A(x), \gamma_A(x) \rangle$$

**Example.3.2.** Let  $X = \{a, b, c, d, e\}$  and

$A = \langle x, \mu_A, \sigma_A, \nu_A \rangle$  given by:

X	$\mu_A(x)$	$\nu_A(x)$	$\sigma_A(x)$	$\mu_A(x) \wedge \nu_A(x) \wedge \sigma_A(x)$
a	0.6	0.3	0.5	0.3
b	0.5	0.3	0.6	0.3
c	0.4	0.4	0.5	0.4
d	0.3	0.5	0.3	0.3
e	0.3	0.6	0.4	0.3

Then the family  $G = \{O_{\sim}, A\}$  is an GNSS on  $X$ .

We can easily generalize the operations of generalized intersection and union in definition 3.4 to arbitrary family of GNSS as follow:

#### Definition

Let  $\{A_j : j \in J\}$  be a arbitrary family of NSS in  $X$ , then

$\cap A_j$  maybe defined as:

$$1) \quad \cap A_j = \left\langle x, \bigwedge_{j \in J} \mu_{A_j}(x), \bigwedge_{j \in J} \sigma_{A_j}(x), \bigvee_{j \in J} \gamma_{A_j}(x) \right\rangle$$

$$2) \quad \cap A_j = \left\langle x, \bigwedge_{j \in J} \mu_{A_j}(x), \bigvee_{j \in J} \sigma_{A_j}(x), \bigvee_{j \in J} \gamma_{A_j}(x) \right\rangle$$

$\cup A_j$  maybe defined as:

$$1) \cup A_j = \left\langle x, \bigvee_{j \in J} \mu_{A_j}, \wedge \sigma_{A_j}, \wedge \nu_{A_j} \right\rangle$$

$$2) \cup A_j = \left\langle x, \bigvee_{j \in J} \mu_{A_j}, \bigvee \sigma_{A_j}, \wedge \nu_{A_j} \right\rangle$$

**Definition**

Let  $A$  and  $B$  are generalized neutrosophic sets then  $A|B$  may be defined as

$$A|B = \langle x, \mu_A \wedge \gamma_B, \sigma_A(x) \sigma_B(x), \gamma_A \vee \mu_B(x) \rangle$$

**Proposition**

For all  $A, B$  two generalized neutrosophic sets then the following are true

i)  $C(A \cap B) = C(A) \cup C(B)$

ii)  $C(A \cup B) = C(A) \cap C(B)$

### 4. Generalized Neutrosophic Topological Spaces

Here we extend the concepts of and intuitionistic fuzzy topological space [5, 7], and neutrosophic topological Space [9] to the case of generalized neutrosophic sets.

**Definition**

A generalized neutrosophic topology (GNT for short) on a non empty set  $X$  is a family  $\tau$  of generalized neutrosophic subsets in  $X$  satisfying the following axioms

$$(GNT_1) O_N, 1_N \in \tau,$$

$$(GNT_2) G_1 \cap G_2 \in \tau \text{ for any } G_1, G_2 \in \tau,$$

$$(GNT_3) \bigcup G_i \in \tau \quad \forall \{G_i : i \in J\} \subseteq \tau$$

In this case the pair  $(X, \tau)$  is called a generalized neutrosophic topological space (GNTS for short) and any neutrosophic set in  $\tau$  is known as neutrosophic open set (NOS for short) in  $X$ . The elements of  $\tau$  are called open generalized neutrosophic sets, A generalized neutrosophic set  $F$  is closed if and only if  $C(F)$  is generalized neutrosophic open.

**Remark** A generalized neutrosophic topological spaces are very natural generalizations of intuitionistic fuzzy topological spaces allow more general functions to be members of intuitionistic fuzzy topology.

**Example**

Let  $X = \{x\}$  and

$$A = \{ \langle x, 0.5, 0.5, 0.4 \rangle : x \in X \}$$

$$B = \{ \langle x, 0.4, 0.6, 0.8 \rangle : x \in X \}$$

$$D = \{ \langle x, 0.5, 0.6, 0.4 \rangle : x \in X \}$$

$$C = \{ \langle x, 0.4, 0.5, 0.8 \rangle : x \in X \}$$

Then the family  $\tau = \{O_n, 1_n, A, B, C, D\}$  of GNSs in  $X$  is generalized neutrosophic topology on  $X$

**Example**

Let  $(X, \tau_0)$  be a fuzzy topological space in Changes [4] sense such that  $\tau_0$  is not indiscrete suppose now that

$\tau_0 = \{0_N, 1_N\} \cup \{V_j : j \in J\}$  then we can construct two GNTSS on  $X$  as follows

$$\tau_{0,1} = \{0_N, 1_N\} \cup \{ \langle x, V_j, \sigma(x), 0 \rangle : j \in J \}$$

$$\tau_{0,2} = \{0_N, 1_N\} \cup \{ \langle x, V_j, 0, \sigma(x), 1 - V_j \rangle : j \in J \}$$

**Proposition**

Let  $(X, \tau)$  be a GNT on  $X$ , then we can also construct several GNTSS on  $X$  in the following way:

a)  $\tau_{0,1} = \{ [ ]G : G \in \tau \}$ ,

b)  $\tau_{0,2} = \{ \langle \rangle G : G \in \tau \}$ ,

**Proof a)**

$(GNT_1)$  and  $(GNT_2)$  are easy.

$(GNT_3)$  Let  $\{ [ ]G_j : j \in J, G_j \in \tau \} \subseteq \tau_{0,1}$ . Since

$$\cup G_j = \left\langle x, \bigvee \mu_{G_j}, \bigvee \sigma_{G_j}, \wedge \gamma_{G_j} \right\rangle \text{ or } \left\langle x, \bigvee \mu_{G_j}, \wedge \sigma_{G_j}, \wedge \gamma_{G_j} \right\rangle \text{ or } \left\langle x, \bigvee \mu_{G_j}, \wedge \sigma_{G_j}, \bigvee \gamma_{G_j} \right\rangle \in \tau,$$

we have

$$\cup [ ]G_j = \left\langle x, \bigvee \mu_{G_j}, \bigvee \sigma_{G_j}, \wedge (1 - \mu_{G_j}) \right\rangle \text{ or } \left\langle x, \bigvee \mu_{G_j}, \bigvee \sigma_{G_j}, (1 - \bigvee \mu_{G_j}) \right\rangle \in \tau_{0,1}$$

This similar to (a)

**Definition**

Let  $(X, \tau_1), (X, \tau_2)$  be two generalized neutrosophic topological spaces on  $X$ . Then  $\tau_1$  is said be contained in  $\tau_2$  (in symbols  $\tau_1 \subseteq \tau_2$ ) if  $G \in \tau_2$  for each  $G \in \tau_1$ . In this case, we also say that  $\tau_1$  is coarser than  $\tau_2$ .

**Proposition**

Let  $\{ \tau_j : j \in J \}$  be a family of NTSS on  $X$ . Then  $\bigcap \tau_j$  is A generalized neutrosophic topology on  $X$ . Furthermore,  $\bigcap \tau_j$  is the coarsest NT on  $X$  containing all.  $\tau_j$ ,<sup>s</sup>

**Proof.** Obvious

**Definition**

The complement of  $A$  ( $C(A)$  for short) of NOS.  $A$  is called a generalized neutrosophic closed set (GNCS for short) in  $X$ .

Now, we define generalized neutrosophic closure and interior operations in generalized neutrosophic topological spaces:

**Definition**

Let  $(X, \tau)$  be GNTS and  $A = \langle x, \mu_A(x), \gamma_A(x), \sigma_A(x) \rangle$  be a GNS in  $X$ .

Then the generalized neutrosophic closer and generalized neutrosophic interior of A are defined by

$$GNCl(A) = \bigcap \{ K : K \text{ is an NCS in } X \text{ and } A \subseteq K \}$$

$GNInt(A) = \bigcup \{ G : G \text{ is an NOS in } X \text{ and } G \subseteq A \}$ . It can be also shown that

It can be also shown that  $GNCl(A)$  is GNCS and  $GNInt(A)$  is a GNOS in  $X$

$A$  is in  $X$  if and only if  $GNCl(A)$ .

$A$  is  $GNCS$  in  $X$  if and only if  $GNInt(A) = A$ .

**Proposition**

For any generalized neutrosophic set  $A$  in  $(X, \tau)$  we have

(a)  $GNCl(C(A)) = C(GNInt(A))$ ,

(b)  $GNInt(C(A)) = C(GNCl(A))$ .

**Proof.**

Let  $A = \{ \langle x, \mu_A, \sigma_A, \nu_A \rangle : x \in X \}$  and suppose that the family of generalized neutrosophic subsets contained in  $A$  are indexed by the family if  $GNSS$  contained in  $A$  are indexed by the

family  $A = \{ \langle x, \mu_{G_i}, \sigma_{G_i}, \nu_{G_i} \rangle : i \in J \}$ . Then we see

that  $GNInt(A) = \{ \langle x, \vee \mu_{G_i}, \vee \sigma_{G_i}, \wedge \nu_{G_i} \rangle \}$  and

hence  $C(GNInt(A)) = \{ \langle x, \wedge \mu_{G_i}, \vee \sigma_{G_i}, \vee \nu_{G_i} \rangle \}$ .

Since  $C(A)$  and  $\mu_{G_i} \leq \mu_A$  and  $\nu_{G_i} \geq \nu_A$  for each  $i \in J$ , we obtaining  $C(A)$ . i.e

$GNCl(C(A)) = \{ \langle x, \wedge \nu_{G_i}, \vee \sigma_{G_i}, \vee \mu_{G_i} \rangle \}$ . Hence

$GNCl(C(A)) = C(GNInt(A))$ , follows immediately

This is analogous to (a).

**Proposition**

Let  $(X, \tau)$  be a  $GNTS$  and  $A, B$  be two neutrosophic sets in  $X$ . Then the following properties hold:

$GNInt(A) \subseteq A$ ,

$A \subseteq GNCl(A)$ ,

$A \subseteq B \Rightarrow GNInt(A) \subseteq GNInt(B)$ ,

$A \subseteq B \Rightarrow GNCl(A) \subseteq GNCl(B)$ ,

$GNInt(GNInt(A)) = GNInt(A) \wedge GNInt(B)$ ,

$GNCl(A \cup B) = GNCl(A) \vee GNCl(B)$ ,

$GNInt(1_N) = 1_N$ ,

$GNCl(0_N) = 0_N$ ,

**Proof** (a), (b) and (e) are obvious (c) follows from (a) and Definitions.

---

## REFERENCES

- [1] K. Atanassov, intuitionistic fuzzy sets, in V.Sgurev, ed., Vii ITKRS Session, Sofia(June 1983 central Sci. and Techn. Library, Bulg. Academy of Sciences( 1984)).
- [2] K. Atanassov, intuitionistic fuzzy sets, Fuzzy Sets and Systems 20(1986)87-96.
- [3] K. Atanassov, Review and new result on intuitionistic fuzzy sets , preprint IM-MFAIS-1-88, Sofia, 1988.
- [4] C.L. Chang, Fuzzy Topological Spaces, J. Math. Anal. Appl. 24 (1968)182-1 90.
- [5] Dogan Coker, An introduction to intuitionistic fuzzy topological spaces, Fuzzy Sets and Systems. 88(1997)81-89.
- [6] Reza Saadati, Jin HanPark, On the intuitionistic fuzzy topological space, Chaos, Solitons and Fractals 27(2006)331-344 .
- [7] Florentin Smarandache , Neutrosophy and Neutrosophic Logic , First International Conference on Neutrosophy , Neutrosophic Logic , Set, Probability, and Statistics University of New Mexico, Gallup, NM 87301, USA(2002) , smarand@unm.edu
- [8] F. Smarandache. A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability. American Research Press, Rehoboth, NM, 1999.
- [9] A.A. Salama and S.A. AL-Blowi , NEUTROSOPHIC SET and NEUTROSOPHIC TOPOLOGICAL SPACES, IOSR Journal of Math. ISSN:2278-5728.Vol.(3) ISSUE4PP31-35(2012)
- [10] L.A. Zadeh, Fuzzy Sets, Inform and Control 8(1965)338-353.