General Relativity as curvature of space

Peter H. Michalicka

February 6, 2014

Abstract

With the Planck 'constants' length, time, mass and acceleration will be shown, that a Quantum Gravity of the cosmos exists. This paper shows how Einstein's Field Equations in Friedmann Robertson Walker Metric solves the Planck Era context.

1 The Planck 'constants'

Planck length $\Delta x = \sqrt{\frac{Gh}{c^3}}$

Planck time $\Delta t = \sqrt{\frac{Gh}{c^5}}$

Planck mass $\Delta m = \sqrt{\frac{hc}{G}}$

Planck acceleration $\Delta a = \frac{c}{\Delta t} = \sqrt{\frac{c^7}{hG}}$

2 Modern Cosmology

Within modern cosmology the Einstein's Field Equations would be written with cosmological term Λ as follows (see [2] and [3]):

$$R_{ik} - \frac{1}{2}g_{ik}R - \Lambda g_{ik} = \frac{8\pi G}{c^4}T_{ik} \quad (2.0)$$

The solutions of the Field Equations in Friedmann-Roberson-Walker-Metric are:

FRW Equation (I)

$$\frac{\dot{R}^2}{R^2} = \frac{8\pi G\rho}{3} + \frac{\Lambda c^2}{3} - \frac{kc^2}{R^2} \quad (2.1)$$

FRW Equation (II)

$$\frac{\ddot{R}}{R} = \frac{\Lambda c^2}{3} - \frac{4\pi G}{3} \left(\rho + \frac{3p}{c^2}\right) \quad (2.2)$$

Einstein abandoned the cosmological term Λ as his "greatest blunder" after Hubble's 1928 discovery that the distant galaxies are expanding away from each other. Within a Universe with ideal Quantum gas and without cosmological term Λ and geometry factor k = 0 the equation (2.1) and (2.2) will become:

FRW Equation (I)

$$\frac{\dot{R}^2}{R^2} = \frac{8\pi G\rho}{3} \quad (2.3)$$

and FRW Equation (II)

$$\frac{\ddot{R}}{R} = -\frac{4\pi G}{3} \left(\rho + \frac{3p}{c^2}\right) \quad (2.4)$$

With the relation $p = \frac{\rho c^2}{3}$ (Quantum gas) will change (2.4) as follows: FRW Equation (II)

$$\frac{R}{R} = -\frac{8\pi G\rho}{3} \quad (2.5)$$

Within Thermodynamics we assume dE = TdS - pdV and an adiabatic process it holds TdS = 0. We become $d(\epsilon V) = d\epsilon V + \epsilon dV = -pdV$ and it follows: With $d\epsilon = -(\epsilon + p)\frac{dV}{V}$ and the relation $p = \frac{\epsilon}{3}$ we assume:

$$\frac{d\epsilon}{\epsilon} = -\frac{4dV}{3V} \text{ or } \epsilon \sim V^{-4/3} \sim R^{-4} \quad (2.6)$$

3 The Cosmic Background Radiation

Now we know the energy-density in the Planck-Era $\rho c^2 = \tilde{a}\Delta T^4 = \frac{3c^7}{8\pi hG} (\tilde{a} = Radiation \ constant = 7.5657e^{-16})$, If we assume $c = h = G = k_B = 1$ we get:

$$\frac{3}{8\pi} = \tilde{a}\Delta T^4 = \frac{8\pi^5}{15}T^4 \text{ or } T^4 = \frac{45}{64\pi^6}$$

Now we get the Planck-Temperature $\Delta T = (\frac{45}{64\pi^6})^{1/4} = \frac{1}{6.08088337383}$

In Planck-Era the following 6 relations are valid:

$$\Delta m \ \Delta x = \frac{h}{c} \quad (3.1)$$

$$\Delta m \ \Delta t = \frac{h}{c^2} \quad (3.2)$$

$$\frac{\Delta m}{\Delta a} = \frac{h}{c^3} \quad (3.3)$$

$$\frac{\Delta m}{\Delta x} = \frac{c^2}{G} \quad (3.4)$$

$$\frac{\Delta m}{\Delta t} = \frac{c^3}{G} \quad (3.5)$$

$$\Delta F = \Delta m \ \Delta a = \frac{c^4}{G} \quad (3.6)$$

Now we could calculate the CBR (T_{γ}) as follows:

$$E_{\gamma} = \frac{hc}{x} = 6.08088337383kT_{\gamma} \text{ with } T_{\gamma} = 2.725K$$

We could calculate $m_{\gamma} = 2.5444e^{-39}kg$ and $x = \lambda_{\gamma} = 8.6828e^{-4}m$, also a $t_{\gamma} = \frac{\lambda_{\gamma}}{c} = 2.8963e^{-12}s$.

For the CBR we receive:

$$E_{\gamma} = m_{\gamma}ax = \frac{h\nu}{c^2}\frac{mc^3}{h}\frac{c}{\nu} = m_{\gamma}c^2$$

4 Gravitation as curvature of space

In macroscopic Scale the equation (3.1) til (3.6) will we rewritten as: (Entropy constant $\zeta=\frac{\Delta T}{T_{\gamma}}=2.1432e^{31})$

 $M R = \zeta^4 \frac{h}{c} \quad (4.1)$ $M t = \zeta^4 \frac{h}{c^2} \quad (4.2)$ $\frac{M}{a} = \zeta^4 \frac{h}{c^3} \quad (4.3)$ $\frac{M}{R} = \frac{c^2}{G} \quad (4.4)$ $\frac{M}{t} = \frac{c^3}{G} \quad (4.5)$ $\Delta F = M \ a = \frac{c^4}{G} \quad (4.6)$

With $a = \frac{G M}{R^2} = \frac{M c^3}{\zeta^4 h}$ follows:

$$\frac{G}{R^2} = \frac{c^3}{\zeta^4 h} \quad (4.7)$$

Furthermore we receive from GR:

$$\begin{split} R &= \zeta^2 \; \Delta x => Radius \; of \; Universe \; R = 1.861 e^{28} m \\ \frac{M}{R} &= \frac{c^2}{2G} = \frac{\Delta m}{\Delta x} => Mass \; of \; Universe \; M = 1.253 e^{55} kg \\ \frac{M}{t} &= \frac{c^3}{G} = \frac{\Delta m}{\Delta t} => Age \; of \; Universe \; t = 6.207 e^{19} s \end{split}$$

For $\dot{R}^2 = \frac{2GM}{R}$ is with (4.7): $\dot{R}^2 = 2MR\frac{c^3}{\zeta^4 h} = c^2$

The FRW Gleichung (I) (2.3) is as follows:

$$\frac{c^2}{R^2} = \frac{8\pi G\rho}{3}$$

or

$$\frac{1}{R^4} = \frac{8\pi\rho c^2}{3\zeta^4 hc}$$

We become the \mathbb{R}^4 dependency of (2.6) as follows:

$$\frac{3\zeta^4 hc}{8\pi R^4} = \rho c^2 = \tilde{a}T_\gamma^4$$

5 References

- A.Einstein, Sitz. Preuss. Akad. d. Wiss., Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie (1917)
- 2. V.Sahni, The Case for a Positive Cosmological Λ -Term, astro-ph/9904398
- 3. S.M.Carroll, The Cosmological Constant, astro-ph/0004075