General Relativity as curvature of space

Peter H. Michalicka

Email: Peter.Michalicka@gmx.at

February 17, 2015

Abstract

With the Planck 'constants' length, time, mass and acceleration will be shown, that a Quantum Gravity of the cosmos exists. This paper shows how Einstein's Field Equations in Friedman Robertson Walker Metric solves the Planck Era context.

1 The Planck 'constants'

Planck length $\Delta x = \sqrt{\frac{Gh}{c^3}}$

Planck time $\Delta t = \sqrt{\frac{Gh}{c^5}}$

Planck mass $\Delta m = \sqrt{\frac{hc}{G}}$

Planck acceleration $\Delta a = \frac{c}{\Delta t} = \sqrt{\frac{c^7}{hG}}$

2 Modern Cosmology

Within modern cosmology the Einstein's Field Equations would be written with cosmological term Λ as follows (see [2] and [3]):

$$R_{ik} - \frac{1}{2}g_{ik}R - \Lambda g_{ik} = \frac{8\pi G}{c^4}T_{ik} \quad (2.0)$$

The solutions of the Field Equations in Friedman-Roberson-Walker-Metric are:

FRW Equation (I)

$$\frac{\dot{R}^2}{R^2} = \frac{8\pi G\rho}{3} + \frac{\Lambda c^2}{3} - \frac{kc^2}{R^2} \quad (2.1)$$

FRW Equation (II)

$$\frac{\ddot{R}}{R} = \frac{\Lambda c^2}{3} - \frac{4\pi G}{3} \left(\rho + \frac{3p}{c^2}\right) \quad (2.2)$$

Einstein abandoned the cosmological term Λ as his "greatest blunder" after Hubble's 1928 discovery that the distant galaxies are expanding away from each other. Within a Universe with ideal Quantum gas and without cosmological term Λ and geometry factor k = 0 the equation (2.1) and (2.2) will become:

FRW Equation (I)

$$\frac{\dot{R}^2}{R^2} = \frac{8\pi G\rho}{3}$$
 (2.3)

and FRW Equation (II)

$$\frac{\ddot{R}}{R} = -\frac{4\pi G}{3} \left(\rho + \frac{3p}{c^2}\right) \quad (2.4)$$

With the relation $p = \frac{\rho c^2}{3}$ (Quantum gas) will change (2.4) as follows: FRW Equation (II)

$$\frac{R}{R} = -\frac{8\pi G\rho}{3} \quad (2.5)$$

Within Thermodynamics we assume dE = TdS - pdV and an adiabatic process it holds TdS = 0. We become $d(\epsilon V) = d\epsilon V + \epsilon dV = -pdV$ and it follows: With $d\epsilon = -(\epsilon + p)\frac{dV}{V}$ and the relation $p = \frac{\epsilon}{3}$ we assume:

$$\frac{d\epsilon}{\epsilon} = -\frac{4dV}{3V} \text{ or } \epsilon \sim V^{-4/3} \sim R^{-4} \quad (2.6)$$

3 The Cosmic Background Radiation

Now we know the energy-density in the Planck-Era $\rho c^2 = \tilde{a}\Delta T^4 = \frac{3c^7}{8\pi hG^2}$ ($\tilde{a} = Radiation \ constant = 7.5657e^{-16}$), If we assume $c = h = G = k_B = 1$ we get:

$$\frac{3}{8\pi} = \tilde{a}\Delta T^4 = \frac{8\pi^5}{15}T^4 \text{ or } T^4 = \frac{45}{64\pi^6}$$

Now we get the Planck-Temperature $\Delta T = (\frac{45}{64\pi^6})^{1/4} = \frac{1}{6.08088337383} = 5.8404e^{31} K$

In Planck-Era the following 6 relations are valid:

 $\Delta m \ \Delta x = \frac{h}{c} \quad (3.1)$ $\Delta m \ \Delta t = \frac{h}{c^2} \quad (3.2)$ $\frac{\Delta m}{\Delta a} = \frac{h}{c^3} \quad (3.3)$ $\frac{\Delta m}{\Delta x} = \frac{c^2}{G} \quad (3.4)$ $\frac{\Delta m}{\Delta t} = \frac{c^3}{G} \quad (3.5)$

With Planck-Force $\Delta F = \Delta m \ \Delta a = \frac{c^4}{G}$ (3.6)

4 Gravitation as curvature of space

In macroscopic scale the equation (3.1) til (3.6) will we rewritten as: (Entropy constant $\zeta = \sqrt{\frac{t}{\Delta t}} = 1.7952e^{30}$)

 $M R = \zeta^4 \frac{h}{c} \quad (4.1)$ $M t = \zeta^4 \frac{h}{c^2} \quad (4.2)$ $\frac{M}{a} = \zeta^4 \frac{h}{c^3} \quad (4.3)$ $\frac{M}{R} = \frac{c^2}{G} \quad (4.4)$ $\frac{M}{t} = \frac{c^3}{G} \quad (4.5)$

With Planck-Force $\Delta F = M \ a = \frac{c^4}{G}$ (4.6)

With Planck-Acceleration $a = \frac{G M}{R^2} = \frac{M c^3}{\zeta^4 h}$ follows:

$$\frac{G}{R^2} = \frac{c^3}{\zeta^4 h} \quad (4.7)$$

Furthermore we receive from GR:

$$\frac{M}{t} = \frac{c^3}{G} = \frac{\Delta m}{\Delta t} \Longrightarrow Age \ of \ Universe \ t = 4.355 e^{17} s$$
$$R = \zeta^2 \ \Delta x \Longrightarrow Radius \ of \ Universe \ R = 1.306 e^{26} m$$
$$\frac{M}{R} = \frac{c^2}{G} = \frac{\Delta m}{\Delta x} \Longrightarrow Mass \ of \ Universe \ M = 1.758 e^{53} kg$$

For $\dot{R}^2 = \frac{GM}{R}$ is with (4.7): $\dot{R}^2 = MR\frac{c^3}{\zeta^4 h} = c^2$

The FRW Equation (I) (2.3) is with $\dot{R} = c$ as follows:

$$\frac{c^2}{R^2} = \frac{8\pi G\rho}{3}$$

or with (4.7):

$$\frac{1}{R^4} = \frac{8\pi\rho c^2}{3\zeta^4 hc}$$

We become the R^4 dependency of (2.6) as follows:

$$\frac{3\zeta^4 hc}{8\pi R^4} = \rho c^2 = \tilde{a}T_\gamma^4$$

We get also:

$$\frac{1}{R} = \frac{Mc}{\zeta^4 h} \; q.e.d$$

5 References

- A.Einstein, Sitz. Preuss. Akad. d. Wiss., Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie (1917)
- A.Friedman, Über die Krümmung des Raumes, Zeitschrift für Physik 10 (1): 377 386. (1922)
- 3. V.Sahni, The Case for a Positive Cosmological A-Term, astro-ph/9904398
- 4. S.M.Carroll, The Cosmological Constant, astro-ph/0004075
- 5. A.I.Arbab, Nonstandard Cosmology With Constant and Variable Gravitational and Variable Cosmological Constants and Bulk Viscosity, http://arxiv.org/pdf/gr-qc/0105027.pdf