

# The idea of the Arithmetica

Hajime Mashima

June 22, 2014

## Abstract

During the 360 years of Fermat's last theorem is to be proved, this proposition was the presence appear full-length novel in "The Lord of the Rings", such as the "One Ring". And finally in 1994, it was proved completely by Andrew Wiles. However interesting proof is Fermat has been is still unknown. This will be assumed in the category of algebra probably.

## introduction

Natural number X,Y and Z solution of 3 or more that this equation holds  $X^n + Y^n = Z^n$  does not exist. Fermat is proven for the conditions of  $n = 4$ . It is sufficient if  $n$  is examining the conditions of prime numbers greater than or equal to 3 for this.

**Theorem 1** *Triangle the hypotenuse of Pythagorean theorem is  $z$ , can be expressed by the following relation by using the  $l$  and  $m$ .*

$$(l^2 - m^2)^2 + 2^2 (lm)^2 = (l^2 + m^2)^2$$

$$x^2 = (l^2 - m^2)^2$$

$$y^2 = 2^2 (lm)^2$$

$$z^2 = (l^2 + m^2)^2$$

$$(xyz \neq 0)$$

To simplify the algebra as a real number  $M$ , and  $N$ .

$$M, N \in \mathbb{R} \quad l^2 = M, \quad m^2 = N$$

$$(M - N)^2 + 2^2 MN = (M + N)^2$$

Put  $X, Y, Z \in \mathbb{N}$  *prime number*  $= p \geq 3$

$$\begin{aligned} X^p &= (M - N)^2 \\ Y^p &= 2^2 MN \\ Z^p &= (M + N)^2 \end{aligned}$$

$(XYZ \neq 0)$

Add the following conditions.  $X, Y, Z \in \text{even number}$

$$\begin{aligned} X^p &= 2^p X_1^p \\ Y^p &= 2^p Y_1^p \\ Z^p &= 2^p Z_1^p \\ (X_1, Y_1, Z_1 &\in \mathbb{N}) \end{aligned}$$

$$MN = 2^{p-2} Y_1^p \in \mathbb{N}$$

Thus  $M, N$  is a rational or irrational both.

## 1 $M, N$ is a condition of both rational

$$X^p = 2^p X_1^p, \quad Z^p = 2^p Z_1^p$$

$M - N, M + N \in \text{even number}$ , and it will be a divisor of  $2^{\frac{p+1}{2}}$  at least.

Consequently,  $Y_1^p \in \text{even number}$  so  $X_1^p, Z_1^p \in \text{even number}$ . (1)

## 2 $M, N$ is a condition of both irrational

$$\begin{aligned} MN &= 2^{p-2} Y_1^p \\ &= 2^{p-2} (Z_1^p - X_1^p) \\ &= 2^{p-2} \left( \sqrt{Z_1^p} + \sqrt{X_1^p} \right) \left( \sqrt{Z_1^p} - \sqrt{X_1^p} \right) \end{aligned}$$

$$X^p = 2^p X_1^p, \quad Z^p = 2^p Z_1^p$$

$$M = \left( \sqrt{2^{p-2} Z_1^p} + \sqrt{2^{p-2} X_1^p} \right) \quad N = \left( \sqrt{2^{p-2} Z_1^p} - \sqrt{2^{p-2} X_1^p} \right) \quad (M > N)$$

$$\begin{aligned} \text{Put } (c, d \in \text{odd number}) \quad h, i \in \mathbb{N} \\ M = 2^{\frac{h}{2}} c^{\frac{1}{2}} + 2^{\frac{i}{2}} d^{\frac{1}{2}} \quad N = 2^{\frac{h}{2}} c^{\frac{1}{2}} - 2^{\frac{i}{2}} d^{\frac{1}{2}} \end{aligned} \quad (\text{I})$$

In addition, assuming that there is no difference and sum,

$$\begin{aligned} &\text{Put}(U, V \in \text{odd number}) \\ M = 2^{\frac{l}{2}}U &\quad N = 2^{\frac{m}{2}}V \end{aligned} \quad (\text{II})$$

$$MN = 2^{p-2}Y_1^p = 2^{\frac{l+m}{2}}UV \in \mathbb{N}$$

M, N because irrational both, therefore  $(l, m \in \text{odd number})$ .

## 2.1 Conditions of (II)

### 2.1.1 Conditions of $(Y_1^p \in \text{odd number})$

$$Y_1^p = Z_1^p - X_1^p$$

$Z_1^p, X_1^p$  is the relationship of "odd and even" or "even and odd".

$Z_1^p$  and  $X_1^p$  are assumed to be coprime. Common divisor  $R^p (\in \text{odd number})$ , if present in the  $Z_1^p$  and  $X_1^p$ , is included as a common divisor of  $R^p$  also  $Y_1^p$ .  
 $(\frac{Y_1^p}{R^p} \in \mathbb{N})$

It is possible to remove common divisor, it is sufficient  $Z_1^p$  and  $X_1^p$  is examining the conditions of coprime.

$$MN = 2^{p-2}Y_1^p = 2^{\frac{l+m}{2}}UV \quad (p = \frac{l+m}{2} + 2 \quad Y_1^p = UV)$$

**Proposition 2**  $l > m \quad \frac{l+m}{2} > m \ (l, m \in \text{odd number} \quad U, V \in \text{odd number})$

$$\underline{\text{odd number} = 2^{\frac{l-m}{2}+2}X_1^p}$$

$$\begin{aligned} X^p &= (M - N)^2 \\ &= M^2 + N^2 - 2MN \\ &= 2^l U^2 + 2^m V^2 - 2 \cdot 2^{\frac{l+m}{2}} UV \\ &= 2^m \left( 2^{l-m} U^2 + V^2 - 2 \cdot 2^{\frac{l-m}{2}} UV \right) \\ &= 2^m (\text{odd number}) \end{aligned}$$

$$X^p = 2^m \left( 2^{\frac{l-m}{2}+2} X_1^p \right)$$

$$\text{odd number} \neq 2^{\frac{l-m}{2}+2} X_1^p \quad (2)$$

**Lemma 3**  $l = p - 2 \quad m = p - 2 \ (l, m \in \text{odd number} \quad U, V \in \text{odd number})$

Other things being does not hold all applies the infinite descent.

$$X^p = (M - N)^2 = \left(2^{\frac{p-2}{2}}U - 2^{\frac{p-2}{2}}V\right)^2 = 2^{p-2}(U - V)^2$$

$$\begin{aligned} 2^2 X_1^p &= (U - V)^2 & (U > V) \\ 2\sqrt{X_1^p} &= U - V & \cdots \textcircled{1} \end{aligned}$$

$$Z^p = (M + N)^2 = \left(2^{\frac{p-2}{2}}U + 2^{\frac{p-2}{2}}V\right)^2 = 2^{p-2}(U + V)^2$$

$$\begin{aligned} 2^2 Z_1^p &= (U + V)^2 & (U > V) \\ 2\sqrt{Z_1^p} &= U + V & \cdots \textcircled{2} \end{aligned}$$

$X_1^p, Z_1^p$  is a square number  $U \pm V$  because it is a natural number.

$$X_1^p = (X_H^p)^2 \quad Z_1^p = (Z_H^p)^2 \quad (X_H^p, Z_H^p \in \mathbb{N})$$

simultaneous equation:  $\textcircled{1} \pm \textcircled{2}$

$$U = Z_H^p + X_H^p \quad V = Z_H^p - X_H^p$$

If  $U, V$  is not a coprime, and a common divisor  $r$  ( $\in \text{odd number}$ ).

$$U = Z_H^p + X_H^p = rf \quad \cdots \textcircled{3}$$

$$V = Z_H^p - X_H^p = rg \quad \cdots \textcircled{4}$$

$$(U, V \in \text{odd number} \quad f, g \in \text{odd number})$$

simultaneous equation:  $\textcircled{3} \pm \textcircled{4}$

$$2Z_H^p = r(f + g)$$

$$2X_H^p = r(f - g)$$

$X_H^p, Z_H^p$  comprises a common divisor  $r$ . but  $X_H^p, Z_H^p$  must also be coprime  $X_1^p, Z_1^p$  is coprime. Thus  $U, V$  is coprime.

**Theorem 4** ( $Y_1^p = UV$ )  $U, V$  is at a coprime, which is a power of a prime number.

$$U = U_H^p, \quad V = V_H^p \quad Y_1^p = (U_H V_H)^p$$

Substitute  $U_H^p, V_H^p$  for  $\textcircled{3}, \textcircled{4}$ .

$$U_H^p = Z_H^p + X_H^p \quad V_H^p + X_H^p = Z_H^p \quad (3)$$

### 2.1.2 Conditions of ( $Y_1^p \in \text{even number}$ )

It is equivalent to  $X_1^p$  because it is odd number.

$$MN = 2^{p-2} X_1^p = 2^{\frac{l+m}{2}} UV \quad (p = \frac{l+m}{2} + 2 \quad X_1^p = UV)$$

#### Proposition 5

$$x^p + y^p = z^p$$

$x$  and  $z$  is a square number, both when this condition is satisfied.

$$\begin{aligned} st &= \left( \frac{s+t}{2} \right)^2 - \left( \frac{s-t}{2} \right)^2 \\ \left( \frac{s-t}{2} \right)^2 + st &= \left( \frac{s+t}{2} \right)^2 \\ \left( \frac{s-t}{s+t} \right)^2 + st \left( \frac{2}{s+t} \right)^2 &= 1 \end{aligned}$$

#### Corollary 6

$$\begin{aligned} \left( \frac{(st)^{\frac{1}{2}} (s-t)}{s+t} \right)^2 + \left( \frac{2st}{s+t} \right)^2 &= st \\ e^p + f^p &= (c^p)^2 \end{aligned}$$

Put

$$\begin{aligned} e^p &= \left( \frac{(st)^{\frac{1}{2}} (s-t)}{s+t} \right)^2 \\ f^p &= \left( \frac{2st}{s+t} \right)^2 \\ c^{2p} &= st \quad e, f, c \in \mathbb{N} \quad s, t \in \mathbb{R} \end{aligned}$$

“ $st$ ” is a natural number, when “ $s$ ” and “ $t$ ” is an irrational number, assumed as follows.

$$\begin{aligned} s &= \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right) K \\ t &= \left( G^{\frac{1}{2}} - H^{\frac{1}{2}} \right) K^{-1} \quad G, H \in \mathbb{N}, \quad K \in \mathbb{R} \end{aligned}$$

$$\begin{aligned} e^{\frac{p}{2}} &= \frac{c^p (s-t)}{s+t} & f^{\frac{p}{2}} &= \frac{2c^{2p}}{s+t} \\ \frac{e^{\frac{p}{2}}}{s-t} &= \frac{c^p}{s+t} & \dots \textcircled{5} & \quad \frac{f^{\frac{p}{2}}}{2c^p} = \frac{c^p}{s+t} & \dots \textcircled{6} \end{aligned}$$

By referring to the ⑤, ⑥.

$$\begin{aligned}
\frac{e^{\frac{p}{2}}}{s-t} &= \frac{f^{\frac{p}{2}}}{2c^p} \\
2c^p e^{\frac{p}{2}} &= f^{\frac{p}{2}} (s-t) \\
2c^p e^{\frac{p}{2}} &= f^{\frac{p}{2}} \left( \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right) K - \left( G^{\frac{1}{2}} - H^{\frac{1}{2}} \right) K^{-1} \right) \\
2c^p e^{\frac{p}{2}} K &= f^{\frac{p}{2}} \left( \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right) K^2 - \left( G^{\frac{1}{2}} - H^{\frac{1}{2}} \right) \right) \\
0 &= f^{\frac{p}{2}} \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right) K^2 - 2c^p e^{\frac{p}{2}} K - f^{\frac{p}{2}} \left( G^{\frac{1}{2}} - H^{\frac{1}{2}} \right) \\
\\
K &= \frac{2c^p e^{\frac{p}{2}} \pm \sqrt{2^2 c^{2p} e^p + 2^2 f^p (G-H)}}{2f^{\frac{p}{2}} \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right)} \\
&= \frac{c^p e^{\frac{p}{2}} \pm \sqrt{c^{2p} e^p + c^{2p} f^p}}{f^{\frac{p}{2}} \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right)} \\
&= \frac{c^p e^{\frac{p}{2}} \pm \sqrt{c^{2p} (e^p + f^p)}}{f^{\frac{p}{2}} \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right)} \\
&= \frac{c^p e^{\frac{p}{2}} \pm c^{2p}}{f^{\frac{p}{2}} \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right)} \\
s &= K \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right) \\
&= \frac{c^p e^{\frac{p}{2}} \pm c^{2p}}{f^{\frac{p}{2}}} \\
&= \frac{c^p (e^{\frac{p}{2}} \pm c^p)}{f^{\frac{p}{2}}} \\
t &= K^{-1} \left( G^{\frac{1}{2}} - H^{\frac{1}{2}} \right) \\
&= \frac{f^{\frac{p}{2}} \left( G^{\frac{1}{2}} + H^{\frac{1}{2}} \right)}{c^p e^{\frac{p}{2}} \pm c^{2p}} \left( G^{\frac{1}{2}} - H^{\frac{1}{2}} \right) \\
&= \frac{c^{2p} f^{\frac{p}{2}}}{c^p e^{\frac{p}{2}} \pm c^{2p}} \\
&= \frac{c^p f^{\frac{p}{2}}}{e^{\frac{p}{2}} \pm c^p}
\end{aligned}$$

$$\begin{aligned}
s + t &= \frac{c^p (e^{\frac{p}{2}} \pm c^p)}{f^{\frac{p}{2}}} + \frac{c^p f^{\frac{p}{2}}}{e^{\frac{p}{2}} \pm c^p} \\
&= \frac{c^p (e^{\frac{p}{2}} \pm c^p)^2 + c^p f^p}{f^{\frac{p}{2}} (e^{\frac{p}{2}} \pm c^p)} \\
s - t &= \frac{c^p (e^{\frac{p}{2}} \pm c^p)^2 - c^p f^p}{f^{\frac{p}{2}} (e^{\frac{p}{2}} \pm c^p)} \quad \dots \textcircled{7}
\end{aligned}$$

By substituting  $\textcircled{7}$ ,

$$\begin{aligned}
\left(\frac{2st}{s+t}\right)^2 &= st \left(1 - \left(\frac{s-t}{s+t}\right)^2\right) \\
f^p &= c^{2p} \left(1 - \left(\frac{s-t}{s+t}\right)^2\right)
\end{aligned}$$

$$\begin{aligned}
\left(\frac{s-t}{s+t}\right)^2 &= \left(\frac{c^p (e^{\frac{p}{2}} \pm c^p)^2 - c^p f^p}{f^{\frac{p}{2}} (e^{\frac{p}{2}} \pm c^p)} \cdot \frac{f^{\frac{p}{2}} (e^{\frac{p}{2}} \pm c^p)}{c^p (e^{\frac{p}{2}} \pm c^p)^2 + c^p f^p}\right)^2 \\
&= \left(\frac{c^p (e^{\frac{p}{2}} \pm c^p)^2 - c^p f^p}{c^p (e^{\frac{p}{2}} \pm c^p)^2 + c^p f^p}\right)^2 \\
&= \left(\frac{(e^{\frac{p}{2}} \pm c^p)^2 - f^p}{(e^{\frac{p}{2}} \pm c^p)^2 + f^p}\right)^2 \\
&= \left(\frac{e^p + c^{2p} \pm 2c^p e^{\frac{p}{2}} - f^p}{e^p + c^{2p} \pm 2c^p e^{\frac{p}{2}} + f^p}\right)^2 \\
&= \left(\frac{2e^p \pm 2c^p e^{\frac{p}{2}}}{2c^{2p} \pm 2c^p e^{\frac{p}{2}}}\right)^2 \\
&= \left(\frac{e^p \pm c^p e^{\frac{p}{2}}}{c^{2p} \pm c^p e^{\frac{p}{2}}}\right)^2 \\
\left(\frac{s+t}{s-t}\right)^2 &= \left(\frac{c^{2p} \pm c^p e^{\frac{p}{2}}}{e^p \pm c^p e^{\frac{p}{2}}}\right)^2 \\
&= \left(\frac{f^p + e^p \pm c^p e^{\frac{p}{2}}}{e^p \pm c^p e^{\frac{p}{2}}}\right)^2 \\
&= \left(\frac{f^p}{e^p \pm c^p e^{\frac{p}{2}}} + 1\right)^2
\end{aligned}$$

$\left(\frac{s-t}{s+t}\right)^2$  Because it is a rational number, “e” is the number of square or “s”, “t” is a rational number.

**Corollary 7** *When the following equation is satisfied,  $x^p$  and  $z^p$  is a square number*

$$x^p + y^p = z^p \quad (x, y \text{ and } z \text{ are disjoint.}) \quad (x^p, y^p, z^p \in \mathbb{N})$$

*And multiplied by  $z^p$  to both sides.*

$$z^p x^p + z^p y^p = z^{2p}$$

*Since " $z^p x^p$ " is also a square number,*

$$x^p = (x_1^p)^2, \quad z^p = (z_1^p)^2$$

$$(x_1^2)^p + y^p = (z_1^2)^p \quad (x_1, y \text{ and } z_1 \text{ are disjoint.})$$

By referring to the (3),

$$U_{II}^p = Z_{II}^p + X_{II}^p \quad V_{II}^p + X_{II}^p = Z_{II}^p$$

$$(2U_{II})^p = (2Z_{II})^p + (2X_{II})^p \quad (2V_{II})^p + (2X_{II})^p = (2Z_{II})^p$$

$$(2Z_{II})^p < Z^p = (2Z_1)^p \quad (2X_{II})^p < X^p = (2X_1)^p$$

Thus the lemma has been shown.

$$x^n + y^n \neq z^n \quad (xyz \neq 0 \quad n \geq 3)$$