

The Simplest Possible Cosmological Model with Explanation to Dark Energy

Mueiz Gafer KamalEdeen
mueizphysics@gmail.com
September 2014

Abstract

In this paper , a new and very simple cosmological model is introduced . It is shown how the model can explain some observations including Hubble's law and dark energy and affect our understanding of some concepts .


Introduction

The gradual accumulation of information about the large-scale universe produced an increasing desire for a simple and consistent description of the universe as a whole that led to the current standard model of cosmology and many competing models .

The following is a proposed alternative simpler model involving abandoning the global validity of some notions about space-time gained from our local experience.

The Model

According to this model the space-time is a 4-sphere in which the 3-dimensional surface represents the 3-space of the universe and the radius represents the time .



As a result of this relation between the *time* and the *3-space* we have to stop thinking about a single direction of time when we are exploring distances comparable to the radius (age) of the universe .*The idea that the time dimension in any point in the 3-space is the line from the center to that point represents the main feature of this model which distinguishes it from the existing ones.*


Now let us see how this model can be used to understand some facts and explain some observational data :

Hubble's Law and red shift : The world line of light as it travel through the 4-dimensional space-time between the source of light and the observer is a logarithmic spiral (tends to straight line in large values of the age of the universe) that keep making an angle ($\Pi / 4$) with the 3-dimensional surface in every time because the speed is equal to the tangent of this angle.

Thus the relation between the time of emission (T_e) and the time of observation (T_o) is :

$$T_o = T_e (e^{\theta})$$

Where (θ) is the angle between the world lines of the observer and the source .



The red-shift (z) resulted from this relation between the time of emission and the time of observation is :

$$z = (e^{\theta}) - 1$$

Now let us show how Hubble's law can be derived from this relation and the reason why it is valid only in small distances.

Firstly: from the properties of the exponential function for small values of (θ) the relation between (z) and (θ) is :

$$z \approx \theta \text{ for small value of } (\theta)$$

We have $\theta = D / T$ where (D) is the proper distance between the observer and the source.

Secondly : We have $z \approx v$ for small (v) where (v) is the velocity of the source moving away from the observer .

Thirdly: It can be deduced directly from this model that any increase in time is associated with an increase in the distance between any two objects proportional to this distance and the inverse of the age of the universe so the velocity of the increase of the proper distance is obtained from the relation : $v = D / T$.

Taking these three points into account we arrive at the following explanation of Hubble's law and its validity :

When applying the global law of red-shift $\{ z = (e^{\theta}) - 1 \}$ locally , the result is the same as that obtained when the red-shift is wrongly attributed to the velocity of the increase of the proper distance between the source and the observer .

The results can be drawn from this is that: $(T = 1/ H)$ and $(v= D/T)$ but this velocity cannot be used to calculate the red-shift although it gives nearly correct results in small distances .

It should also be noted that the relation $\{ z = (e^{\theta}) - 1 \}$ can be used to explain red-shift data claimed to be results of accelerated expansion (which is impossible according to this model) such as supernova observations .

The Large-scale Universe is Homogeneous and Isotropic : This can be thought of as a natural result of the shape of the *3-space* in this model , because there is no preferred region or direction in a spherical surface.

Dark Energy and the Cosmological Constant : As seen in the model and regardless of the material content at any moment in any place there exist a non-zero value for the curvature of spherical surface that decreases with the age of the universe this can be thought of as the so-called cosmological constant (which is no longer constant according to this model) which is used to explain dark energy . In addition to the Hubble's Law (as understood in this model) this curvature represents another way to determine the age of the universe.

The examination of the equality of the age of the universe obtained from Hubble's constant and the age obtained from this curvature (or the equivalent density of what is considered as dark energy) serves as a good test for this model :

Let (Λ) be the cosmological constant which is interpreted as dark energy:

The curvature of the 3-space = $12/ R^2$


The curvature (cosmological constant) $\sim 10^{-52} \text{ m}^{-2}$

Thus $R = (12 / \Lambda)^{(1/2)} \sim 3.464 * 10^{26} \text{ m}$

We have $(c / H) = 1.306 * 10^{26} \text{ m}$

They are nearly the same , the difference ranges within the uncertainty in the value of (Λ) .


Einstein's Field Equations : According to this model the geometry of large-scale universe is simply spherical geometry with the time dimension in the direction of the radius , this global geometry has nothing to do with the material content of the universe (**this can easily be obtained if we think of the cosmological constant as composed of two parts: the first is the general curvature of the spherical space and the second is the average density of the universe subtracted from the right-hand side of the equation**) while the local topological structure of the space-time is affected by the stress-energy tensor and the global geometry which is represented in the cosmological constant as obtained from *Einstein's Field Equation*.



Horizon Problem : The problem with the standard cosmological model that different regions of the universe have not contacted each other (according to the standard model) but have the same physical properties is known as the horizon problem . The cosmic microwave background radiation (CMB) which fills the space between galaxies is precisely the same everywhere.

This model provides a simple explanation to this homogeneity. According to the equation of the world line of light obtained from this model $\{ \mathbf{T}o =$

$T_e (e^\theta)$ } all the radiation emitted from a source with a world line of an angle ($\theta = n\pi$ where n is any integer) with the world line of the observer, at the time $\{ T_e = T_o (e^\theta - n\pi) = 0.043214 T_o \text{ for } n = 1 \}$ reaches the observer at time (T_o) from all direction. This supports a simple assumption that CMB. is the radiation coming from the same source which is converged again at the local space-time of the observer.



Conclusion

The space-time continuum can be divided into four dimensions the way we wish , but if we want to choose one of them to be measured by a clock , there must be some restrictions obtained from this model by which dark energy can be easily explained.