

# The Positron Fraction in Primary Cosmic Rays and New Cosmological Problems

Sylwester Kornowski

**Abstract:** Here, within the Scale-Symmetric Theory (SST), we calculated the positron fraction in primary cosmic rays as a function of energy and we described positron flux. Obtained results are consistent with the data from the Alpha Magnetic Spectrometer (AMS) on the International Space Station (ISS). These data lead to the internal structure of the core of baryons and to phenomena characteristic for regions in the Universe filled with baryonic plasma described within SST. Here as well we described new cosmological problems which lead to new cosmology. They are as follows. There are not in existence the B-modes associated with gravitational waves - it leads to conclusion that there was a separation in time of the inflation (the big bang) and the expansion of the Universe (the “soft” big bang) or/and that gravitational waves do not exist (SST shows that both conclusions are correct). In the Universe there is too small number of dwarf galaxies - there should be about 10 times more dwarf galaxies than we observe. In the very distant Universe there are numerous massive elliptical galaxies which do not significantly evolve so a time for their formation was too short. In the very distant Universe there are too many barred galaxies - it is inconsistent with simulations grounded on the Cosmological Standard Model (CSM). There is a substantial asymmetry in the CMB signal observed in the two opposite hemispheres of the sky. These new problems suggest that CSM starts from wrong initial conditions.

## 1. Introduction

The General Relativity leads to the non-gravitating Higgs field composed of tachyons [1A]. On the other hand, the Scale-Symmetric Theory (SST) shows that the succeeding phase transitions of such Higgs field lead to the different scales of sizes [1A]. Due to the saturation of interactions via the Higgs field and due to the law of conservation of the half-integral spin that is obligatory for all scales, there consequently appear the superluminal binary systems of closed strings (entanglons) responsible for the quantum entanglement (it is the quantum-entanglement scale), stable neutrinos and luminal neutrino-antineutrino pairs which are the components of the luminal Einstein spacetime (it is the Planck scale), cores of baryons (it is the electric-charges scale), and the cosmic structures (protoworlds; it is the cosmological scale) that evolution leads to the dark matter, dark energy and expanding universes (the “soft” big bangs) [1A], [1B]. The non-gravitating tachyons have infinitesimal spin so all listed structures have internal helicity (helicities) which distinguishes particles from their antiparticles [1A]. SST shows that a fundamental theory should start from infinite nothingness

and pieces of space [1A]. Sizes of pieces of space depend on their velocities [1A]. The inflation field started as the liquid-like field composed of non-gravitating pieces of space [1A]. Cosmoses composed of universes are created because of collisions of big pieces of space [1A], [1B]. During the inflation, the liquid-like inflation field (the non-gravitating superluminal Higgs field) transformed partially into the luminal Einstein spacetime (the big bang) [1A], [1B]. In our Cosmos, the two-component spacetime is surrounded by timeless wall – it causes that the fundamental constants are invariant [1A], [1B].

Due to the symmetrical decays of bosons on the equator of the core of baryons, there appears the atom-like structure of baryons described by the Titius-Bode orbits for the nuclear strong interactions [1A].

Applying 7 parameters only and a few new symmetries we calculated a thousand of basic physical (and mathematical) quantities (there are derived the physical and mathematical constants as well) consistent or very close to experimental data and observational facts ([http://vixra.org/author/sylwester\\_kornowski](http://vixra.org/author/sylwester_kornowski) ). In SST there do not appear approximations, mathematical tricks, and free parameters which are characteristic for the mainstream particle physics and mainstream cosmology.

Within the Standard Model (SM), since 1964, we cannot calculate the exact mass and spin of proton. It suggests that the 3-valence-quarks model cannot be realized by Nature at low energy. Just at low energies the perturbation theory breaks down so we do not understand the dynamics that lead to confinement and entanglement. Non-perturbative methods are required and the non-perturbative Scale-Symmetric Theory (SST) provides such methods [1]. Within SST, among many theoretical results, there are the rigorous mass and spin of proton, [1A], and the exact electron and muon radii of proton [2].

The SST shows that in centre of the baryons there is the core composed of torus/charge and condensate. The Einstein spacetime is grainy and consists of the neutrino-antineutrino pairs. Mass of each pair is very small (about  $6.7 \cdot 10^{-67}$  kg) and the total weak charge is equal to zero [1A] – it causes that detection of the Einstein-spacetime components is much difficult than neutrinos. The internal structure of the Einstein-spacetime components causes that they can be confined or/and entangled [1A]. The torus is built of entangled Einstein-spacetime components whereas the condensate consists of confined ones only. Mass of the charged core is  $H^+ = 727.44$  MeV whereas of the neutral one is  $H^0 = 724.78$  MeV [1A].

**In baryonic matter (protons and neutrons) the core cannot be charged negatively [1A]. It causes that due to the internal helicity of fermions, there appears the positron-electron asymmetry. The positron-electron pairs produced inside the positively charged core of baryons appear as pairs composed of real positron,  $e^+$ , and virtual electron,  $(e^-)^*$ .**

The calculated fine-structure constant is  $\alpha_{EM} = 1/137.036$  [1A]. The calculated coupling constant for weak interactions of proton and electron is  $\alpha'_{W(e-p)} = 1.11943581 \cdot 10^{-5}$  [1A]. The calculated weak binding energy of the torus/charge with the central condensate is  $E_W = 14.980$  MeV [1A]. The calculated electromagnetic binding energy of the torus and an electron placed on the circular axis of the torus is  $E_{EM} = 3.097$  MeV [1A] (this energy leads to the Higgs boson with a mass of 125 GeV [3]). The calculated mass of the condensate is  $Y = 424.124$  MeV [1A]. The calculated mass of muon is  $m_{muon} = 105.656314$  MeV [1A]. The calculated lower limit for the mass of a loop produced by the core of baryons is  $m_{LL} = 67.54441$  MeV whereas the upper limit is  $m_{UL} = 727.44$  MeV [1A]. The calculated mass of electron is  $m_{electron} = 0.5109989$  MeV, of mass of charged pion is  $m_{pion(+,-)} =$

139.57041 MeV [1A]. Calculated mass of the boson characteristic for baryonic plasma is  $m_{boson} \approx 283$  MeV [1D].

Table 1 *Characteristic features of the core of baryons*

Physical Quantity	Symbol	Theoretical value [1A], [1D]
Mass of positively charged core of baryons	$H^+$	727.44 MeV
Mass of neutral core of baryons	$H^0$	724.78 MeV
Fine-structure constant	$\alpha_{EM}$	1/137.036
Coupling constant for weak interactions of proton and electron	$\alpha'_{W(e-p)}$	$1.11943581 \cdot 10^{-5}$
Weak binding energy	$E_W$	14.980 MeV
Electromagnetic binding energy	$E_{EM}$	3.097 MeV
Mass of condensate	$Y$	424.124 MeV
Mass of muon	$m_{muon}$	105.656314 MeV
Lower limit for mass of a loop	$m_{LL}$	67.54441 MeV
Upper limit for mass of a loop	$m_{UL}$	727.44 MeV
Mass of electron/positron	$m_{electron}$	0.5109989 MeV
Mass of the boson characteristic for baryonic plasma	$m_{boson}$	$\sim 283$ MeV
Mass of charged pion	$m_{pion(+,-)}$	139.57041 MeV

The calculated masses of proton and neutron are respectively  $m_{proton} = 938.27$  MeV and  $m_{neutron} = 939.56$  MeV [1A].

Table 2 *Masses of nucleons*

Physical Quantity	Symbol	Theoretical value [1A]
Mass of proton	$m_{proton}$	938.27 MeV
Mass of neutron	$m_{neutron}$	939.56 MeV

Gluons and photons are the rotational energies of the Einstein-spacetime components. Inside the strong fields (they have internal helicity) the rotating Einstein-spacetime components behave as gluons whereas outside the strong fields (internal helicity is equal to zero) behave as photons. On the surfaces of the strong fields there is the gluon-photon transition [1A].

**Assume that due to the collisions of nucleons inside baryonic plasma (it consists of the cores of baryons [1A]), inside the core of baryons appear weak condensates with energy/mass  $E$ . Due to the gluon-photon transition, they can leak outside the plasma but then their energy/mass must increase  $F \gg 1$  times (i.e. there is the cooling of the baryonic plasma)**

$$F = \alpha_{EM} / \alpha'_{W(e-p)} = 651.88. \quad (1)$$

**In the SST there appears the four-particle symmetry [1A], [1B], [1D]. The four-particle symmetry follows from the fact that four particles can have total spin and total internal helicity equal to zero i.e. such objects do not create turbulences in spacetime at low and high energy.**

Here we will show that the positron fraction in primary cosmic rays as a function of energy and as a function of positron flux follow from the internal structure of the core of baryons and from the phenomena characteristic for the baryonic plasma.

We as well describe the origin of new cosmological problems which lead to new cosmology. They are as follows. There are not in existence the  $B$ -modes associated with gravitational waves – it leads to conclusion that there was a separation in time of the inflation (the big bang) and the expansion of the Universe (the “soft” big bang) or/and that gravitational waves do not exist (SST shows that both conclusions are correct). In the Universe there is too small number of dwarf galaxies - there should be about 10 times more dwarf galaxies than we observe. In the very distant Universe there are numerous massive elliptical galaxies which do not significantly evolve so a time for their formation was too short. In the very distant Universe there are too many barred galaxies – it is inconsistent with simulations grounded on the Cosmological Standard Model (CSM). There is a substantial asymmetry in the CMB signal observed in the two opposite hemispheres of the sky. These new problems suggest that CSM starts from wrong initial conditions.

## 2. The positron fraction and flux

The four-particle symmetry is very important inside baryonic plasma. Notice also that  $E_W$  can decay to  $4E_{EM}$ . Calculate energy of positron produced in following process

$$Nucleon + nucleon \rightarrow F ( E_W + 4E_{EM} ) \rightarrow 2 e^+ + 2 (e^-)*. \quad (2)$$

Energy of the positron is  $E_{LL,positron} = 8.9 \text{ GeV}$  (precisely, for  $4E_{EM}$  is 8.08 GeV whereas for  $E_W$  is 9.77 GeV). It is the lower limit for energy of positron produced by the core of baryons in baryonic plasma for the interval with increasing positron fraction.

Notice that  $Y$  can decay to  $4m_{muon}$ . Calculate energy of positron in following process

$$Nucleon + nucleon \rightarrow F ( Y + 4m_{muon} ) \rightarrow 2 e^+ + 2 (e^-)*. \quad (3)$$

Energy of the positron is  $E_{UL,positron} = 276.0 \text{ GeV}$  (precisely, for  $Y$  is 276.5 GeV whereas for  $4m_{muon}$  is 275.5 GeV). It is the upper limit for energy of positron produced by the core of baryons in baryonic plasma for the interval with increasing positron fraction.

Calculate the positron fraction in cosmic rays for the lower and upper limits.

The lowest mass of a loop is  $m_{LL}$  and such loops appear on the circular axis of the torus whereas the highest mass of a loop is  $m_{UL}$  and such loops appear on the equator of the torus i.e. in the  $d = 0$  state [1A]. Since the four-particle symmetry is very important in baryonic plasma so the number of positrons and electrons for the lower limit for the interval with increasing positron fraction ( $positron \text{ fraction} = R = e^+ / (e^+ + e^-)$ ) is proportional to  $4m_{LL}$  whereas for the upper limit is proportional to  $4m_{UL}$ . On the other hand, the number of positrons for the lower limit is proportional to the mean mass of condensates carrying mass equal to  $E_W$  or  $4E_{EM}$  i.e.  $E_{LL,mean} = ( E_W + 4E_{EM} ) / 2 = 13.684 \text{ MeV}$  whereas for the

upper limit is proportional to the mean mass of condensates carrying mass equal to  $Y$  or  $4m_{muon}$  i.e.  $E_{UL,mean} = (Y + 4m_{muon}) / 2 = 423.375$  MeV.

The positron fraction  $R$  for the lower limit is

$$R_{LL} = \{ e^+ / (e^+ + e^-) \}_{LL} = E_{LL,mean} / 4m_{LL} = 0.05065 \approx 0.05. \quad (4)$$


The positron fraction for the upper limit is

$$R_{UL} = \{ e^+ / (e^+ + e^-) \}_{UL} = E_{UL,mean} / 4m_{UL} = 0.14550 \approx 0.15. \quad (5)$$

Since there is the logarithmic scale so the positron fraction, for the interval with increasing positron fraction, as a function of energy of positrons  $E$  looks as follows

$$\log \{ e^+ / (e^+ + e^-) \} = a \log ( E / E_0 ) + b, \quad (6)$$

where  $E_0 = 8.9$  GeV,  $a = 0.31$  whereas  $b = -1.30$ . For energy  $E = 8.9$  GeV is  $R = 0.050$ , for  $E = 30$  GeV is  $R = 0.073$ , for  $E = 70$  GeV is  $R = 0.095$  and for  $E = 276$  GeV is  $R = 0.145$ .



The characteristic features of the positron fraction in cosmic rays are as follows (Fig.1).

### 2.1.

The mean energy at which the positron fraction begins to increase is 8.9 GeV (8.08 and 9.77 GeV) and is about 0.05.

## 2.2.

There exists the sharp structure (i.e. the core of baryons) with a mass of 727.44 MeV which leads to energy of positron about 474 GeV.

## 2.3.

The mean energy at which the positron fraction reaches its maximum is 276 GeV (275.5 and 276.5 GeV) and is about 0.15.

## 2.4.

The rate of increase with energy defines following function


$$\log R = a \log ( E / E_0 ) + b. \quad (7)$$

## 2.5.

The increase in positron fraction follows from the phenomena characteristic for baryonic plasma. Such plasma consists of the cores of baryons which structure is described within the Scale-Symmetric Theory.

The obtained theoretical results for the positron fraction are consistent with the data from the Alpha Magnetic Spectrometer (AMS) on the International Space Station (ISS) [4].

Calculate the characteristic features for the positron flux (Fig.2).



Due to the thermal motions of the cores of baryons in baryonic plasma, the positron flux should increase for following interval  $(F m_{positron}, F E_{LL,mean}) \equiv (0.33 \text{ GeV}, 8.9 \text{ GeV})$ . Next, for interval  $(F E_{LL,mean}, F m_{muon}) \equiv (8.9 \text{ GeV}, 68.9 \text{ GeV})$ , each core of nucleon can produce one positron so the positron flux should level out. Above 69 GeV, once again, due to the thermal motions, increases number of positively charged muons so number of positrons increases as well. Such increasing positron flux we should observe up to the upper limit  $FE_{UL,positron} = 276 \text{ GeV}$ . For energies higher than 276 GeV there are mostly produced the muon-antimuon pairs so positron flux should rapidly decrease.

We should observe a small increase in positron flux for  $Fm_{pion(+)} \approx 91$  GeV and for  $Fm_{boson} \approx 184$  GeV.

For  $Fm_{muon} = 68.9$  GeV, the positron flux should be  $H^+ / m_{LL} = 10.77$  times lower than electron flux.

Since the mass distance between neutron and proton is about two times smaller than the mass distance between the charged and neutral cores of baryons so for the lower limit 8.9 GeV the electron flux should be about two times higher than the positron flux. It follows from the fact that for energy 8.08 GeV there is the transition from the positively charged core of baryons to the neutral one ( $H^+ - H^0 \approx 2.66$  MeV) i.e. the transition from proton to neutron – in the beta decay of this neutron there appears electron (the total maximum energy of electron is  $m_{neutron} - m_{proton} \approx 1.3$  MeV), proton and electron-antineutrino. We can see that  $2.66 / 1.3 \approx 2$ .

The obtained theoretical results for the flux are consistent with the data from the Alpha Magnetic Spectrometer (AMS) on the International Space Station (ISS) [5].

### 3. New cosmological problems

There appeared new cosmological problems which lead to new cosmology described within SST. They are as follows.

#### 3.1.

There are not in existence the  $B$ -modes associated with gravitational waves – it leads to conclusion that there was a separation in time of the inflation and the “soft” big bang of the Universe or/and that luminal gravitational waves are not in existence.

Both conclusions are consistent with new cosmology described within SST [1B], [6].

#### 3.2.

In the very distant Universe there is too small number of dwarf galaxies (about 10 times too small).

According to the Cosmological Standard Model (CSM), the bigger cosmic structures have been forming due to the mergers of smaller structures. It leads to conclusion that galaxies should have much more satellite/dwarf galaxies. On the other hand, the SST shows that the very early Universe was the double cosmic loop composed of binary systems of protogalaxies which consisted of the neutron black holes [1B]. The protogalaxies evolved due to the inflows of the dark matter and dark energy into the double cosmic loop [1B]. Why we cannot see the first stage of evolution of the double cosmic loop? Why we already see the massive galaxies and massive black holes at the very distant Universe? The answer follows from the duality of relativity [7]. SST shows that the most distant galaxies are in the time distance equal to  $13.866 \pm 0.096$  Gyr but they are already 7.75 Gyr old [1B], [7], [8]. Just the real age of the Universe is  $21.614 \pm 0.096$  Gyr [1B].

#### 3.3.

In the very distant Universe there are the elliptical massive galaxies which do not significantly evolve [9] so a time for their formation was too short. In the very distant Universe there are too many barred galaxies – it is inconsistent with simulations grounded on the Cosmological Standard Model (CSM) ([9], at  $z \sim 1.5$ , there among the disc galaxies is about 14% barred galaxies).

It is consistent with SST. The shape of the double cosmic loop, [1B], leads to numerous barred galaxies.

### 3.4.

There is a substantial asymmetry in the CMB signal observed in the two opposite hemispheres of the sky [10]. One of the two hemispheres appears to have a significantly stronger signal on average. It is inconsistent with CSM.

Such result is consistent with SST. SST shows that the Universe is moving with a sound speed about 355 m/s in relation to the Einstein spacetime. It causes that the frontal hemisphere is a little hotter [12].

Described above new cosmological problems suggest that CSM starts from wrong initial conditions. We can solve the new problems within the Scale-Symmetric Theory.

### 4. Summary

Here, within the Scale-Symmetric Theory (SST), we calculated the positron fraction in primary cosmic rays as a function of energy and we described positron flux. Obtained results are consistent with the data from the Alpha Magnetic Spectrometer (AMS) on the International Space Station (ISS). These data lead to the internal structure of the core of baryons and to phenomena characteristic for regions in the Universe filled with baryonic plasma described within SST.

Here as well we described new cosmological problems which lead to new cosmology. They are as follows. There are not in existence the B-modes associated with gravitational waves – it leads to conclusion that there was a separation in time of the inflation (the big bang) and the expansion of the Universe (the “soft” big bang) or/and that gravitational waves do not exist (SST shows that both conclusions are correct). In the Universe there is too small number of dwarf galaxies - there should be about 10 times more dwarf galaxies than we observe. In the very distant Universe there are numerous massive elliptical galaxies which do not significantly evolve so a time for their formation was too short. In the very distant Universe there are too many barred galaxies – it is inconsistent with simulations grounded on the Cosmological Standard Model (CSM). There is a substantial asymmetry in the CMB signal observed in the two opposite hemispheres of the sky. These new problems suggest that CSM starts from wrong initial conditions.

### References

- [1] Sylwester Kornowski (2015). *Scale-Symmetric Theory*  
 [1A]: <http://vixra.org/abs/1511.0188> (Particle Physics)  
 [1B]: <http://vixra.org/abs/1511.0223> (Cosmology)  
 [1C]: <http://vixra.org/abs/1511.0284> (Chaos Theory)  
 [1D]: <http://vixra.org/abs/1512.0020> (Reformulated QCD)
- [2] Sylwester Kornowski (2015). “The Root-Mean-Square Charge Radius of Proton”  
<http://vixra.org/abs/1301.0174>
- [3] Sylwester Kornowski (2015). “The Higgs Mechanism and Tower of Higgs Bosons”  
<http://vixra.org/abs/1310.0094>
- [4] L. Accardo *et al.* (AMS Collaboration), (18 September 2014). “High Statistics Measurement of the Positron Fraction in Primary Cosmic Rays of 0.5–500 GeV with the Alpha Magnetic Spectrometer on the International Space Station”  
 Phys. Rev. Lett. **113**, 121101
- [5] M. Aguilar *et al.* (AMS Collaboration), (18 September 2014). “Electron and Positron Fluxes in Primary Cosmic Rays Measured with the Alpha Magnetic Spectrometer on the International Space Station”  
 Phys. Rev. Lett. **113**, 121102

- [6] Sylwester Kornowski (2016). "The E- and B-Modes in the CMB"  
<http://vixra.org/abs/1403.0501>
- [7] Sylwester Kornowski (2016). "The Duality of Relativity"  
<http://vixra.org/abs/1405.0327>
- [8] Sylwester Kornowski (2016). "The Evolution of Massive Galaxies"  
<http://vixra.org/abs/1406.0014>
- [9] B. D. Simmons *et al.*, "Galaxy Zoo: CANDELS Barred Disks and Bar Fractions"  
*Monthly Notices of the Royal Astronomical Society*, 2014  
DOI:10.1093/mnras/stu1817
- [10] [http://www.aanda.org/index.php?option=com\\_toc&url=/articles/aa/abs/2014/11/contents/contents.html](http://www.aanda.org/index.php?option=com_toc&url=/articles/aa/abs/2014/11/contents/contents.html)
- [11] Sylwester Kornowski (2015). "The Origin and Fate of the Cosmos and Universes and the Classical-Quantum Asymmetry"  
<http://vixra.org/abs/1308.0138>