

$$= \bigcup_{\alpha \in \lambda} \mu_{\alpha}(x) \Rightarrow \bigcup_{\alpha \in \lambda} \mu_{\alpha}(0) \geq \bigcup_{\alpha \in \lambda} \mu_{\alpha}(x) \quad \forall x \in X .$$

ii. Let $x, z \in Q, y \in X$. $\bigcup_{\alpha \in \lambda} \mu_{\alpha}(x * z^n) = \sup\{ \mu_{\alpha}(x), \alpha \in \lambda \} \geq$

$$\sup\{ \min\{ \mu_{\alpha}((x * y) * z^n), \mu_{\alpha}(y) \}, \alpha \in \lambda \}$$

[Since μ_{α} is a Q-Smarandache fuzzy n-fold strong ideal, $\forall \alpha \in \lambda$. By definition 3(ii)]

But $\{ \alpha, \alpha \in \lambda \}$ is a chain \Rightarrow there exist, $j \in \lambda$ such that $\sup\{ \min\{ \mu_{\alpha}((x * y) * z^n), \mu_{\alpha}(y) \}, \alpha \in \lambda \} = \min\{ \mu_j((x * y) * z^n), \mu_j(y) \}$

$$= \min\{ \sup\{ \mu_{\alpha}((x * y) * z^n), \alpha \in \lambda \}, \sup\{ \mu_{\alpha}(y), \alpha \in \lambda \} \}$$

$$\Rightarrow \bigcup_{\alpha \in \lambda} \mu_{\alpha}(x * z^n) \geq \min\{ \mu_j((x * y) * z^n), \mu_j(y) \}$$

$$\geq \min\{ \sup\{ \mu_{\alpha}((x * y) * z^n), \alpha \in \lambda \}, \sup\{ \mu_{\alpha}(y), \alpha \in \lambda \} \} = \min\{$$

$$\bigcup_{\alpha \in \lambda} \mu_{\alpha}((x * y) * z^n), \bigcup_{\alpha \in \lambda} \mu_{\alpha}(y) \}$$

$$\Rightarrow \bigcup_{\alpha \in \lambda} \mu_{\alpha}(x * z^n) \geq \min\{ \bigcup_{\alpha \in \lambda} \mu_{\alpha}((x * y) * z^n), \bigcup_{\alpha \in \lambda} \mu_{\alpha}(y) \} \Rightarrow$$

$$\bigcup_{\alpha \in \lambda} \mu_{\alpha} \text{ is a Q-Smarandache fuzzy n-fold strong ideal of } X.$$

■

References

- [1] A.B.Saeid and A. Namdar, "Smarandache BCH-algebras", World Applied Sciences Journal 7 (11), 1818-4952, pp.77-83, 2009.
- [2] C. H. Park, "Interval-valued Fuzzy Ideal in BH-algebras", Advance in fuzzy set and systems 1(3), pp.231-240, 2006.
- [3] E. M.kim and S.S. Ahn, " On fuzzy n-fold strong ideals of BH-algebras" , J. Appl. Math. & Informatics Vol. 30, No.3 - 4, pp. 665 - 676, 2012 .
- [4] E. M. kim and S. S. Ahn, " Fuzzy strong ideals of BH-algebras with degrees in the interval (0; 1]", J. Appl. Math. & Informatics Vol. 31, No. 1- 2, pp. 211 - 220, 2013.
- [5] H. H. Abbass , "The Q-Smarandache closed ideal and Q-Smarandache Fuzzy Closed Ideal with Respect to an Element of a Q-Smarandache BCH-Algebra", Journal kerbala university, vol. 10, No.3 scientific, pp. 302-312, 2012 .
- [6] H. H. Abbass and S. J. Mohammed "On a Q-Smarandache Fuzzy Completely Closed Ideal with Respect to an Element of a BH-algebra", Kufa University, M.Sc. thesis, 2013.
- [7] L. A. Zadeh, "Fuzzy sets", Information and control ,Vol. 8, PP. 338-353, 1965.
- [8] O. G. Xi, "Fuzzy BCK-algebra", Math. Japonica 36, No. 5, pp.935-942, 1991.
- [9] Q. Zhang, Y. B. Jun and E. H. Roh, "On the branch of BH-algebras", Scientiae Mathematicae Japonicae Online, Vol.4 , pp.917- 921, 2001.
- [10] Q. Zhang, E. H. Roh and Y. B. Jun, "On Fuzzy BH-algebras", J.Huanggang, Normal Univ. 21(3), pp.14-19, 2001.

- [11] Y. B. Jun, "Closed Fuzzy ideals in BCI-algebras", Math. Japonica 38, No. 1, pp.199-202, 1993.
- [12] Y. B. Jun, E. H. Roh and H. S. Kim, "On BH-algebras", Scientiae Mathematicae , 1(1) , pp.347-354, 1998.
- [13] Y.B.Jun, "Fuzzy Closed Ideals and Fuzzy filters in BCH-algebras", J. Fuzzy Math. 7 , No. 2, pp.435-444, 1999.