
On Strong Interval Valued Neutrosophic Graphs

Said Broumi¹, Mohamed Talea², Assia Bakali³, Florentin
Smarandache⁴

^{1,2}Laboratory of Information processing, Faculty of Science Ben M'Sik, University Hassan II,
B.P 7955, Sidi Othman, Casablanca, Morocco. broumisaid78@gmail.com,
taleamohamed@yahoo.fr

³Ecole Royale Navale, Boulevard Sour Jdid, B.P 16303 Casablanca, Morocco.
assiabakali@yahoo.fr

⁴Department of Mathematics, University of New Mexico, 705 Gurley Avenue, Gallup, NM
87301, USA

fsmarandache@gmail.com

Abstract: In this paper, we discuss a subclass of interval valued neutrosophic graphs called strong interval valued neutrosophic graphs which were introduced by Broumi et al [41]. The operations of Cartesian product, composition, union and join of two strong interval valued neutrosophic graphs are defined. Some propositions involving strong interval valued neutrosophic graphs are stated and proved.

Keywords: Single valued neutrosophic graph, interval valued neutrosophic graph, strong interval valued neutrosophic graph, cartesian product, composition, union, and join.

1. Introduction

Neutrosophic set proposed by Smarandache [13, 14] is a powerful tool to deal with incomplete, indeterminate and inconsistent information in real world. It is a generalization of the theory of fuzzy set [30], intuitionistic fuzzy sets [27, 29], interval-valued fuzzy sets [22] and interval-valued intuitionistic fuzzy sets [28], then the neutrosophic set is characterized by a truth-membership degree (t), an indeterminacy-membership degree (i) and a falsity-membership degree (f) independently, which are within the real standard or nonstandard unit interval $]-0, 1+[$. Therefore, if their range is restrained within the real standard unit interval $[0, 1]$, the neutrosophic set is easily applied to engineering problems. For this purpose, Wang et al. [17] introduced the concept of a single valued neutrosophic set (SVNS) as a subclass of the neutrosophic set. The same authors introduced the notion of interval valued neutrosophic sets [18] as subclass of neutrosophic sets in

which the value of truth-membership, indeterminacy-membership and falsity-membership degrees are intervals of numbers instead of the real numbers. Recently, the concept of single valued neutrosophic set and interval valued neutrosophic sets have been applied in a wide variety of fields including computer science, engineering, mathematics, medicine and economic [3, 4, 5, 6, 16, 19, 20, 21, 23, 24, 25, 26, 32, 34, 35, 36, 37, 38, 43].

Lots of works on fuzzy graphs and intuitionistic fuzzy graphs [7, 8, 9, 31, 33] have been carried out and all of them have considered the vertex sets and edge sets as fuzzy and /or intuitionistic fuzzy sets. But, when the relations between nodes(or vertices) in problems are indeterminate, the fuzzy graphs and intuitionistic fuzzy graphs are failed. For this purpose, Smarandache [10, 11] have defined four main categories of neutrosophic graphs, two based on literal indeterminacy (I), which called them; I-edge neutrosophic graph and I-vertex neutrosophic graph, these concepts are studied deeply and has gained popularity among the researchers due to its applications via real world problems [1, 12, 15, 44, 45, 46]. The two others graphs are based on (t, i, f) components and called them; The (t, i, f)-edge neutrosophic graph and the (t, i, f)-vertex neutrosophic graph, these concepts are not developed at all. Later on, Broumi et al. [40] introduced a third neutrosophic graph model combined the (t, i, f)-edge and and the (t, i, f)-vertex neutrosophic graph and investigated some of their properties. The third neutrosophic graph model is called single valued neutrosophic graph (SVNG for short). The single valued neutrosophic graph is the generalization of fuzzy graph and intuitionistic fuzzy graph. Also, the same authors [39] introduced neighborhood degree of a vertex and closed neighborhood degree of vertex in single valued neutrosophic graph as a generalization of neighborhood degree of a vertex and closed neighborhood degree of vertex in fuzzy graph and intuitionistic fuzzy graph. On the other hand, Broumi et al. [41] introduced the concept of interval valued neutrosophic graphs , which is a generalization of fuzzy graph, intuitionistic fuzzy graph, interval valued fuzzy graph, interval valued intuitionistic fuzzy graph and single valued neutrosophic graph,. Also, Broumi et al. [42] studied some operations on interval valued neutrosophic graphs. Motivated by the operations on (crisp) graphs such as Cartesian product, composition, union and join, In this paper, we define the operations of cartesian product, composition, union and join on strong interval valued neutrosophic graphs and investigate some their properties.

2.Preliminaries

In this section, we mainly recall some notions related to neutrosophic sets, single valued neutrosophic sets, interval valued neutrosophic sets, fuzzy graph and intuitionistic fuzzy graph, interval valued intuitionstic fuzzy graph, and interval valued neutrosophic graph. relevant to the present work. See especially [2, 7, 8, 13, 17, 40, 41] for further details and background.

Definition 2.1 [13]. Let X be a space of points (objects) with generic elements in X denoted by x ; then the neutrosophic set A (NS A) is an object having the form $A = \{ \langle x: T_A(x), I_A(x), F_A(x) \rangle, x \in X \}$, where the functions $T, I, F: X \rightarrow]-0,1+[$ define respectively the a truth-membership function, an indeterminacy-membership function, and a falsity-membership function of the element $x \in X$ to the set A with the condition:

$$-0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3^+ \quad (1)$$

The functions $T_A(x), I_A(x)$ and $F_A(x)$ are real standard or nonstandard subsets of $] -0,1+[$.

Since it is difficult to apply NSs to practical problems, Wang et al. [16] introduced the concept of a SVN, which is an instance of a NS and can be used in real scientific and engineering applications.

Definition 2.2 [17]. Let X be a space of points (objects) with generic elements in X denoted by x . A single valued neutrosophic set A (SVNS A) is characterized by truth-membership function $T_A(x)$, an indeterminacy-membership function $I_A(x)$, and a falsity-membership function $F_A(x)$. For each point x in X $T_A(x), I_A(x), F_A(x) \in [0, 1]$. A SVN A can be written as

$$A = \{ \langle x: T_A(x), I_A(x), F_A(x) \rangle, x \in X \} \quad (2)$$

Definition 2.3[7]. A fuzzy graph is a pair of functions $G = (\sigma, \mu)$ where σ is a fuzzy subset of a non empty set V and μ is a symmetric fuzzy relation on σ . i.e $\sigma: V \rightarrow [0,1]$ and

$\mu: V \times V \rightarrow [0,1]$ such that $\mu(uv) \leq \sigma(u) \wedge \sigma(v)$ for all $u, v \in V$ where uv denotes the edge between u and v and $\sigma(u) \wedge \sigma(v)$ denotes the minimum of $\sigma(u)$ and $\sigma(v)$. σ is called the fuzzy vertex set of V and μ is called the fuzzy edge set of E .




Figure 1: Fuzzy Graph

Definition 2.4 [7]. The fuzzy subgraph $H = (\tau, \rho)$ is called a fuzzy subgraph of $G = (\sigma, \mu)$ if $\tau(u) \leq \sigma(u)$ for all $u \in V$ and $\rho(u, v) \leq \mu(u, v)$ for all $u, v \in V$

Definition 2.5 [8]. An intuitionistic fuzzy graph is of the form $G=(V, E)$ where

- i. $V=\{v_1, v_2, \dots, v_n\}$ such that $\mu_1: V \rightarrow [0,1]$ and $\gamma_1: V \rightarrow [0,1]$ denote the degree of membership and nonmembership of the element $v_i \in V$, respectively, and $0 \leq \mu_1(v_i) + \gamma_1(v_i) \leq 1$ for every $v_i \in V, (i = 1, 2, \dots, n)$,
- ii. $E \subseteq V \times V$ where $\mu_2: V \times V \rightarrow [0,1]$ and $\gamma_2: V \times V \rightarrow [0,1]$ are such that $\mu_2(v_i, v_j) \leq \min [\mu_1(v_i), \mu_1(v_j)]$ and $\gamma_2(v_i, v_j) \geq \max [\gamma_1(v_i), \gamma_1(v_j)]$ and $0 \leq \mu_2(v_i, v_j) + \gamma_2(v_i, v_j) \leq 1$ for every $(v_i, v_j) \in E, (i, j = 1, 2, \dots, n)$




Figure 2: Intuitionistic Fuzzy Graph

Definition 2.6 [40]. Let $A = (T_A, I_A, F_A)$ and $B = (T_B, I_B, F_B)$ be single valued neutrosophic sets on a set X . If $A = (T_A, I_A, F_A)$ is a single valued neutrosophic relation on a set X , then $A = (T_A, I_A, F_A)$ is called a single valued neutrosophic relation on $B = (T_B, I_B, F_B)$ if

$$T_B(x, y) \leq \min(T_A(x), T_A(y))$$

$$I_B(x, y) \geq \max(I_A(x), I_A(y)) \text{ and}$$

$$F_B(x, y) \geq \max(F_A(x), F_A(y)) \text{ for all } x, y \in X.$$

A single valued neutrosophic relation A on X is called symmetric if $T_A(x, y) = T_A(y, x), I_A(x, y) = I_A(y, x), F_A(x, y) = F_A(y, x)$ and $T_B(x, y) = T_B(y, x), I_B(x, y) = I_B(y, x)$ and $F_B(x, y) = F_B(y, x)$, for all $x, y \in X$.

Definition 2.7 [2] An interval valued intuitionistic fuzzy graph with underlying set V is defined to be a pair $G= (A, B)$ where

1) The functions $M_A : V \rightarrow D [0, 1]$ and $N_A : V \rightarrow D [0, 1]$ denote the degree of membership and non membership of the element $x \in V$, respectively, such that $0 \leq M_A(x) + N_A(x) \leq 1$ for all $x \in V$.

2) The functions $M_B : E \subseteq V \times V \rightarrow D [0, 1]$ and $N_B : E \subseteq V \times V \rightarrow D [0, 1]$ are defined by $M_{BL}(x, y) \leq \min (M_{AL}(x), M_{AL}(y))$ and $N_{BL}(x, y) \geq \max (N_{AL}(x), N_{AL}(y))$

$M_{BU}(x, y) \leq \min (M_{AU}(x), M_{AU}(y))$ and $N_{BU}(x, y) \geq \max (N_{AU}(x), N_{AU}(y))$ such that

$0 \leq M_{BU}(x, y) + N_{BU}(x, y) \leq 1$ for all $(x, y) \in E$.

Definition 2.8. [41] By an interval-valued neutrosophic graph of a graph $G^* = (V, E)$ we mean a pair $G = (A, B)$, where $A = \langle [T_{AL}, T_{AU}], [I_{AL}, I_{AU}], [F_{AL}, F_{AU}] \rangle$ is an interval-valued neutrosophic set on V and $B = \langle [T_{BL}, T_{BU}], [I_{BL}, I_{BU}], [F_{BL}, F_{BU}] \rangle$ is an interval-valued neutrosophic relation on E satisfies the following condition:

1. $V = \{v_1, v_2, \dots, v_n\}$ such that $T_{AL}: V \rightarrow [0, 1]$, $T_{AU}: V \rightarrow [0, 1]$, $I_{AL}: V \rightarrow [0, 1]$, $I_{AU}: V \rightarrow [0, 1]$ and $F_{AL}: V \rightarrow [0, 1]$, $F_{AU}: V \rightarrow [0, 1]$ denote the degree of truth-membership, the degree of indeterminacy-membership and falsity-membership of the element $v \in V$, respectively, and

$$0 \leq T_A(v_i) + I_A(v_i) + F_A(v_i) \leq 3 \text{ for all } v_i \in V (i=1, 2, \dots, n)$$

2. The functions $T_{BL}: V \times V \rightarrow [0, 1]$, $T_{BU}: V \times V \rightarrow [0, 1]$, $I_{BL}: V \times V \rightarrow [0, 1]$, $I_{BU}: V \times V \rightarrow [0, 1]$ and $F_{BL}: V \times V \rightarrow [0, 1]$, $F_{BU}: V \times V \rightarrow [0, 1]$ are such that

$$T_{BL}(\{v_i, v_j\}) \leq \min [T_{AL}(v_i), T_{AL}(v_j)]$$

$$T_{BU}(\{v_i, v_j\}) \leq \min [T_{AU}(v_i), T_{AU}(v_j)]$$

$$I_{BL}(\{v_i, v_j\}) \geq \max [I_{BL}(v_i), I_{BL}(v_j)]$$

$$I_{BU}(\{v_i, v_j\}) \geq \max [I_{BU}(v_i), I_{BU}(v_j)] \text{ And}$$

$$F_{BL}(\{v_i, v_j\}) \geq \max [F_{BL}(v_i), F_{BL}(v_j)]$$

$$F_{BU}(\{v_i, v_j\}) \geq \max [F_{BU}(v_i), F_{BU}(v_j)]$$

Denotes the degree of truth-membership, indeterminacy-membership and falsity-membership of the edge $(v_i, v_j) \in E$ respectively, where

$$0 \leq T_B(\{v_i, v_j\}) + I_B(\{v_i, v_j\}) + F_B(\{v_i, v_j\}) \leq 3 \text{ for all } \{v_i, v_j\} \in E (i, j = 1, 2, \dots, n)$$

We call A the interval valued neutrosophic vertex set of V , B the interval valued neutrosophic edge set of E , respectively, Note that B is a symmetric interval valued neutrosophic relation on A . We use the notation (v_i, v_j) for an element of E Thus, $G = (A, B)$ is a interval valued neutrosophic graph of $G^* = (V, E)$ if

$$T_{BL}(v_i, v_j) \leq \min [T_{AL}(v_i), T_{AL}(v_j)]$$

$$T_{BU}(v_i, v_j) \leq \min [T_{AU}(v_i), T_{AU}(v_j)]$$

$$I_{BL}(v_i, v_j) \geq \max [I_{BL}(v_i), I_{BL}(v_j)]$$

$$I_{BU}(v_i, v_j) \geq \max [I_{BU}(v_i), I_{BU}(v_j)] \text{ And}$$

$$F_{BL}(v_i, v_j) \geq \max [F_{BL}(v_i), F_{BL}(v_j)]$$

$$F_{BU}(v_i, v_j) \geq \max [F_{BU}(v_i), F_{BU}(v_j)] \text{ for all } (v_i, v_j) \in E.$$

Here after, we use the notation xy for (x,y) an element of E .

3. Strong interval valued neutrosophic graph

Throught this paper, we denote $G^* = (V, E)$ a crisp graph, and $G=(A, B)$ an interval valued neutrosophic graph.

Definition 3.1 An interval valued neutrosophic graph $G = (A, B)$ is called strong interval valued neutrosophic graph if

$$T_{BL}(xy) = \min (T_{AL}(x) , T_{AL}(y)), I_{BL}(xy) = \max (I_{AL}(x) , I_{AL}(y)) \text{ and } F_{BL}(xy) = \max (F_{AL}(x) , F_{AL}(y))$$

$$T_{BU}(xy) = \min (T_{AU}(x) , T_{AU}(y)), I_{BU}(xy) = \max (I_{AU}(x) , I_{AU}(y)) \text{ and } F_{BU}(xy) = \max (F_{AU}(x) , F_{AU}(y)) \text{ such that}$$

$$0 \leq T_{BU}(x, y) + I_{BU}(x, y) + F_{BU}(x, y) \leq 3 \text{ for all } xy \in E.$$

Example 3.2 Figure 1 is an example for IVNG, $G=(A, B)$ defined on a graph $G^* = (V, E)$ such that $V = \{x, y, z\}$, $E = \{xy, yz, zx\}$, A is an interval valued neutrosophic set of V

$$A = \{ \langle x, [0.5, 0.7], [0.2, 0.3], [0.1, 0.3] \rangle, \langle y, [0.6, 0.7], [0.2, 0.4], [0.1, 0.3] \rangle, \langle z, [0.4, 0.6], [0.1, 0.3], [0.2, 0.4] \rangle \}$$

$$B = \{ \langle xy, [0.3, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle yz, [0.3, 0.5], [0.2, 0.3], [0.2, 0.4] \rangle, \langle xz, [0.3, 0.5], [0.1, 0.5], [0.2, 0.4] \rangle \}$$




Figure 3: Interval valued neutrosophic graph

Example 3.2 Figure 2 is a SIVNG $G = (A, B)$, where

$$A = \{ \langle x, [0.5, 0.7], [0.1, 0.4], [0.1, 0.3] \rangle, \langle y, [0.6, 0.7], [0.2, 0.3], [0.1, 0.3] \rangle, \langle z, [0.4, 0.6], [0.2, 0.3], [0.2, 0.4] \rangle \}$$

$$B = \{ \langle xy, [0.5, 0.7], [0.2, 0.4], [0.1, 0.3] \rangle, \langle yz, [0.4, 0.6], [0.2, 0.3], [0.2, 0.4] \rangle, \langle xz, [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$$




Figure 4:Strong Interval valued neutrosophic graph.

Proposition 3.3: A strong interval valued neutrosophic graph is the generalization of strong interval valued fuzzy graph

Proof: Suppose $G=(V, E)$ be a strong interval valued neutrosophic graph. Then by setting the indeterminacy- membership and falsity- membership values of vertex set and edge set equals to zero reduces the strong interval valued neutrosophic graph to strong interval valued fuzzy graph.

Proposition 3.4: A strong interval valued neutrosophic graph is the generalization of strong interval valued intuitionistic fuzzy graph

Proof: Suppose $G=(V, E)$ be a strong interval valued neutrosophic graph. Then by setting the indeterminacy- membership values of vertex set and edge set equals to zero reduces the strong interval valued neutrosophic graph to strong interval valued intuitionistic fuzzy graph.

Proposition 3.5: A strong interval valued neutrosophic graph is the generalization of strong intuitionistic fuzzy graph

Proof: Suppose $G=(V, E)$ be a strong interval valued neutrosophic graph. Then by setting the indeterminacy- membership, upper truth-membership and upper falsity-membership values of vertex set and edge set equals to zero reduces the strong interval valued neutrosophic graph to strong intuitionistic fuzzy graph.

Proposition 3.6: A strong interval valued neutrosophic graph is the generalization of strong single neutrosophic graph.

Proof: Suppose $G= (V, E)$ be a strong interval valued neutrosophic graph. Then by setting the upper truth-membership equals lower truth-membership, upper indeterminacy- membership equals lower indeterminacy-membership

and upper falsity-membership equals lower falsity- membership values of vertex set and edge set reduces the strong interval valued neutrosophic graph to strong single valued neutrosophic graph.

Definition 3.7 Let A_1 and A_2 be interval-valued neutrosophic subsets of V_1 and V_2 respectively. Let B_1 and B_2 interval-valued neutrosophic subsets of E_1 and E_2 respectively. The Cartesian product of two SIVNGs G_1 and G_2 is denoted by $G_1 \times G_2 = (A_1 \times A_2, B_1 \times B_2)$ and is defined as follows:

- 1) $(T_{A_1L} \times T_{A_2L})(x_1, x_2) = \min(T_{A_1L}(x_1), T_{A_2L}(x_2))$
 $(T_{A_1U} \times T_{A_2U})(x_1, x_2) = \min(T_{A_1U}(x_1), T_{A_2U}(x_2))$
 $(I_{A_1L} \times I_{A_2L})(x_1, x_2) = \max(I_{A_1L}(x_1), I_{A_2L}(x_2))$
 $(I_{A_1U} \times I_{A_2U})(x_1, x_2) = \max(I_{A_1U}(x_1), I_{A_2U}(x_2))$
 $(F_{A_1L} \times F_{A_2L})(x_1, x_2) = \max(F_{A_1L}(x_1), F_{A_2L}(x_2))$
 $(F_{A_1U} \times F_{A_2U})(x_1, x_2) = \max(F_{A_1U}(x_1), F_{A_2U}(x_2))$ for all $(x_1, x_2) \in V$
- 2) $(T_{B_1L} \times T_{B_2L})((x, x_2)(x, y_2)) = \min(T_{A_1L}(x), T_{B_2L}(x_2y_2))$
 $(T_{B_1U} \times T_{B_2U})((x, x_2)(x, y_2)) = \min(T_{A_1U}(x), T_{B_2U}(x_2y_2))$
 $(I_{B_1L} \times I_{B_2L})((x, x_2)(x, y_2)) = \max(I_{A_1L}(x), I_{B_2L}(x_2y_2))$
 $(I_{B_1U} \times I_{B_2U})((x, x_2)(x, y_2)) = \max(I_{A_1U}(x), I_{B_2U}(x_2y_2))$
 $(F_{B_1L} \times F_{B_2L})((x, x_2)(x, y_2)) = \max(F_{A_1L}(x), F_{B_2L}(x_2y_2))$
 $(F_{B_1U} \times F_{B_2U})((x, x_2)(x, y_2)) = \max(F_{A_1U}(x), F_{B_2U}(x_2y_2)) \forall x \in V_1$ and
 $\forall x_2y_2 \in E_2$
- 3) $(T_{B_1L} \times T_{B_2L})((x_1, z)(y_1, z)) = \min(T_{B_1L}(x_1y_1), T_{A_2L}(z))$
 $(T_{B_1U} \times T_{B_2U})((x_1, z)(y_1, z)) = \min(T_{B_1U}(x_1y_1), T_{A_2U}(z))$
 $(I_{B_1L} \times I_{B_2L})((x_1, z)(y_1, z)) = \max(I_{B_1L}(x_1y_1), I_{A_2L}(z))$
 $(I_{B_1U} \times I_{B_2U})((x_1, z)(y_1, z)) = \max(I_{B_1U}(x_1y_1), I_{A_2U}(z))$
 $(F_{B_1L} \times F_{B_2L})((x_1, z)(y_1, z)) = \max(F_{B_1L}(x_1y_1), F_{A_2L}(z))$
 $(F_{B_1U} \times F_{B_2U})((x_1, z)(y_1, z)) = \max(F_{B_1U}(x_1y_1), F_{A_2U}(z)) \forall z \in V_2$ and
 $\forall x_1y_1 \in E_1$

Proposition 3.7 If G_1 and G_2 are the strong interval valued neutrosophic graphs, then the cartesian product $G_1 \times G_2$ is a strong interval valued neutrosophic graph.

Proof:

Let G_1 and G_2 are SIVNGs, there exist $x_i, y_i \in E_i, i=1, 2$ such that

$$\begin{aligned} T_{B_iL}(x_i, y_i) &= \min(T_{A_iL}(x_i), T_{A_iL}(y_i)), i=1, 2. \\ T_{B_iU}(x_i, y_i) &= \min(T_{A_iU}(x_i), T_{A_iU}(y_i)), i=1, 2. \\ I_{B_iL}(x_i, y_i) &= \max(I_{A_iL}(x_i), I_{A_iL}(y_i)), i=1, 2. \\ I_{B_iU}(x_i, y_i) &= \max(I_{A_iU}(x_i), I_{A_iU}(y_i)), i=1, 2. \end{aligned}$$

$$F_{B_iL}(x_i, y_i) = \max (F_{A_iL}(x_i), F_{A_iL}(y_i)), i = 1, 2.$$

$$F_{B_iU}(x_i, y_i) = \max (F_{A_iU}(x_i), F_{A_iU}(y_i)), i = 1, 2.$$

Let $E = \{(x, x_2) (x, y_2) / x \in V_1, x_2y_2 \in E_2\} \cup \{(x_1, z) (y_1, z) / z \in V_2, x_1y_1 \in E_1\}$

Consider, $(x, x_2) (x, y_2) \in E$, we have

$$(T_{B_1L} \times T_{B_2L}) ((x, x_2) (x, y_2)) = \min (T_{A_1L}(x), T_{B_2L}(x_2y_2))$$

$$= \min (T_{A_1L}(x), T_{A_2L}(x_2), T_{A_2L}(y_2))$$

Similarly,

$$(T_{B_1U} \times T_{B_2U}) ((x, x_2) (x, y_2)) = \min (T_{A_1U}(x), T_{B_2U}(x_2y_2))$$

$$= \min (T_{A_1U}(x), T_{A_2U}(x_2), T_{A_2U}(y_2))$$

$$(T_{A_1L} \times T_{A_2L}) (x_1, x_2) = \min (T_{A_1L}(x_1), T_{A_2L}(x_2))$$

$$(T_{A_1U} \times T_{A_2U}) (x_1, x_2) = \min (T_{A_1U}(x_1), T_{A_2U}(x_2))$$

$$(T_{A_1L} \times T_{A_2L}) (x_1, y_2) = \min (T_{A_1L}(x_1), T_{A_2L}(y_2))$$

$$(T_{A_1U} \times T_{A_2U}) (x_1, y_2) = \min (T_{A_1U}(x_1), T_{A_2U}(y_2))$$

$$\text{Min} ((T_{A_1U} \times T_{A_2U}) (x, x_2), (T_{A_1U} \times T_{A_2U}) (x, y_2))$$

$$= \min (\min (T_{A_1U}(x), T_{A_2U}(x_2)), \min (T_{A_1U}(x), T_{A_2U}(y_2)))$$

$$= \min (T_{A_1U}(x), T_{A_2U}(x_2), T_{A_1U}(y_2))$$

Hence,

$$(T_{B_1L} \times T_{B_2L}) ((x, x_2) (x, y_2)) = \min ((T_{A_1L} \times T_{A_2L}) (x, x_2), (T_{A_1L} \times T_{A_2L}) (x, y_2))$$

$$(T_{B_1U} \times T_{B_2U}) ((x, x_2) (x, y_2)) = \min ((T_{A_1U} \times T_{A_2U}) (x, x_2), (T_{A_1U} \times T_{A_2U}) (x, y_2))$$

Similarly, we can show that

$$(I_{B_1L} \times I_{B_2L}) ((x, x_2) (x, y_2)) = \max ((I_{A_1L} \times I_{A_2L}) (x, x_2), (I_{A_1L} \times I_{A_2L}) (x, y_2))$$

$$(I_{B_1U} \times I_{B_2U}) ((x, x_2) (x, y_2)) = \max ((I_{A_1U} \times I_{A_2U}) (x, x_2), (I_{A_1U} \times I_{A_2U}) (x, y_2))$$

And also,

$$(F_{B_1L} \times F_{B_2L}) ((x, x_2) (x, y_2)) = \max ((F_{A_1L} \times F_{A_2L}) (x, x_2), (F_{A_1L} \times F_{A_2L}) (x, y_2))$$

$$(F_{B_1U} \times F_{B_2U}) ((x, x_2) (x, y_2)) = \max ((F_{A_1U} \times F_{A_2U}) (x, x_2), (F_{A_1U} \times F_{A_2U}) (x, y_2))$$

Hence, $G_1 \times G_2$ strong interval valued neutrosophic graph. This completes the proof.

Proposition 3.8 If $G_1 \times G_2$ is strong interval valued neutrosophic graph then at least G_1 or G_2 must be strong.

Proof:

Let G_1 and G_2 are no strong interval valued neutrosophic graphs, there exist $x_i, y_i \in E_i, i=1, 2$ such that

$$\begin{aligned} T_{B_iL}(x_i, y_i) &< \min (T_{A_iL}(x_i), T_{A_iL}(y_i)), i=1, 2. \\ T_{B_iU}(x_i, y_i) &< \min (T_{A_iU}(x_i), T_{A_iU}(y_i)), i=1, 2. \\ I_{B_iL}(x_i, y_i) &> \max (I_{A_iL}(x_i), I_{A_iL}(y_i)), i=1, 2. \\ I_{B_iU}(x_i, y_i) &> \max (I_{A_iU}(x_i), I_{A_iU}(y_i)), i=1, 2. \\ F_{B_iL}(x_i, y_i) &> \max (F_{A_iL}(x_i), F_{A_iL}(y_i)), i=1, 2. \\ F_{B_iU}(x_i, y_i) &> \max (F_{A_iU}(x_i), F_{A_iU}(y_i)), i=1, 2. \end{aligned}$$

Let $E = \{(x, x_2) (x, y_2) / x \in V_1, x_2 y_2 \in E_2\} \cup \{(x_1, z) (y_1, z) / z \in V_2, x_1 y_1 \in E_1\}$

Consider, $(x, x_2) (x, y_2) \in E$, we have

$$\begin{aligned} (T_{B_1L} \times T_{B_2L}) ((x, x_2) (x, y_2)) &= \min (T_{A_1L}(x), T_{B_2L}(x_2 y_2)) \\ &< \min (T_{A_1L}(x), T_{A_2L}(x_2), T_{A_2L}(y_2)) \end{aligned}$$

Similarly,

$$\begin{aligned} (T_{B_1U} \times T_{B_2U}) ((x, x_2) (x, y_2)) &= \min (T_{A_1U}(x), T_{B_2U}(x_2 y_2)) \\ &< \min (T_{A_1U}(x), T_{A_2U}(x_2), T_{A_2U}(y_2)) \end{aligned}$$

$$(T_{A_1L} \times T_{A_2L}) (x_1, x_2) = \min (T_{A_1L}(x_1), T_{A_2L}(x_2))$$

$$(T_{A_1U} \times T_{A_2U}) (x_1, x_2) = \min (T_{A_1U}(x_1), T_{A_2U}(x_2))$$

$$(T_{A_1L} \times T_{A_2L}) (x_1, y_2) = \min (T_{A_1L}(x_1), T_{A_2L}(y_2))$$

$$(T_{A_1U} \times T_{A_2U}) (x_1, y_2) = \min (T_{A_1U}(x_1), T_{A_2U}(y_2))$$

$$\begin{aligned} \text{Min} ((T_{A_1U} \times T_{A_2U}) (x, x_2), (T_{A_1U} \times T_{A_2U}) (x, y_2)) \\ &= \min (\min (T_{A_1U}(x), T_{A_2U}(x_2)), \min (T_{A_1U}(x), T_{A_2U}(y_2))) \\ &= \min (T_{A_1U}(x), T_{A_2U}(x_2), T_{A_1U}(y_2)) \end{aligned}$$

Hence,

$$(T_{B_1L} \times T_{B_2L}) ((x, x_2) (x, y_2)) < \min ((T_{A_1L} \times T_{A_2L}) (x, x_2), (T_{A_1L} \times T_{A_2L}) (x, y_2))$$

$$(T_{B_1U} \times T_{B_2U}) ((x, x_2) (x, y_2)) < \min ((T_{A_1U} \times T_{A_2U}) (x, x_2), (T_{A_1U} \times T_{A_2U}) (x, y_2))$$

Similarly, we can show that

$$(I_{B_1L} \times I_{B_2L}) ((x, x_2) (x, y_2)) > \max ((I_{A_1L} \times I_{A_2L}) (x, x_2), (I_{A_1L} \times I_{A_2L}) (x, y_2))$$

$$(I_{B_1U} \times I_{B_2U}) ((x, x_2) (x, y_2)) > \max ((I_{A_1U} \times I_{A_2U}) (x, x_2), (I_{A_1U} \times I_{A_2U}) (x, y_2))$$

And also,

$$(F_{B_1L} \times F_{B_2L}) ((x, x_2) (x, y_2)) > \max ((F_{A_1L} \times F_{A_2L}) (x, x_2), (F_{A_1L} \times F_{A_2L}) (x, y_2))$$

$$(F_{B_1U} \times F_{B_2U}) ((x, x_2) (x, y_2)) > \max ((F_{A_1U} \times F_{A_2U}) (x, x_2), (F_{A_1U} \times F_{A_2U}) (x, y_2))$$

Hence, $G_1 \times G_2$ is not strong interval valued neutrosophic graph, which is a contradiction . This completes the proof.

Remark: 3.9 If G_1 is a SIVNG and G_2 is not a SIVNG, then $G_1 \times G_2$ is need not be an SIVNG.

Example 3.10 Let $G_1 = (A_1, B_1)$ be a SIVNG, where $A_1 = \{< a, [0.6, 0.7], [0.2, 0.5], [0.1, 0.3]>, < b, [0.6, 0.7], [0.2, 0.5], [0.1, 0.3]>\}$ and $B_1 = \{< ab, [0.6, 0.7], [0.2, 0.5], [0.1, 0.3]>\}$




Figure 5: Interval valued neutrosophic G_1 .

$G_2 = (A_2, B_2)$ is not a SIVNG, where $A_2 = \{< c, [0.4, 0.6], [0.2, 0.4], [0.1, 0.3]>, < d, [0.4, 0.6], [0.1, 0.3], [0.2, 0.4]>\}$ and $B_2 = \{< cd, [0.3, 0.5], [0.1, 0.2], [0.3, 0.5]>\}$.




Figure 6: Interval valued neutrosophic G_2 .

$G_1 \times G_2 = (A_1 \times A_2, B_1 \times B_2)$ is not a SIVNG, where

$A_1 \times A_2 = \{< (a, c), [0.4, 0.6], [0.2, 0.3], [0.2, 0.4]>, < (a, d), [0.4, 0.6], [0.2, 0.3], [0.2, 0.4]>, < (b, c), [0.4, 0.6], [0.2, 0.6], [0.2, 0.4]>, < (b, d), [0.4, 0.6], [0.3, 0.4], [0.2, 0.4]>\}$,

$B_1 \times B_2 = \{< ((a, c), (a, d)), [0.3, 0.5], [0.3, 0.5], [0.3, 0.5]>, < ((a, c), (b, c)), [0.4, 0.6], [0.1, 0.4], [0.3, 0.4]>, < ((b, c), (b, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4]>, < ((a, d), (b, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4]>\}$. In this example, G_1 is a SIVNG and G_2 is not a SIVNG, then $G_1 \times G_2$ is not a SIVNG.




Figure 7: Cartesian product $G_1 \times G_2$

Example 3.11 Let $G_1 = (A_1, B_1)$ be a SIVNG, where $A_1 = \{ \langle a, [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle b, [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle \}$ and $B_1 = \{ \langle ab, [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle \}$,




Figure 8: Interval valued neutrosophic G_1 .

$G_2 = (A_2, B_2)$ is not a SIVIFG, where $A_2 = \{ \langle c, [0.6, 0.7], [0.1, 0.3], [0.1, 0.3] \rangle, \langle d, [0.6, 0.7], [0.1, 0.3], [0.2, 0.4] \rangle \}$ and $B_2 = \{ \langle cd, [0.5, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$,




Figure 9: Interval valued neutrosophic G_2 .

$G_1 \times G_2 = (A_1 \times A_2, B_1 \times B_2)$ is a SIVNG, where

$A_1 \times A_2 = \{ \langle (a, c), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle (a, d), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle (b, c), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle (b, d), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$ and

$B_1 \times B_2 = \{ \langle ((a, c), (a, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle ((a, c), (b, c)), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle ((b, c), (b, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle ((a, d), (b, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$. In this example, G_1 is a SIVNG and G_2 is not a SIVNG, then $G_1 \times G_2$ is a SIVNG.




Figure 10: Cartesian product

Proposition 3.12 Let G_1 be a strong interval valued neutrosophic graph.

Then for any interval valued neutrosophic graph $G_2, G_1 \times G_2$ is strong interval valued neutrosophic graph iff

$$T_{A_1L}(x_1) \leq T_{B_1L}(x_2y_2), I_{A_1L}(x_1) \geq I_{B_1L}(x_2y_2) \text{ and } F_{A_1L}(x_1) \geq F_{B_1L}(x_2y_2),$$

$$T_{A_1U}(x_1) \leq T_{B_1U}(x_2y_2), I_{A_1U}(x_1) \geq I_{B_1U}(x_2y_2) \text{ and } F_{A_1U}(x_1) \geq F_{B_1U}(x_2y_2),$$

$$\forall x_1 \in V_1, x_2y_2 \in E_2$$

Definition 3.13 Let A_1 and A_2 be interval valued neutrosophic subsets of V_1 and V_2 respectively. Let B_1 and B_2 interval-valued neutrosophic subsets of E_1 and E_2 respectively. The **composition** of two strong interval valued neutrosophic graphs G_1 and G_2 is denoted by $G_1 \circ G_2 = (A_1 \circ A_2, B_1 \circ B_2)$ and is defined as follows:

- 1) $(T_{A_1L} \circ T_{A_2L})(x_1, x_2) = \min(T_{A_1L}(x_1), T_{A_2L}(x_2))$
 $(T_{A_1U} \circ T_{A_2U})(x_1, x_2) = \min(T_{A_1U}(x_1), T_{A_2U}(x_2))$
 $(I_{A_1L} \circ I_{A_2L})(x_1, x_2) = \max(I_{A_1L}(x_1), I_{A_2L}(x_2))$
 $(I_{A_1U} \circ I_{A_2U})(x_1, x_2) = \max(I_{A_1U}(x_1), I_{A_2U}(x_2))$
 $(F_{A_1L} \circ F_{A_2L})(x_1, x_2) = \max(F_{A_1L}(x_1), F_{A_2L}(x_2))$
 $(F_{A_1U} \circ F_{A_2U})(x_1, x_2) = \max(F_{A_1U}(x_1), F_{A_2U}(x_2)) \forall x_1 \in V_1, x_2 \in V_2$
- 2) $(T_{B_1L} \circ T_{B_2L})((x, x_2)(x, y_2)) = \min(T_{A_1L}(x), T_{B_2L}(x_2y_2))$
 $(T_{B_1U} \circ T_{B_2U})((x, x_2)(x, y_2)) = \min(T_{A_1U}(x), T_{B_2U}(x_2y_2))$
 $(I_{B_1L} \circ I_{B_2L})((x, x_2)(x, y_2)) = \max(I_{A_1L}(x), I_{B_2L}(x_2y_2))$
 $(I_{B_1U} \circ I_{B_2U})((x, x_2)(x, y_2)) = \max(I_{A_1U}(x), I_{B_2U}(x_2y_2))$
 $(F_{B_1L} \circ F_{B_2L})((x, x_2)(x, y_2)) = \max(F_{A_1L}(x), F_{B_2L}(x_2y_2))$
 $(F_{B_1U} \circ F_{B_2U})((x, x_2)(x, y_2)) = \max(F_{A_1U}(x), F_{B_2U}(x_2y_2)) \forall x \in V_1, \forall x_2y_2 \in E_2$
- 3) $(T_{B_1L} \circ T_{B_2L})((x_1, z)(y_1, z)) = \min(T_{B_1L}(x_1y_1), T_{A_2L}(z))$

$$(T_{B_1U} \circ T_{B_2U}) ((x_1, z) (y_1, z)) = \min (T_{B_1U}(x_1y_1), T_{A_2U}(z))$$

$$(I_{B_1L} \circ I_{B_2L}) ((x_1, z) (y_1, z)) = \max (I_{B_1L}(x_1y_1), I_{A_2L}(z))$$

$$(I_{B_1U} \circ I_{B_2U}) ((x_1, z) (y_1, z)) = \max (I_{B_1U}(x_1y_1), I_{A_2U}(z))$$

$$(F_{B_1L} \circ F_{B_2L}) ((x_1, z) (y_1, z)) = \max (F_{B_1L}(x_1y_1), F_{A_2L}(z))$$

$$(F_{B_1U} \circ F_{B_2U}) ((x_1, z) (y_1, z)) = \max (F_{B_1U}(x_1y_1), F_{A_2U}(z)) \forall z \in V_2, \forall x_1y_1 \in E_1$$

$$4) (T_{B_1L} \circ T_{B_2L}) ((x_1, x_2) (y_1, y_2)) = \min (T_{A_2L}(x_2), T_{A_2L}(y_2), T_{B_1L}(x_1y_1))$$

$$(T_{B_1U} \circ T_{B_2U}) ((x_1, x_2) (y_1, y_2)) = \min (T_{A_2U}(x_2), T_{A_2U}(y_2), T_{B_1U}(x_1y_1))$$

$$(I_{B_1L} \circ I_{B_2L}) ((x_1, x_2) (y_1, y_2)) = \max(I_{A_2L}(x_2), I_{A_2L}(y_2), I_{B_1L}(x_1y_1))$$

$$(I_{B_1U} \circ I_{B_2U}) ((x_1, x_2) (y_1, y_2)) = \max (I_{A_2U}(x_2), I_{A_2U}(y_2), I_{B_1U}(x_1y_1))$$

$$(F_{B_1L} \circ F_{B_2L}) ((x_1, x_2) (y_1, y_2)) = \max(F_{A_2L}(x_2), F_{A_2L}(y_2), F_{B_1L}(x_1y_1))$$

$$(F_{B_1U} \circ F_{B_2U})((x_1, x_2) (y_1, y_2)) = \max (F_{A_2U}(x_2), F_{A_2U}(y_2), F_{B_1U}(x_1y_1))$$

$\forall (x_1, x_2)(y_1, y_2) \in E^0 - E$, where $E^0 = E \cup \{(x_1, x_2) (y_1, y_2) | x_1y_1 \in E_1, x_2 \neq y_2\}$

The following propositions are stated without their proof.

Proposition 3.14 If G_1 and G_2 are the strong interval valued neutrosophic graphs, then the composition $G_1[G_2]$ is a strong interval valued neutrosophic graph.

Proposition 3.15 If $G_1[G_2]$ is strong interval valued neutrosophic graphs, then at least composition G_1 or G_2 must be strong.

Example 3.16 Let $G_1 = (A_1, B_1)$ be a SIVNG, where $A_1 = \{< a, [0.6, 0.7], [0.2, 0.3], [0.1, 0.3], >, < b, [0.6, 0.7], [0.2, 0.3], [0.1, 0.3] >\}$ and $B_1 = \{< ab, [0.6, 0.7], [0.2, 0.3], [0.1, 0.3] >\}$.




Figure 11: Interval valued neutrosophic G_1 .

$G_2 = (A_2, B_2)$ is not a SIVNG, where $A_2 = \{< c, [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] >, < d, [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] >\}$ and $B_2 = \{< cd, [0.3, 0.5], [0.2, 0.5], [0.3, 0.5] >\}$.




Figure 12: Interval valued neutrosophic G_2 .

$G_1[G_2] = (A_1 \circ A_2, B_1 \circ B_2)$ is not a SIVNG, where

$A_1 \circ A_2 = \{ \langle (a,c), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle (a,d), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle (b,c), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle (b,d), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle \}$,

$B_1 \circ B_2 = \{ \langle ((a,c), (a,d)), [0.3, 0.5], [0.2, 0.4], [0.3, 0.5] \rangle, \langle ((a,c), (b,c)), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle ((b,c), (b,d)), [0.3, 0.5], [0.2, 0.4], [0.3, 0.5] \rangle, \langle ((a,d), (b,d)), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle ((a,c), (b,d)), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle, \langle ((a,d), (b,c)), [0.4, 0.6], [0.2, 0.4], [0.1, 0.3] \rangle \}$. In this example, G_1 is a SIVNG and G_2 is not a SIVNG, then $G_1[G_2]$ is not a SIVNG.




Figure 13: Composition

Example 3.17 Let $G_1 = (A_1, B_1)$ be a SIVNG, where $A_1 = \{ \langle a, [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle b, [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$ and $B_1 = \{ \langle ab, [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$.




Figure 14: Interval valued neutrosophic G_1 .

$G_2 = (A_2, B_2)$ is not a SIVNG, where $A_2 = \{ \langle c, [0.6, 0.7], [0.1, 0.3], [0.1, 0.3] \rangle, \langle d, [0.6, 0.7], [0.2, 0.4], [0.1, 0.3] \rangle \}$ and $B_2 = \{ \langle cd, [0.5, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$.




Figure 15: Interval valued neutrosophic G_2 .

$G_1[G_2] = (A_1 \circ A_2, B_1 \circ B_2)$ is a SIVNG. Where

$A_1 \circ A_2 = \{ \langle (a, c), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle (a, d), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle (b, c), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle (b, d), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$ and

$B_1 \circ B_2 = \{ \langle ((a, c), (a, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle ((a, c), (b, c)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle ((b, c), (b, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle ((a, d), (b, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle ((a, c), (b, d)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle, \langle ((a, d), (b, c)), [0.4, 0.6], [0.2, 0.4], [0.2, 0.4] \rangle \}$. In this example, G_1 is a SIVIFG and G_2 is not a SIVNG, then $G_1[G_2]$ is a SIVNG.




Figure 16: Composition of G_1 and G_2

Proposition 3.18 Let G_1 be a strong interval valued neutrosophic graph. Then for any

interval valued neutrosophic graph G_2 , $G_1 [G_2]$ is strong interval valued neutrosophic graph iff

$$T_{A_1L}(x_1) \leq T_{B_1L}(x_2y_2), I_{A_1L}(x_1) \geq I_{B_1L}(x_2y_2) \text{ and } F_{A_1L}(x_1) \geq F_{B_1L}(x_2y_2),$$

$$T_{A_1U}(x_1) \leq T_{B_1U}(x_2y_2), I_{A_1U}(x_1) \geq I_{B_1U}(x_2y_2) \text{ and } F_{A_1U}(x_1) \geq F_{B_1U}(x_2y_2),$$

$$\forall x_1 \in V_1, x_2y_2 \in E_2.$$

Definition 3.19 Let A_1 and A_2 be interval valued neutrosophic subsets of V_1 and V_2 respectively. Let B_1 and B_2 interval valued neutrosophic subsets of E_1 and E_2 respectively. The join of two strong interval valued neutrosophic graphs G_1 and G_2 is denoted by $G_1 + G_2 = (A_1 + A_2, B_1 + B_2)$ and is defined as follows:

$$1) \quad (T_{A_1L} + T_{A_2L})(x) = \begin{cases} (T_{A_1L} \cup T_{A_2L})(x) & \text{if } x \in V_1 \cup V_2 \\ T_{A_1L}(x) & \text{if } x \in V_1 \\ T_{A_2L}(x) & \text{if } x \in V_2 \end{cases}$$

$$(T_{A_1U} + T_{A_2U})(x) = \begin{cases} (T_{A_1U} \cup T_{A_2U})(x) & \text{if } x \in V_1 \cup V_2 \\ T_{A_1U}(x) & \text{if } x \in V_1 \\ T_{A_2U}(x) & \text{if } x \in V_2 \end{cases}$$

$$(I_{A_1L} + I_{A_2L})(x) = \begin{cases} (I_{A_1L} \cap I_{A_2L})(x) & \text{if } x \in V_1 \cup V_2 \\ I_{A_1L}(x) & \text{if } x \in V_1 \\ I_{A_2L}(x) & \text{if } x \in V_2 \end{cases}$$

$$(I_{A_1U} + I_{A_2U})(x) = \begin{cases} (I_{A_1U} \cap I_{A_2U})(x) & \text{if } x \in V_1 \cup V_2 \\ I_{A_1U}(x) & \text{if } x \in V_1 \\ I_{A_2U}(x) & \text{if } x \in V_2 \end{cases}$$

$$(F_{A_1L} + F_{A_2L})(x) = \begin{cases} (F_{A_1L} \cap F_{A_2L})(x) & \text{if } x \in V_1 \cup V_2 \\ F_{A_1L}(x) & \text{if } x \in V_1 \\ F_{A_2L}(x) & \text{if } x \in V_2 \end{cases}$$

$$(F_{A_1U} + F_{A_2U})(x) = \begin{cases} (F_{A_1U} \cap F_{A_2U})(x) & \text{if } x \in V_1 \cup V_2 \\ F_{A_1U}(x) & \text{if } x \in V_1 \\ F_{A_2U}(x) & \text{if } x \in V_2 \end{cases}$$

$$2) (T_{B_1L} + T_{B_2L})(xy) = \begin{cases} (T_{B_1L} \cup T_{B_2L})(xy) & \text{if } xy \in E_1 \cup E_2 \\ T_{B_1L}(xy) & \text{if } xy \in E_1 \\ T_{B_2L}(xy) & \text{if } xy \in E_2 \end{cases}$$

$$(T_{B_1U} + T_{B_2U})(xy) = \begin{cases} (T_{B_1U} \cup T_{B_2U})(xy) & \text{if } xy \in E_1 \cup E_2 \\ T_{B_1U}(xy) & \text{if } xy \in E_1 \\ T_{B_2U}(xy) & \text{if } xy \in E_2 \end{cases}$$

$$(I_{B_1L} + I_{B_2L})(xy) = \begin{cases} (I_{B_1L} \cap I_{B_2L})(xy) & \text{if } xy \in E_1 \cup E_2 \\ I_{B_1L}(xy) & \text{if } xy \in E_1 \\ I_{B_2L}(xy) & \text{if } xy \in E_2 \end{cases}$$

$$(I_{B_1U} + I_{B_2U})(xy) = \begin{cases} (I_{B_1U} \cap I_{B_2U})(xy) & \text{if } xy \in E_1 \cup E_2 \\ I_{B_1U}(xy) & \text{if } xy \in E_1 \\ I_{B_2U}(xy) & \text{if } xy \in E_2 \end{cases}$$

$$(F_{B_1L} + F_{B_2L})(xy) = \begin{cases} (F_{B_1L} \cap F_{B_2L})(xy) & \text{if } xy \in E_1 \cup E_2 \\ F_{B_1L}(xy) & \text{if } xy \in E_1 \\ F_{B_2L}(xy) & \text{if } xy \in E_2 \end{cases}$$

$$(F_{B_1U} + F_{B_2U})(xy) = \begin{cases} (F_{B_1U} \cap F_{B_2U})(xy) & \text{if } xy \in E_1 \cup E_2 \\ F_{B_1U}(xy) & \text{if } xy \in E_1 \\ F_{B_2U}(xy) & \text{if } xy \in E_2 \end{cases}$$

$$3) (T_{B_1L} + T_{B_2L})(xy) = \min(T_{B_1L}(x), T_{B_2L}(x))$$

$$(T_{B_1U} + T_{B_2U})(xy) = \min(T_{B_1U}(x), T_{B_2U}(x))$$

$$(I_{B_1L} + I_{B_2L})(xy) = \max(I_{B_1L}(x), I_{B_2L}(x))$$

$$(I_{B_1U} + I_{B_2U})(xy) = \max(I_{B_1U}(x), I_{B_2U}(x))$$

$$(F_{B_1L} + F_{B_2L})(xy) = \max(F_{B_1L}(x), F_{B_2L}(x))$$

$(F_{B_1U} + F_{B_2U})(xy) = \max(F_{B_1U}(x), F_{B_2U}(x))$ if $xy \in E'$, where E' is the set of all edges joining the nodes of V_1 and V_2 and where we assume $V_1 \cap V_2 = \emptyset$.

4. Conclusion

Interval valued neutrosophic sets is a generalization of the notion of fuzzy sets and intuitionistic fuzzy sets, interval valued fuzzy sets, interval valued intuitionistic fuzzy sets and single valued neutrosophic sets. Interval valued neutrosophic models gives more precisions, flexibility and compatibility to the system as compared to the classical, fuzzy, intuitionistic fuzzy and single valued neutrosophic models. In this paper, we have discussed a sub class of interval valued neutrosophic graph called strong interval valued neutrosophic graph, and we have introduced some operations such as, cartesian product, composition and join of two strong interval valued neutrosophic graph with proofs. In future study, we plan to extend our research to regular interval valued neutrosophic graphs, irregular interval valued neutrosophic.

5. References

- [1] A. V. Devadoss, A. Rajkumar & N. J. P .Praveena, A Study on Miracles through Holy Bible using Neutrosophic Cognitive Maps (NCMS). International Journal of Computer Applications, 69(3) (2013).
- [2] A. Mohamed Ismayil and A. Mohamed Ali, On Strong Interval-Valued Intuitionistic Fuzzy Graph International Journal of Fuzzy Mathematics and Systems. Volume 4, Number 2 (2014), 161-168.
- [3] A. Aydoğdu, On Similarity and Entropy of Single Valued Neutrosophic Sets, Gen. Math. Notes, Vol. 29, No. 1, July 2015, pp. 67-74.
- [4] A. Q. Ansari, R. Biswas & S. Aggarwal, (2012). Neutrosophic classifier: An extension of fuzzy classifier. Elsevier- Applied Soft Computing, 13 (2013) 563-573, <http://dx.doi.org/10.1016/j.asoc.2012.08.002>
- [5] A. Q. Ansari, R. Biswas & S. Aggarwal. (Poster Presentation) Neutrosophication of Fuzzy Models, IEEE Workshop On Computational Intelligence: Theories, Applications and Future Directions (hosted by IIT Kanpur), 14th July'13.
- [6] A. Q. Ansari, R. Biswas & S. Aggarwal Extension to fuzzy logic representation: Moving towards neutrosophic logic - A new laboratory rat, Fuzzy Systems (FUZZ), 2013 IEEE International Conference, 1-8, DOI:10.1109/FUZZ-IEEE.2013.6622412.
- [7] A. Nagoor Gani . and M. Basheer Ahamed, Order and Size in Fuzzy Graphs, Bulletin of Pure and Applied Sciences, Vol 22E (No.1) (2003) 145-148.
- [8] A. Nagoor Gani. A and S. Shajitha Begum, Degree, Order and Size in Intuitionistic Fuzzy Graphs, International Journal of Algorithms, Computing and Mathematics, (3)3 (2010).
- [9] A. Nagoor Gani and S. R. Latha, On Irregular Fuzzy Graphs, Applied Mathematical Sciences, Vol.6, 2012, no.11, 517-523.
- [10] F. Smarandache. Refined Literal Indeterminacy and the Multiplication

- Law of Sub-Indeterminacies, Neutrosophic Sets and Systems, Vol. 9, 58- 63, 2015,
- [11] F. Smarandache, Types of Neutrosophic Graphs and neutrosophic Algebraic Structures together with their Applications in Technology, seminar, Universitatea Transilvania din Brasov, Facultatea de Design de Produs si Mediu, Brasov, Romania 06 June 2015.
- [12] F. Smarandache, Symbolic Neutrosophic Theory, Europeanova asbl and the author, 195p, 2015.
- [13] F. Smarandache , Neutrosophic set - a generalization of the intuitionistic fuzzy set, Granular Computing, 2006 IEEE International Conference, 38 – 42, 2006, DOI: 10.1109/GRC.2006.1635754.
- [14] F. Smarandache, A geometric interpretation of the neutrosophic set — A generalization of the intuitionistic fuzzy set Granular Computing (GrC), 2011 IEEE International Conference , 602 – 606, 2011, DOI 10.1109/GRC.2011.6122665.
- [15] Gaurav Garg, Kanika Bhutani, Megha Kumar and Swati Aggarwal, Hybrid model for medical diagnosis using Neutrosophic Cognitive Maps with Genetic Algorithms, FUZZ-IEEE 2015(IEEE International conference on fuzzy systems).
- [16] H.Wang, Y. Zhang, R. Sunderraman, Truth-value based interval neutrosophic sets, Granular Computing, IEEE International Conference , (2005) 274 - 277 Vol. 1, 2005 DOI: 10.1109/GRC.2005.1547284.
- [17] H. Wang, F. Smarandache, Y. Zhang, and R. Sunderraman, Single valued Neutrosophic Sets, Multispace and Multistructure 4 (2010) 410-413.
- [18] H.Y. Zhang , J.Q .Wang , X.H .Chen, Interval neutrosophic sets and their application in multicriteria decision making problems, The Scientific World Journal , DOI:10.1155/2014/ 645953, 2014 .
- [19] H.Y. Zhang , P. Ji, J. Q.Wang & X. HChen, An Improved Weighted Correlation Coefficient Based on Integrated Weight for Interval Neutrosophic Sets and its Application in Multi-criteria Decision-making Problems, International Journal of Computational Intelligence Systems ,V8, Issue 6, 2015, DOI:10.1080/18756891.2015.1099917
- [20] H, Zhang, J.Wang, X. Chen, An outranking approach for multi-criteria decision-making problems with interval-valued neutrosophic sets, Neural Computing and Applications, pp 1-13 (2015).
- [21] I. Deli, M. Ali, F. Smarandache , Bipolar neutrosophic sets and their application based on multi-criteria decision making problems, Advanced Mechatronic Systems (ICAMechS), International Conference, (2015) 249 - 254, DOI: 10.1109/ICAMechS.2015.7287068.
- [22] I. Turksen, Interval valued fuzzy sets based on normal forms, Fuzzy Sets and Systems, vol. 20, (1986)191-210.
- [23] J. Ye, vector similarity measures of simplified neutrosophic sets and their application in multicriteria decision making, International Journal of Fuzzy Systems, Vol. 16, No. 2, p.204-211 (2014).
- [24] J. Ye, Single-Valued Neutrosophic Minimum Spanning Tree and Its Clustering Method, Journal of Intelligent Systems 23(3): 311–324,(2014).
- [25] J. Ye Similarity measures between interval neutrosophic sets and their

applications in Multi-criteria decision-making, *Journal of Intelligent and Fuzzy Systems*, 26, 165-172 (2014).

[26] J. Ye Some aggregation operators of interval neutrosophic linguistic numbers for multiple attribute decision making, *Journal of Intelligent & Fuzzy Systems* (27) 2231-2241, (2014).

[27] K. Atanassov, "Intuitionistic fuzzy sets," *Fuzzy Sets and Systems*, vol. 20, p. 87-96 (1986).

[28] K. Atanassov and G. Gargov, "Interval valued intuitionistic fuzzy sets," *Fuzzy Sets and Systems*, vol.31, (1989) 343-349.

[29] K. Atanassov. *Intuitionistic fuzzy sets: theory and applications*. Physica, New York, 1999.

[30] L. Zadeh, *Fuzzy sets*, *Inform and Control*, 8(1965), 338-353

[31] P. Bhattacharya, Some remarks on fuzzy graphs, *Pattern Recognition Letters* 6: 297-302, 1987.

[32] P. Liu and L. Shi The generalized hybrid weighted average operator based on interval neutrosophic hesitant set and its application to multiple attribute decision making, *Neural Computing and Applications*, 26 (2) (2015) 457-471.

[33] R. Parvathi and M. G. Karunambigai, *Intuitionistic Fuzzy Graphs, Computational Intelligence, Theory and applications*, International Conference in Germany, Sept 18 -20, 2006.

[34] R. Rıdvan, A. Küçük, Subsethood measure for single valued neutrosophic sets, *Journal of Intelligent & Fuzzy Systems*, vol. 29, no. 2, pp. 525-530, 2015, DOI: 10.3233/IFS-141304.

[35] R. Şahin, Cross-entropy measure on interval neutrosophic sets and its applications in multicriteria decision making, *Neural Computing and Applications*, pp 1-11 (2015).

[36] S. Aggarwal, R. Biswas, A. Q. Ansari, *Neutrosophic modeling and control ,Computer and Communication Technology (ICCCT)*, International Conference , (2010) 718 – 723, DOI:10.1109/ICCCT.2010.5640435

[37] S. Broumi, F. Smarandache, New distance and similarity measures of interval neutrosophic sets, *Information Fusion (FUSION)*, 2014 IEEE 17th International Conference, 2014,p 1 – 7.

[38] S. Broumi, F. Smarandache, Single valued neutrosophic trapezoid linguistic aggregation operators based multi-attribute decision making, *Bulletin of Pure & Applied Sciences- Mathematics and Statistics*,(2014)135-155, DOI : 10.5958/2320-3226.2014.00006.X.

[39] S.Broumi, M. Talea, F. Smarandache, *Single Valued Neutrosophic Graphs: Degree, Order and Size*, (2016) submitted

[40] S.Broumi, M. Talea, A.Bakali, F. Smarandache, *Single Valued Neutrosophic Graphs*, *Journal of new theory* , N 10(2016) 86-101

[41] S.Broumi, M. Talea, A.Bakali, F. Smarandache, *Interval Valued Neutrosophic Graphs*, critical review (2016) in presse

[42] S.Broumi, M. Talea, A.Bakali, F. Smarandache, *Operations on Interval Valued Neutrosophic Graphs*, (2016) submitted.

[43] Y. Hai-Long, G. She, Yanhonge, L. Xiuwu, On single valued neutrosophic relations, *Journal of Intelligent & Fuzzy Systems*, vol. Preprint, no. Preprint, p. 1-12 (2015)

[44] W. B. Vasantha Kandasamy and F. Smarandache, Fuzzy Cognitive Maps and Neutrosophic Congtive Maps,2013.

[45] W. B. Vasantha Kandasamy, K. Ilanthenral and Florentin Smarandache, Neutrosophic Graphs: A New Dimension to Graph Theory, Kindle Edition, 2015.

[46] W.B. Vasantha Kandasamy and F. Smarandache “Analysis of social aspects of migrant laborers living with HIV/AIDS using Fuzzy Theory and Neutrosophic Cognitive Maps”, Xiquan, Phoenix (2004).