

Does a classical probability space for two-dimensional quantum measurement theory exist?

Koji Nagata¹ and Tadao Nakamura²

¹*Department of Physics, Korea Advanced Institute of Science and Technology, Daejeon 305-701, Korea*

E-mail: ko_mi_na@yahoo.co.jp

²*Department of Information and Computer Science, Keio University,*

3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan

E-mail: nakamura@pipelining.jp

(Dated: June 8, 2016)

Recently, a new measurement theory based on the truth values is proposed [37]. The results of measurements are either 0 or 1. The measurement theory accepts a hidden variables model for a single Pauli observable. Therefore we can introduce a classical probability space for the measurement theory in this case. On the other hand, we discuss the fact that the projective measurement theory (the results of measurements are either +1 or -1) does not meet a hidden variables model for a single Pauli observable. Hence we cannot introduce a classical probability space for the projective measurement theory in this case. Our discussion provides new insight to formulate quantum measurement theory, by using the measurement theory based on the truth values.

PACS numbers: 03.65.Ta (Quantum measurement theory), 03.65.Ud (Quantum non locality), 03.65.Ca (Formalism)

I. INTRODUCTION

The projective measurement theory (cf. [1–6]) gives accurate and at times remarkably accurate numerical predictions. Much experimental data has fit to the quantum predictions for long time.

From the incompleteness argument of Einstein, Podolsky, and Rosen (EPR) [7], a hidden-variables interpretation of quantum mechanics has been an attractive topic of research [3, 4]. One is the Bell-EPR theorem [8]. The other is the no-hidden-variables theorem of Kochen and Specker (the KS theorem) [9]. Greenberger, Horne, and Zeilinger discover [10, 11] the so-called GHZ theorem for four-partite GHZ state. And, the Bell-KS theorem becomes very simple form (see also Refs. [12–16]).

Leggett-type nonlocal hidden-variable theory [17] is experimentally investigated [18–20]. The experiments report that the quantum theory does not accept Leggett-type nonlocal hidden-variable theory. These experiments are done in four-dimensional space (two parties) in order to study nonlocality of hidden-variable theories. However there are debates for the conclusions of the experiments. See Refs. [21–23].

As for the applications of the projective measurement theory, implementation of a quantum algorithm to solve Deutsch's problem [24, 25] on a nuclear magnetic resonance quantum computer is reported firstly [26]. Implementation of the Deutsch-Jozsa algorithm on an ion-trap quantum computer is also reported [27]. There are several attempts to use single-photon two-qubit states for quantum computing. Oliveira *et al.* implement Deutsch's algorithm with polarization and transverse spatial modes of the electromagnetic field as qubits [28]. Single-photon Bell states are prepared and measured [29]. Also the decoherence-free implementation of Deutsch's algorithm is reported by using such single-photon and

by using two logical qubits [30]. More recently, a one-way based experimental implementation of Deutsch's algorithm is reported [31]. In 1993, the Bernstein-Vazirani algorithm was reported [32]. It can be considered as an extended Deutsch-Jozsa algorithm. In 1994, Simon's algorithm was reported [33]. Implementation of a quantum algorithm to solve the Bernstein-Vazirani parity problem without entanglement on an ensemble quantum computer is reported [34]. Fiber-optics implementation of the Deutsch-Jozsa and Bernstein-Vazirani quantum algorithms with three qubits is discussed [35]. A quantum algorithm for approximating the influences of Boolean functions and its applications is recently reported [36].

More recently, a new measurement theory based on the truth values is proposed [37]. The results of measurements are either 0 or 1. The measurement theory accepts a hidden variables model for a single Pauli observable. Therefore we can introduce a classical probability space for the measurement theory in this case. On the other hand, we discuss the fact that the projective measurement theory (the results of measurements are either +1 or -1) does not meet a hidden variables model for a single Pauli observable. Hence we cannot introduce a classical probability space for the projective measurement theory in this case. Our discussion provides new insight to formulate quantum measurement theory, by using the measurement theory based on the truth values.

II. MEASUREMENT THEORY BASED ON THE TRUTH VALUES MEETS A HIDDEN VARIABLES MODEL OF A SINGLE SPIN OBSERVABLE

We discuss the new measurement theory meets a hidden variables model of a single spin observable. Assume

a spin-1/2 state ρ . Let σ_x be a single Pauli observable. We have a quantum expected value as

$$\text{Tr}[\rho\sigma_x]. \quad (1)$$

We derive a necessary condition for the quantum expected value for the system in a spin-1/2 state given in (1). We have

$$0 \leq (\text{Tr}[\rho\sigma_x])^2 \leq 1. \quad (2)$$

It is worth noting here that we have $(\text{Tr}[\rho\sigma_x])^2 = 1$ if ρ is a pure state lying in the x -direction. Hence we derive the following proposition concerning quantum mechanics when the system is in a state lying in the x -direction

$$(\text{Tr}[\rho\sigma_x])_{\max}^2 = 1. \quad (3)$$

$(\text{Tr}[\rho\sigma_x])_{\max}^2$ is the maximal possible value of the product. It is worth noting here that we have $(\text{Tr}[\rho\sigma_x])^2 = 0$ when the system is in a pure state lying in the z -direction. Thus we have

$$(\text{Tr}[\rho\sigma_x])_{\min}^2 = 0. \quad (4)$$

$(\text{Tr}[\rho\sigma_x])_{\min}^2$ is the minimal possible value of the product. In short, we have

$$(\text{Tr}[\rho\sigma_x])_{\min}^2 = 0 \text{ and } (\text{Tr}[\rho\sigma_x])_{\max}^2 = 1. \quad (5)$$

In what follows, we derive the above proposition (5) assuming the following form:

$$\text{Tr}[\rho\sigma_x] = \int d\lambda \rho(\lambda) f(\sigma_x, \lambda). \quad (6)$$

where λ denotes some hidden variable and $f(\sigma_x, \lambda)$ is the hidden result of measurements of the Pauli observable σ_x . We assume the values of $f(\sigma_x, \lambda)$ are 1 and 0 (in $\hbar/2$ unit).

Let us assume a hidden variables theory of a single spin observable based on the new measurement theory. In this case, the quantum expected value in (1), which is the average of the hidden results of the new measurements, is given by

$$\text{Tr}[\rho\sigma_x] = \int d\lambda \rho(\lambda) f(\sigma_x, \lambda). \quad (7)$$

The possible values of the hidden result $f(\sigma_x, \lambda)$ are 1 and 0 (in $\hbar/2$ unit). Same expected value is given by

$$\text{Tr}[\rho\sigma_x] = \int d\lambda' \rho(\lambda') f(\sigma_x, \lambda'), \quad (8)$$

because we only change the notation as $\lambda \rightarrow \lambda'$. Of course, the possible values of the hidden result $f(\sigma_x, \lambda')$ are 1 and 0 (in $\hbar/2$ unit). By using these facts, we derive a necessary condition for the expected value for the system in a spin-1/2 state lying in the x -direction. We derive the

possible values of the product $(\text{Tr}[\rho\sigma_x])^2$. We have

$$\begin{aligned} & (\text{Tr}[\rho\sigma_x])^2 \\ &= \int d\lambda \rho(\lambda) f(\sigma_x, \lambda) \times \int d\lambda' \rho(\lambda') f(\sigma_x, \lambda') \\ &= \int d\lambda \rho(\lambda) \cdot \int d\lambda' \rho(\lambda') f(\sigma_x, \lambda) f(\sigma_x, \lambda') \\ &\leq \int d\lambda \rho(\lambda) \cdot \int d\lambda' \rho(\lambda') |f(\sigma_x, \lambda) f(\sigma_x, \lambda')| \\ &= \int d\lambda \rho(\lambda) \cdot \int d\lambda' \rho(\lambda') = 1. \end{aligned} \quad (9)$$

Clearly, the above inequality can have the upper limit since the following case is possible:

$$\|\{\lambda|f(\sigma_x, \lambda) = 1\}\| = \|\{\lambda'|f(\sigma_x, \lambda') = 1\}\|, \quad (10)$$

and

$$\|\{\lambda|f(\sigma_x, \lambda) = 0\}\| = \|\{\lambda'|f(\sigma_x, \lambda') = 0\}\|. \quad (11)$$

Thus we derive a proposition concerning the hidden variables theory based on the new measurement theory (in a spin-1/2 system), that is, $(\text{Tr}[\rho\sigma_x])^2 \leq 1$. Hence we derive the following proposition concerning the hidden variables theory:

$$(\text{Tr}[\rho\sigma_x])_{\max}^2 = 1. \quad (12)$$

We derive another necessary condition for the expected value for the system in a pure spin-1/2 state lying in the z -direction. We have

$$\begin{aligned} & (\text{Tr}[\rho\sigma_x])^2 \\ &= \int d\lambda \rho(\lambda) f(\sigma_x, \lambda) \times \int d\lambda' \rho(\lambda') f(\sigma_x, \lambda') \\ &= \int d\lambda \rho(\lambda) \cdot \int d\lambda' \rho(\lambda') f(\sigma_x, \lambda) f(\sigma_x, \lambda') \\ &\geq \int d\lambda \rho(\lambda) \cdot \int d\lambda' \rho(\lambda') (0) \\ &= (0) \left(\int d\lambda \rho(\lambda) \cdot \int d\lambda' \rho(\lambda') \right) = 0. \end{aligned} \quad (13)$$

Clearly, the above inequality can have the lower limit since the following case is possible:

$$\|\{\lambda|f(\sigma_x, \lambda) = 1\}\| = \|\{\lambda'|f(\sigma_x, \lambda') = 0\}\|, \quad (14)$$

and

$$\|\{\lambda|f(\sigma_x, \lambda) = 0\}\| = \|\{\lambda'|f(\sigma_x, \lambda') = 1\}\|. \quad (15)$$

Thus we derive a proposition concerning the hidden variables theory based on the new measurement theory (in a spin-1/2 system), that is, $(\text{Tr}[\rho\sigma_x])^2 \geq 0$. Hence we derive the following proposition concerning the hidden variables theory

$$(\text{Tr}[\rho\sigma_x])_{\min}^2 = 0. \quad (16)$$

Thus from (12) and (16) we have

$$(\text{Tr}[\rho\sigma_x])_{\min}^2 = 0 \text{ and } (\text{Tr}[\rho\sigma_x])_{\max}^2 = 1. \quad (17)$$

Clearly, we can assign the truth value “1” for two propositions (5) (concerning quantum mechanics) and (17) (concerning the hidden variables theory based on the new measurement theory), simultaneously. Therefore, the new measurement theory meets the existence of the hidden variables theory of a single spin observable.

III. PROJECTIVE MEASUREMENT THEORY DOES NOT MEET A HIDDEN VARIABLES MODEL OF A SINGLE SPIN OBSERVABLE

In what follows, we cannot derive the proposition (5) assuming the following form:

$$\text{Tr}[\rho\sigma_x] = \int d\lambda\rho(\lambda)f(\sigma_x, \lambda). \quad (18)$$

where λ denotes some hidden variable and $f(\sigma_x, \lambda)$ is the hidden result of measurements of the Pauli observable σ_x . We assume the values of $f(\sigma_x, \lambda)$ are +1 and -1 (in $\hbar/2$ unit).

Let us assume a hidden variables model based on the projective measurement theory of a single spin observable. In this case, the quantum expected value in (1), which is the average of the hidden results of the projective measurements, is given by

$$\text{Tr}[\rho\sigma_x] = \int d\lambda\rho(\lambda)f(\sigma_x, \lambda). \quad (19)$$

The possible values of the hidden result $f(\sigma_x, \lambda)$ are +1 and -1 (in $\hbar/2$ unit). Same expected value is given by

$$\text{Tr}[\rho\sigma_x] = \int d\lambda'\rho(\lambda')f(\sigma_x, \lambda'), \quad (20)$$

because we only change the notation as $\lambda \rightarrow \lambda'$. Of course, the possible values of the hidden result $f(\sigma_x, \lambda')$ are +1 and -1 (in $\hbar/2$ unit). By using these facts, we derive a necessary condition for the expected value for the system in a spin-1/2 state lying in the x -direction. We derive the possible values of the product $(\text{Tr}[\rho\sigma_x])^2$. We have

$$\begin{aligned} & (\text{Tr}[\rho\sigma_x])^2 \\ &= \int d\lambda\rho(\lambda)f(\sigma_x, \lambda) \times \int d\lambda'\rho(\lambda')f(\sigma_x, \lambda') \\ &= \int d\lambda\rho(\lambda) \cdot \int d\lambda'\rho(\lambda')f(\sigma_x, \lambda)f(\sigma_x, \lambda') \\ &\leq \int d\lambda\rho(\lambda) \cdot \int d\lambda'\rho(\lambda')|f(\sigma_x, \lambda)f(\sigma_x, \lambda')| \\ &= \int d\lambda\rho(\lambda) \cdot \int d\lambda'\rho(\lambda') = 1. \end{aligned} \quad (21)$$

Clearly, the above inequality can have the upper limit since the following case is possible:

$$\|\{\lambda|f(\sigma_x, \lambda) = 1\}\| = \|\{\lambda'|f(\sigma_x, \lambda') = 1\}\|, \quad (22)$$

and

$$\|\{\lambda|f(\sigma_x, \lambda) = 0\}\| = \|\{\lambda'|f(\sigma_x, \lambda') = 0\}\|. \quad (23)$$

Thus we derive a proposition concerning the hidden variables theory based on the projective measurement theory (in a spin-1/2 system), that is, $(\text{Tr}[\rho\sigma_x])^2 \leq 1$. Hence we derive the following proposition concerning the hidden variables theory

$$(\text{Tr}[\rho\sigma_x])_{\text{max}}^2 = 1. \quad (24)$$

We derive another necessary condition for the expected value for the system in a pure spin-1/2 state lying in the z -direction. We have

$$\begin{aligned} & (\text{Tr}[\rho\sigma_x])^2 \\ &= \int d\lambda\rho(\lambda)f(\sigma_x, \lambda) \times \int d\lambda'\rho(\lambda')f(\sigma_x, \lambda') \\ &= \int d\lambda\rho(\lambda) \cdot \int d\lambda'\rho(\lambda')f(\sigma_x, \lambda)f(\sigma_x, \lambda') \\ &\geq \int d\lambda\rho(\lambda) \cdot \int d\lambda'\rho(\lambda')(-1) \\ &= (-1) \left(\int d\lambda\rho(\lambda) \cdot \int d\lambda'\rho(\lambda') \right) = -1. \end{aligned} \quad (25)$$

Clearly, the above inequality can have the lower limit since the following case is possible:

$$\|\{\lambda|f(\sigma_x, \lambda) = 1\}\| = \|\{\lambda'|f(\sigma_x, \lambda') = -1\}\|, \quad (26)$$

and

$$\|\{\lambda|f(\sigma_x, \lambda) = -1\}\| = \|\{\lambda'|f(\sigma_x, \lambda') = 1\}\|. \quad (27)$$

Thus we derive a proposition concerning the hidden variables theory based on the projective measurement theory (in a spin-1/2 system), that is, $(\text{Tr}[\rho\sigma_x])^2 \geq -1$. Hence we derive the following proposition concerning the hidden variables theory

$$(\text{Tr}[\rho\sigma_x])_{\text{min}}^2 = -1. \quad (28)$$

Thus from (24) and (28) we have

$$(\text{Tr}[\rho\sigma_x])_{\text{min}}^2 = -1 \text{ and } (\text{Tr}[\rho\sigma_x])_{\text{max}}^2 = 1. \quad (29)$$

Clearly, we cannot assign the truth value “1” for two propositions (5) (concerning quantum mechanics) and (29) (concerning the hidden variables theory based on the projective measurement theory), simultaneously. Therefore, the projective measurement theory does not meet the existence of the hidden variables theory of a single spin observable.

IV. CONCLUSIONS

In conclusions, recently, a new measurement theory based on the truth values has been proposed [37]. The results of measurements have been either 0 or 1. The measurement theory has accepted a hidden variables model

for a single Pauli observable. Therefore we can have introduced a classical probability space for the measurement theory in this case. On the other hand, we have discussed the fact that the projective measurement theory (the results of measurements are either $+1$ or -1)

does not meet a hidden variables model for a single Pauli observable. Hence we cannot have introduced a classical probability space for the projective measurement theory in this case. Our discussion has provided new insight to formulate quantum measurement theory, by using the measurement theory based on the truth values.

-
- [1] J. von Neumann, *Mathematical Foundations of Quantum Mechanics* (Princeton University Press, Princeton, New Jersey, 1955).
- [2] J. J. Sakurai, *Modern Quantum Mechanics* (Addison-Wesley Publishing Company, 1995), Revised ed.
- [3] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer Academic, Dordrecht, The Netherlands, 1993).
- [4] M. Redhead, *Incompleteness, Nonlocality, and Realism* (Clarendon Press, Oxford, 1989), 2nd ed.
- [5] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2000).
- [6] R. P. Feynman, R. B. Leighton, and M. Sands, *Lectures on Physics, Volume III, Quantum mechanics* (Addison-Wesley Publishing Company, 1965).
- [7] A. Einstein, B. Podolsky, and N. Rosen, *Phys. Rev.* **47**, 777 (1935).
- [8] J. S. Bell, *Physics* **1**, 195 (1964).
- [9] S. Kochen and E. P. Specker, *J. Math. Mech.* **17**, 59 (1967).
- [10] D. M. Greenberger, M. A. Horne, and A. Zeilinger, in *Bell's Theorem, Quantum Theory and Conceptions of the Universe*, edited by M. Kafatos (Kluwer Academic, Dordrecht, The Netherlands, 1989), pp. 69-72.
- [11] D. M. Greenberger, M. A. Horne, A. Shimony, and A. Zeilinger, *Am. J. Phys.* **58**, 1131 (1990).
- [12] C. Pagonis, M. L. G. Redhead, and R. K. Clifton, *Phys. Lett. A* **155**, 441 (1991).
- [13] N. D. Mermin, *Phys. Today* **43**(6), 9 (1990).
- [14] N. D. Mermin, *Am. J. Phys.* **58**, 731 (1990).
- [15] A. Peres, *Phys. Lett. A* **151**, 107 (1990).
- [16] N. D. Mermin, *Phys. Rev. Lett.* **65**, 3373 (1990).
- [17] A. J. Leggett, *Found. Phys.* **33**, 1469 (2003).
- [18] S. Gröblacher, T. Paterek, R. Kaltenbaek, Č. Brukner, M. Żukowski, M. Aspelmeyer, and A. Zeilinger, *Nature (London)* **446**, 871 (2007).
- [19] T. Paterek, A. Fedrizzi, S. Gröblacher, T. Jennewein, M. Żukowski, M. Aspelmeyer, and A. Zeilinger, *Phys. Rev. Lett.* **99**, 210406 (2007).
- [20] C. Branciard, A. Ling, N. Gisin, C. Kurtsiefer, A. Lamas-Linares, and V. Scarani, *Phys. Rev. Lett.* **99**, 210407 (2007).
- [21] A. Suarez, *Found. Phys.* **38**, 583 (2008).
- [22] M. Żukowski, *Found. Phys.* **38**, 1070 (2008).
- [23] A. Suarez, *Found. Phys.* **39**, 156 (2009).
- [24] D. Deutsch, *Proc. Roy. Soc. London Ser. A* **400**, 97 (1985).
- [25] D. Deutsch and R. Jozsa, *Proc. Roy. Soc. London Ser. A* **439**, 553 (1992).
- [26] J. A. Jones and M. Mosca, *J. Chem. Phys.* **109**, 1648 (1998).
- [27] S. Gulde, M. Riebe, G. P. T. Lancaster, C. Becher, J. Eschner, H. Häffner, F. Schmidt-Kaler, I. L. Chuang, and R. Blatt, *Nature (London)* **421**, 48 (2003).
- [28] A. N. de Oliveira, S. P. Walborn, and C. H. Monken, *J. Opt. B: Quantum Semiclass. Opt.* **7**, 288-292 (2005).
- [29] Y.-H. Kim, *Phys. Rev. A* **67**, 040301(R) (2003).
- [30] M. Mohseni, J. S. Lundeen, K. J. Resch, and A. M. Steinberg, *Phys. Rev. Lett.* **91**, 187903 (2003).
- [31] M. S. Tame, R. Prevedel, M. Paternostro, P. Böhi, M. S. Kim, and A. Zeilinger, *Phys. Rev. Lett.* **98**, 140501 (2007).
- [32] E. Bernstein and U. Vazirani, *Proceedings of the Twenty-Fifth Annual ACM Symposium on Theory of Computing (STOC '93)*, pp. 11-20 (1993), doi:10.1145/167088.167097; *SIAM J. Comput.* **26**-5, pp. 1411-1473 (1997).
- [33] D. R. Simon, *Foundations of Computer Science*, (1994) *Proceedings.*, 35th Annual Symposium on: 116-123, retrieved 2011-06-06.
- [34] J. Du, M. Shi, X. Zhou, Y. Fan, B. J. Ye, R. Han, and J. Wu, *Phys. Rev. A* **64**, 042306 (2001).
- [35] E. Brainis, L.-P. Lamoureaux, N. J. Cerf, Ph. Emplit, M. Haelterman, and S. Massar, *Phys. Rev. Lett.* **90**, 157902 (2003).
- [36] H. Li and L. Yang, *Quantum Inf. Process.* **14**, 1787 (2015).
- [37] K. Nagata and T. Nakamura, *Int. J. Theor. Phys.* DOI 10.1007/s10773-016-2990-2, (2016).