

**Initial mass in Pre Planckian space-timed defined, and causal discontinuity.**

**Andrew Walcott Beckwith**

Physics Department, Chongqing University, College of Physics, Chongqing  
University Huxi Campus, No. 44 Daxuechen Nanlu, Shapinba District, Chongqing  
401331, People's Republic of China

[abeckwith@uh.edu](mailto:abeckwith@uh.edu), [rwill9955b@gmail.com](mailto:rwill9955b@gmail.com)

## **Abstract**

This document reviews the Landau – Lifshits reformulation of General relativity with a representation of the available mass of a graviton. From looking at a conservation law, using the Landau-Lifshits formulation, we obtain conditions for initial mass, in the Pre Planckian regime of space-time. In doing so, we also indicate a metric tensor and metric pseudo tensor delineation of causal discontinuity.

Key words, Massive gravitons, Landau-Lifshits pseudo tensor, causal discontinuity.

## **1. Introduction**

We will use the Poisson formulation of the Landau-Lifshits formulation of General Relativity [1] in order to obtain an initial mass, in the Pre Planckian Regime of space time. Using a modification of the Heisenberg Uncertainty principle, we also obtain an inter relationship between an inflation, initially, and initial mass. Finally, we reference the issue of a causal discontinuity, with surprising implications.

We first begin by a recapitulation of the different models for HFGW as given by Dr. Li. Et al, 2008 [2]

Table 1: magnitude, sources, and top frequency values for HFGW (from Li et al. 2008) [5]

Sources	Amplitude	frequency	Characteristics
HFGW in Quintessence inflationary models	$h_{rms} \sim 10^{-30} - 10^{-32} / \sqrt{Hz}$	$\nu \sim 10^9 - 10^{10} Hz$	Random background
HFGW in some string theory scenarios	$h_{rms} \sim 10^{-30} - 10^{-34} / \sqrt{Hz}$	$\nu \sim 10^8 - 10^{11} Hz$	Random background
Solar Plasma	$h_{rms} \sim 10^{-39} / \sqrt{Hz}$	$\nu \sim 10^{15} Hz$	On the Earth
High energy particles, e.g. Fermi Ring	$h_{rms} \sim 10^{-39} - 10^{-41} / \sqrt{Hz}$	$\nu \sim 10^4 - 10^5 Hz$	On the center the frequency depends upon the rotational frequency of particles in the Fermi Ring
Stanford Linear Accelerator	$h_{rms} \sim 10^{-39} / \sqrt{Hz}$	$\nu \sim 10^{23} Hz$	On the collision center, the frequency depends upon the self-energy and the Lorentz factor of high energy e <sup>+</sup> e <sup>-</sup> beams
LHC- Large Hadron collider			Spectra of high energy gravitons
Nano-piezo electric crystal array, with size of about 100 nanometers	$h_{rms} \sim 10^{-28} - 10^{-31} / \sqrt{Hz}$	$\nu \sim 10^9 - 10^{10} Hz$	On the wave zone with an effective cross section of or less than .01 meters squared, for gravitational radiation

This re produced PRD table[5] is important since it suggests that relic GW, if properly measured, may be the first ones to await experimental verification. But in order to do this, we add in several caveats, as to after the fact considerations

1. We are considering using massive gravitons [3]
2. Our methodology suggests that if we measure relic gravitational waves, that we will have to consider if an initial mass in the evolution of the universe, actually existed.[1]
3. We also review, by example, if there is a way to use a modified version of the Heisenberg uncertainty principle, as given in [ 4 ] to obtain an inflaton, in the onset of expansion of the Universe
4. By way of our construction, we will also look at if a causal discontinuity exists as an elaboration of [ 5 ]

Having said that, it is time now to unveil by way of the Poisson reference [ 1 ] initial mass, and to tie it into our four issues brought up above. But before this, we will allude to our basic work horse, i.e. the Ng Entropy.

**1a. Entropy generation via Ng's infinite quantum statistics (short review)**

We wish to understand the linkage between dark matter and gravitons. To consider just that, we look at the “size” of the nucleation space,  $V$ .  $V$  for nucleation is HUGE. Graviton space  $V$  for nucleation is tiny, well inside inflation/ therefore, the log factor drops OUT of entropy  $S$  if  $V$  chosen properly for both Eq 1 and Eq. 2. Ng's [6] result begins with a modification of the entropy/ partition function Ng used the following approximation of temperature and its variation with respect to a spatial parameter, starting with temperature  $T \approx R_H^{-1}$  ( $R_H$  can be thought of as a representation of the region of space where we take statistics of the particles in question). Furthermore, assume that the volume of space to be analyzed is of the form  $V \approx R_H^3$  and look at a preliminary numerical factor we shall call  $N \sim (R_H/l_P)^2$ , where the denominator is Planck's length (on the order of  $10^{-35}$  centimeters). We also specify a “wavelength” parameter  $\lambda \approx T^{-1}$ . So the value of  $\lambda \approx T^{-1}$  and of  $R_H$  are approximately the same order of magnitude. Now this is how Jack Ng changes conventional statistics: he outlines how to get  $S \approx N$ , which with additional arguments we refine to be  $S \approx \langle n \rangle$  (where  $\langle n \rangle$  is graviton density). Begin with a partition function

$$Z_N \sim \left(\frac{1}{N!}\right) \cdot \left(\frac{V}{\lambda^3}\right)^N \tag{1}$$

This, according to Ng,[6] leads to entropy of the limiting value of, if  $S = (\log[Z_N])$

$$S \approx N \cdot (\log[V/N\lambda^3] + 5/2) \xrightarrow{\text{Ng-inf inite-Quantum-Statistics}} N \cdot (\log[V/\lambda^3] + 5/2) \approx N \quad (2)$$

But  $V \approx R_H^3 \approx \lambda^3$ , The modification of Ng's entropy expression [6] is in the region of space time for which the general temperature dependent entropy Kolb and Turner expression breaks down. In particular, the evaluation of entropy we do via the modified Ng argument above is in regions of space time where  $g$  before re heat is an unknown, unmeasurable number of degrees of freedom. The Kolb and Turner entropy expression [7] 1991 has a temperature  $T$  related entropy density which leads to that we are able to state total entropy as the entropy density time's space time volume  $V_4$  with  $g_{re-heat} \approx 1000$ , while dropping to  $g_{electro-weak} \approx 100$  in the electro weak era. This value of the space time degrees of freedom, according to de Vega has reached a low of  $g_{today} \approx 2-3$  today. We assert that Eq. (2) above occurs in a region of space time before  $g_{re-heat} \approx 1000$ , so after re heating Eq. (2) no longer holds, and we instead can look at [7]

$$S_{total} \equiv s_{Density} \cdot V_4 = \frac{2\pi^2}{45} \cdot g \cdot T^3 \cdot V_4 \quad (3)$$

Where  $T < 10^{32} K$ .

Such a linkage would open up the possibility that the density of primordial gravitational waves could be examined, and linked to modeling gravity as an effective theory. The details of linking what is done with Eq. (2) and bridging it to Eq. (3) await additional theoretical development, and are probably conceptually understandable if the following is used to link the two regimes. I.e. we can use the number of space time operations used to create Eq. (2), via Seth Lloyds [8]

$$I = S_{total} / k_B \ln 2 = [\#operations]^{3/4} = [\rho \cdot c^5 \cdot t^4 / \hbar]^{3/4} \quad (4)$$

Essentially, what will be done is to use Eq. (4) to show linkage between a largely thermally based production of entropy, as implied by Eq. (3) and a particle counting algorithm, as given by Eq. (2). This due to the problems inherent in making connections between a particle count generation of entropy, and thermal contributions. I.e. two different processes are involved.

## 2. The Poisson rendition of the Landau-Liftshitz effective initial mass

To do this go to [1] and the non Tensorial form of the Einstein field equations starting with the so called Gothic inverse metric, with a 2n line of Eq. (5) being a tensor identity

$$\begin{aligned} \mathfrak{S}^{\alpha\beta} &= \sqrt{-(g = \det g^{\alpha\beta})} \cdot g^{\alpha\beta} \\ H^{\alpha\mu\beta\nu} &= \mathfrak{S}^{\alpha\beta} \mathfrak{S}^{\mu\nu} - \mathfrak{S}^{\alpha\nu} \mathfrak{S}^{\mu\beta} \end{aligned} \quad (5)$$

Furthermore, the two equations in Eq. (5) have a representation of the GR field equations as

$$\partial_{\beta} \partial_{\mu\nu} H^{\alpha\mu\beta\nu} = 0 = \frac{16\pi G \cdot \partial_{\beta} \left( -(g = \det g^{\alpha\beta}) \right) \cdot [T^{\alpha\beta} + t_{LL}^{\alpha\beta}]}{c^4} \quad (6)$$

If so then, we can have a simple solution to this above which is of the type

$$[T^{\alpha\beta} + t_{LL}^{\alpha\beta}] = -\frac{c_1}{-(g = \det g^{\alpha\beta})} \quad (7)$$

Here, in this situation we have that if we are following the ideas in [4] where we have restriction to the zeroth (time component) of the metric tensor, so that

$$\delta(g_{tt} = g_{00}) = a_{initial}^2 \cdot \phi(initial) \quad (8)$$

We will be looking at the value of Eq. (7) if  $\phi \sim \xi^+ \ll M_{Planck}$ . In short, we have then that by [4]

$$\begin{aligned} (\Delta l)_{ij} &= \frac{\delta g_{ij}}{g_{ij}} \cdot \frac{l}{2} \\ (\Delta p)_{ij} &= \Delta T_{ij} \cdot \delta t \cdot \Delta A \end{aligned} \quad (9)$$

If we use the following, from the Roberson-Walker metric [9,10,11].

$$\begin{aligned}
g_{tt} &= 1 \\
g_{rr} &= \frac{-a^2(t)}{1-k \cdot r^2} \\
g_{\theta\theta} &= -a^2(t) \cdot r^2 \\
g_{\phi\phi} &= -a^2(t) \cdot \sin^2 \theta \cdot d\phi^2
\end{aligned} \tag{10}$$

Then, the surviving version of Eq. (9) and Eq. (10) is, then, if  $\Delta T_{tt} \sim \Delta\rho$  [4, 10, 11]

$$\begin{aligned}
V^{(4)} &= \delta t \cdot \Delta A \cdot r \\
\delta g_{tt} \cdot \Delta T_{tt} \cdot \delta t \cdot \Delta A \cdot \frac{r}{2} &\geq \frac{\hbar}{2} \\
\Leftrightarrow \delta g_{tt} \cdot \Delta T_{tt} &\geq \frac{\hbar}{V^{(4)}}
\end{aligned} \tag{11}$$

This Eq. (11) is such that we can extract, up to a point the HUP principle for uncertainty in time and energy, with one very large caveat added, namely if we use the fluid approximation of space-time [4]

$$T_{ii} = \text{diag}(\rho, -p, -p, -p) \tag{12}$$

Then by [10,11]

$$\Delta T_{tt} \sim \Delta\rho \sim \frac{\Delta E}{V^{(3)}} \tag{13}$$

$$\text{Then, } \delta t \Delta E \geq \frac{\hbar}{\delta g_{tt}} \neq \frac{\hbar}{2} \tag{14}$$

$$\text{Unless } \delta g_{tt} \sim O(1)$$

Here, we have a causal discontinuity as given by

$$\begin{aligned}
M &= \frac{1}{c^2} \int_V (-g)(T^{00} + g_{LL}^{00}) \cdot d^3x = \frac{c_1}{c^2} \cdot \int_V d^3x \sim \frac{\hbar}{g_{00} \cdot \delta t} = \frac{\hbar}{a_{\min}^2 \phi_{\min} \cdot \delta t} \\
[T^{00} + t_{LL}^{00}] &= - \frac{\hbar}{a_{\min}^2 \phi_{\min} \cdot \delta t|_{\text{Pre-Planckian}}} \propto \left( - \frac{\hbar}{\delta t|_{\text{Planckian}}} \right) = [T^{\alpha\beta} + t_{LL}^{\alpha\beta}] \\
\Leftrightarrow T^{00}|_{\text{Pre-Planckian}} - T^{\alpha\beta}|_{\text{Planckian}} &\propto t_{LL}^{\alpha\beta}|_{\text{Planckian}} - t_{LL}^{00}|_{\text{Pre-Planckian}}
\end{aligned} \tag{15}$$

We will address the implications of Eq. (15) in the conclusion

If we then put in the initial mass, of say from [1] of

$$M = \frac{1}{c^2} \int_V (-g)(T^{00} + g^{00}) \cdot d^3x = \frac{c_1}{c^2} \cdot \int_V d^3x \sim \frac{\hbar}{g_{00} \cdot \delta t} = \frac{\hbar}{a_{\min}^2 \phi_{nub} \cdot \delta t} \quad (15)$$

With the minimum scale factor a small, but non zero factor by [12] which is shown up in

$$\begin{aligned} [T^{00} + t_{LL}^{00}] &= - \cdot \frac{\hbar}{a_{\min}^2 \phi_{\min} \cdot \delta t|_{\text{Pre-Planckian}}} \propto \left( - \frac{\hbar}{\delta t|_{\text{Planckian}}} \right) = [T^{\alpha\beta} + t_{LL}^{\alpha\beta}] \quad (16) \\ \Leftrightarrow T^{00}|_{\text{Pre-Planckian}} - T^{\alpha\beta}|_{\text{Planckian}} &\propto t_{LL}^{\alpha\beta}|_{\text{Planckian}} - t_{LL}^{00}|_{\text{Pre-Planckian}} \end{aligned}$$

### 3. Conclusion, implication of causal discontinuity as implied by Eq. (16)

Note that having the right hand side of the 2<sup>nd</sup> line of Eq. (16) going to zero is implying that there is an invariance as to the gravitational field pseudo tensor, which may have implications as to the entropy- information transfer from Pre Planckian to Planckian space-time , however we have to consider the Pre Planckian to Planckian physics of the more traditional stress energy tensor.

What we will say, is that by [13] and [14] if we have that the 2<sup>nd</sup> line of Eq. (16) equals zero, with this reflecting the  $T^{00}|_{\text{Pre-Planckian}}$  as a time line fluid momentum density (assuming  $c=1$  in units) and  $T^{\alpha\beta}|_{\text{Planckian}}$  being a general stress energy tensor when we have Planckian, as opposed to Pre Planckian conditions, that we say that

$$T^{\alpha\beta}|_{\text{Planckian}} \neq T^{00}|_{\text{Planckian}} \quad (17)$$

Dispersal of the terms alpha and beta, into the non time components of the stress energy tensor would be saying also that

$$T^{00}\Big|_{Pre-Planckian} \neq T^{00}\Big|_{Planckian} \quad (18)$$

Note that in [14] we have that

Quote

These gapless Goldstone modes are the quantum carriers of information and entropy. Analyzing their effective theory, we observe the information-processing properties strikingly similar to the ones predicted by the black hole portrait. The energy cost per qubit of information-storage vanishes in the large- $N$  limit and the total information-storage capacity increases with  $N$  either exponentially or as a power law.

End of quote

Our idea is that the  $N$  limit, of information entropy, is akin to graviton counting, using the information given in section 2, as given by Ng [6] and that if or not we have gravitons in a counting mode will be strongly affected by a possible causal discontinuity as given by Eq. (18). If or not we have a causal continuity or discontinuity will also be affected by the simple minded rendition of inflaton physics, whereas

$$M = \frac{\hbar}{a_{\min}^2 \phi_{\min} \cdot \delta t} \sim S(\text{entropy}) \times m(\text{graviton})$$

$$\Leftrightarrow \phi_{\min} \sim \frac{\hbar}{a_{\min}^2 (S(\text{entropy}) \times m(\text{graviton})) \cdot \delta t} \sim \frac{\hbar}{a_{\min}^2 (N_{\text{count}} \times m(\text{graviton})) \cdot \delta t} \quad (19)$$

$$\&0 < N_{\text{count}} \sim \text{Initial} - \text{entropy}$$

If  $N$  being a graviton count is sufficiently large, and the initial inflation can be parameterized, we can understand if or not Eq. (18) is a causal structure discontinuity.

Furthermore, understanding Eq. (16) to Eq. (19) more fully may allow us to choose between the different models given in Table 1, among other things looking at if the following is true, namely

‘Note that usual Randall Sundrum brane theory has a production rate [15,16] of

$$\Gamma \sim T^6 / M_{\text{Planck}}^2 \quad (20)$$

as the number of Kaluza Klein gravitons per unit time per unit volume Note that this production rate is for a formula assuming mass for which  $T > M_X$ , and that we are assuming that the

temperature  $T \sim T_*$ . Furthermore, we also are looking at a de facto total production rate of KK gravitons of the form [15]

$$\frac{dn}{dt} \sim \frac{T^6}{M_{Planck}^2} \cdot (T \cdot R)^d \sim T^4 \cdot \left( \frac{T}{M_X} \right)^{2+d} \quad (21)$$

Where R is the assumed higher dimension 'size' and , d is the number of dimensions above 4, and typically we obtain  $T \gg 1/R$ .

This with additional work may allow us to distinguish between the GW and gravity models as given in reference [17] by Corda as well as giving more definition to the issues brought up the the LIGO analysis of gravitational waves, as given in [18] and [19]

## ACKNOWLEDGEMENTS

This work is supported in part by National Nature Science Foundation of China grant No. 11375279

## BIBLIOGRAPHY

- [1] Poisson, E, Will, C. Gravity, *Newtonian, Post Newtonian, and Relativistic*, Cambridge University Press, Cambridge, UK. 2014
- [2] Li, F., Yang, N., et. al., Phys.Rev.D78:024041,2008, <http://arxiv.org/abs/0807.0885>
- [3] Zakharov, Valentin I. (1970). "Linearized gravitation theory and the graviton mass". *JETP Lett.* **12**: 312. [Bibcode:1970JETPL..12..312Z](https://arxiv.org/abs/1970JETPL..12..312Z)
- [4] Beckwith, A. (2016) Gedanken Experiment for Refining the Unruh Metric Tensor Uncertainty Principle via Schwarzschild Geometry and Planckian Space-Time with Initial Nonzero Entropy and Applying the Riemannian-Penrose Inequality and Initial Kinetic Energy for a Lower Bound to Graviton Mass (Massive Gravity). *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 106-124. doi: [10.4236/jhepgc.2016.21012](https://arxiv.org/abs/10.4236/jhepgc.2016.21012).
- [5] Beckwith, A. (2016) Open Question: Could a Causal Discontinuity Explain Fluctuations in the CMBR Radiation Spectrum?. *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 186-208. doi:[10.4236/jhepgc.2016.22018](https://arxiv.org/abs/10.4236/jhepgc.2016.22018).
- [6] Ng, Y, "Article: Spacetime Foam: From Entropy and Holography to Infinite Statistics and Nonlocality" Entropy 2008, 10(4), 441-461; DOI: 10.3390/e10040441

- [7] Kolb, E., and Turner, S. "The Early Universe", Westview Press, Chicago, USA, 1994
- [8] Lloyd , Seth "Computational capacity of the universe", Phys. Rev. Lett. 88, 237901 (2002)
- [9] D. Gorbunov, V. Rubakov, *Introduction to the Theory of the Early Universe, Cosmological Perturbations and Inflationary Theory*, World Scientific Publishing Pte. Ltd, Singapore, Republic of Singapore, 2011
- [10] W. G. Unruh; "Why study quantum theory?", *Canadian Journal of Physics*, 1986, Vol. 64, No. 2 : pp. 128-130; (doi: 10.1139/p86-019)
- [11] W. G. Unruh; "Erratum: Why study quantum gravity?", Can. J. Phys. 64, 128 (1986)
- [12] Beckwith, A. (2016) Gedanken Experiment for Delineating the Regime for the Start of Quantum Effects, and Their End, Using Turok's Perfect Bounce Criteria and Radii of a Bounce Maintaining Quantum Effects, as Delineated by Haggard and Rovelli. *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 287-292. doi:10.4236/jhepgc.2016.23024.
- [13] Beckwith, A. (2016) Open Question: Could a Causal Discontinuity Explain Fluctuations in the CMBR Radiation Spectrum?. *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 186-208. doi:10.4236/jhepgc.2016.22018.
- [14] Dvali G. ,Franca A., Gomez, C., and Wintergerst, N., "Nambu-Goldstone Effective Theory of Information at Quantum Criticality", arXiv:1507.02948v1 [hep-th] 10 Jul 2015 ; Phys. Rev. D 92, 125002 (2015)
- [15] V. A. Rubakov, "Large and infinite extra dimensions"; <http://arxiv.org/pdf/hep-ph/0104152.pdf>
- [16] Valery Rubakov *Classical Theory of Gauge Fields*, 456 pages, Princeton University Press, 2002, Princeton, New Jersey, USA; Translated by Stephen S. Wilson
- [17] Corda, Christian **Interferometric detection of gravitational waves: the definitive test for General Relativity** - Corda, Christian Int.J.Mod.Phys. D18 (2009) 2275-2282 arXiv:0905.2502 [gr-qc]
- [18] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration)  
Phys. Rev. Lett. **116**, 061102 – Published 11 February 2016; <https://physics.aps.org/featured-article-pdf/10.1103/PhysRevLett.116.061102>
- [19] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration); Phys. Rev. Lett. 116, 241103