

# VECTOR LORENTZ TRANSFORMATIONS

A. Blato

Creative Commons Attribution 3.0 License

(2016) Buenos Aires

Argentina

This article presents the vector Lorentz transformations of time, space, velocity and acceleration.

## Introduction

If we consider two inertial reference frames (  $S$  and  $S'$  ) whose origins coincide at time zero ( in both frames ) then the time (  $t'$  ), the position (  $\mathbf{r}'$  ), the velocity (  $\mathbf{v}'$  ) and the acceleration (  $\mathbf{a}'$  ) of a (massive or non-massive) particle relative to the inertial reference frame  $S'$  are given by:

$$t' = \gamma \left( t - \frac{\mathbf{r} \cdot \mathbf{V}}{c^2} \right)$$

$$\mathbf{r}' = \left[ \mathbf{r} + \frac{\gamma^2}{\gamma + 1} \frac{(\mathbf{r} \cdot \mathbf{V}) \mathbf{V}}{c^2} - \gamma \mathbf{V} t \right]$$

$$\mathbf{v}' = \left[ \mathbf{v} + \frac{\gamma^2}{\gamma + 1} \frac{(\mathbf{v} \cdot \mathbf{V}) \mathbf{V}}{c^2} - \gamma \mathbf{V} \right] \frac{1}{\gamma (1 - \frac{\mathbf{v} \cdot \mathbf{V}}{c^2})}$$

$$\mathbf{a}' = \left[ \mathbf{a} - \frac{\gamma}{\gamma + 1} \frac{(\mathbf{a} \cdot \mathbf{V}) \mathbf{V}}{c^2} + \frac{(\mathbf{a} \times \mathbf{v}) \times \mathbf{V}}{c^2} \right] \frac{1}{\gamma^2 (1 - \frac{\mathbf{v} \cdot \mathbf{V}}{c^2})^3}$$

where (  $t$ ,  $\mathbf{r}$ ,  $\mathbf{v}$ ,  $\mathbf{a}$  ) are the time, the position, the velocity and the acceleration of the particle relative to the inertial reference frame  $S$ , (  $\mathbf{V}$  ) is the velocity of the inertial reference frame  $S'$  relative to the inertial reference frame  $S$  and (  $c$  ) is the speed of light in vacuum. (  $\mathbf{V}$  ) is a constant.  $\gamma = (1 - \mathbf{V} \cdot \mathbf{V}/c^2)^{-1/2}$

- $\mathbf{v}' = \frac{d\mathbf{r}'}{dt'} = \frac{d\mathbf{r}'}{dt'} \frac{dt}{dt} = \frac{d\mathbf{r}'}{dt} \frac{dt}{dt'} = \left( \frac{d\mathbf{r}'}{dt} \right) \frac{1}{\left( \frac{dt'}{dt} \right)}$
- $\mathbf{a}' = \frac{d\mathbf{v}'}{dt'} = \frac{d\mathbf{v}'}{dt'} \frac{dt}{dt} = \frac{d\mathbf{v}'}{dt} \frac{dt}{dt'} = \left( \frac{d\mathbf{v}'}{dt} \right) \frac{1}{\left( \frac{dt'}{dt} \right)}$
- $dt' = \gamma \left( dt - \frac{d\mathbf{r} \cdot \mathbf{V}}{c^2} \right)$
- $\left( \frac{dt'}{dt} \right) = \gamma \left( 1 - \frac{\mathbf{v} \cdot \mathbf{V}}{c^2} \right)$
- $d\mathbf{r}' = \left[ d\mathbf{r} + \frac{\gamma^2}{\gamma+1} \frac{(d\mathbf{r} \cdot \mathbf{V}) \mathbf{V}}{c^2} - \gamma \mathbf{V} dt \right]$
- $\left( \frac{d\mathbf{r}'}{dt} \right) = \left[ \mathbf{v} + \frac{\gamma^2}{\gamma+1} \frac{(\mathbf{v} \cdot \mathbf{V}) \mathbf{V}}{c^2} - \gamma \mathbf{V} \right]$
- $d\mathbf{v}' = \left[ d\mathbf{m} \cdot n - \mathbf{m} \cdot dn \right] \frac{1}{n^2}$
- $\mathbf{m} = \left[ \mathbf{v} + \frac{\gamma^2}{\gamma+1} \frac{(\mathbf{v} \cdot \mathbf{V}) \mathbf{V}}{c^2} - \gamma \mathbf{V} \right]$
- $d\mathbf{m} = \left[ d\mathbf{v} + \frac{\gamma^2}{\gamma+1} \frac{(d\mathbf{v} \cdot \mathbf{V}) \mathbf{V}}{c^2} \right]$
- $n = \left[ \gamma \left( 1 - \frac{\mathbf{v} \cdot \mathbf{V}}{c^2} \right) \right]$
- $dn = \left[ -\gamma \frac{d\mathbf{v} \cdot \mathbf{V}}{c^2} \right]$
- $\left( \frac{d\mathbf{v}'}{dt} \right) = \left[ \mathbf{a} - \frac{\gamma}{\gamma+1} \frac{(\mathbf{a} \cdot \mathbf{V}) \mathbf{V}}{c^2} + \frac{(\mathbf{a} \times \mathbf{v}) \times \mathbf{V}}{c^2} \right] \frac{1}{\gamma \left( 1 - \frac{\mathbf{v} \cdot \mathbf{V}}{c^2} \right)^2}$

## Bibliography

[https://it.wikipedia.org/wiki/Trasformazione\\_di\\_Lorentz](https://it.wikipedia.org/wiki/Trasformazione_di_Lorentz)

[https://en.wikipedia.org/wiki/Lorentz\\_transformation](https://en.wikipedia.org/wiki/Lorentz_transformation)

<https://arxiv.org/abs/physics/0507099>

<https://arxiv.org/abs/physics/0702191>

[https://archive.org/details/blato\\_links](https://archive.org/details/blato_links)

[https://archive.org/details/@a\\_blato](https://archive.org/details/@a_blato)

## Appendix I

In the previous bibliography, if the velocity of the inertial reference frame  $S'$  relative to the inertial reference frame  $S$  is non-zero ( $\mathbf{V} \neq 0$ ) then we have:

$$\frac{\gamma - 1}{\mathbf{V}^2} = \frac{\gamma^2}{\gamma + 1} \frac{1}{c^2} \quad (\mathbf{V}^2 = \mathbf{V} \cdot \mathbf{V})$$

where ( $c$ ) is the speed of light in vacuum.  $\gamma = (1 - \mathbf{V} \cdot \mathbf{V}/c^2)^{-1/2}$

## Appendix II

Quotient rule:  $\mathbf{a} = \frac{\mathbf{m}}{n} \quad \rightarrow \quad d\mathbf{a} = \left[ d\mathbf{m} \cdot n - \mathbf{m} \cdot dn \right] \frac{1}{n^2}$

Triple product expansion:  $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b})$

Anticommutativity:  $(\mathbf{a} \times \mathbf{b}) = -(\mathbf{b} \times \mathbf{a})$