

# The Theory of Quantum Gravity and Calculation of Cosmological Constant

M. Christolubova<sup>a</sup>, T. Christolubov<sup>b</sup>

*The Massachusetts Institute of Technology*

## Abstract

To construct quantum gravity we formulate quantum electrodynamics in equivalent form with possibility to generalize, we calculate the cosmological constant assuming that the quantum state is a function of time and radius of universe.

*Key words:* Quantum gravity, cosmological constant, Friedman.

PACS: 11.30Cp, 12.20.-m, 12.60.-i

## 1. Introduction

The new and equivalent formulation of quantum electrodynamics is given in terms of space of states.

$$\Psi = \Psi(t, x_1, \dots, x_n, \{A^\gamma(x)\}) \quad (1)$$

where  $A^\gamma$  is the electromagnetic field. To construct quantum gravity we add metric as the new argument of state. The quantum gravity state is

$$\Psi = \Psi(t, x_1, \dots, x_n, \{A^\gamma(x)\}, \{g_{\mu\nu}(x)\}) \quad (2)$$

For example, the mean value of metric at point  $x_0$  at time  $t$

$$\langle g_{\mu\nu}(t_0, x_0) \rangle = \int Dx_1 \dots Dx_n D\{A^\gamma(x)\} D\{g_{\mu\nu}(x)\} \Psi^*(t) g_{\mu\nu}(x_0) \Psi(t) \quad (3)$$

The quantum gravity must be completely field theory, we substitute particle coordinates with the mass fields.

$$\Psi = \Psi(t, \{\Phi_1(x)\}, \dots, \{\Phi_n(x)\}, \{A^\gamma(x)\}, \{g_{\mu\nu}(x)\}) \quad (4)$$

To find the equation of motion we use the approach

Action  $\rightarrow$  Lagrangian  $\rightarrow$  Hamiltonian  $\rightarrow$  Quantization.

The lagrangian density participates in the integral

$$S = \int dt d^3x \mathcal{L} \quad (5)$$

The momentum density

$$p^\mu = \frac{\partial \mathcal{L}}{\partial(\dot{\varphi}(x))} \quad (6)$$

where  $\varphi$  are the all fields.

$$\varphi = (g^{\alpha\beta}, A^\mu, \phi_1, \dots, \phi_n), \quad (7)$$

where  $\phi_1, \dots, \phi_n$  are the mass fields.

The Hamiltonian density

$$\mathcal{H} = p^\mu \partial_\mu \varphi - \mathcal{L} \quad (8)$$

The Hamiltonian

$$H = \int \mathcal{H} d^3x \quad (9)$$

To perform quantization we substitute the fields and the momentum of fields with the operators. If

$$\mathcal{H} = \mathcal{H}(t, x, \varphi(x), p^\mu(x), \frac{\partial^2 \varphi}{\partial x_\mu \partial x_\nu}) \quad (10)$$

We introduce the operator of the field value at the point  $x_0$

$$\hat{\varphi}(x_0) : \Psi \longrightarrow \varphi(x_0)\Psi \quad (11)$$

The momentum operator of the field value at the point  $x_0$

$$\hat{p}_\varphi(x_0) : \Psi \longrightarrow -i\hbar \lim_{q \rightarrow 0} \frac{\Psi(\{\phi(x) + q\delta(x - x_0)\}) - \Psi(\{\phi(x)\})}{q} \quad (12)$$

The momentum and coordinate operators satisfy

$$\frac{i}{\hbar} [\hat{p}_{\phi(x)}, \hat{\varphi}(x_0)] = \delta(x - x_0) \quad (13)$$

Then the "acceleration of field" can found from

$$\frac{i}{\hbar} [\hat{H}, \hat{\varphi}(x_0)] = \ddot{\varphi}(x_0) \quad (14)$$

## 1. Relativistic invariance

To achieve the relativistic invariance we represent our manifold as set of slices, with invariant parameter w-number of slices, and the invariant points M on each slices.

## 2. The calculation of cosmological constant

To calculate the cosmological constant we perform the quantization of Friedman model. Suppose that the state is the function of time and the universe curvature radius.

$$\Psi = \Psi(t, a) \quad (15)$$

To calculate the quantum Hamiltonian we use our scheme:

$$\text{Action} \rightarrow \text{Lagrangian} \rightarrow \text{Hamiltonian} \rightarrow \text{Quantization.}$$

As well known, the amount of the visible matter in universe is less than 5 %. We assume that the universe is massless and the Einstein-Hilbert action given by

$$S = \frac{c^3}{16\pi G} \int d\Omega R = \frac{c^3}{16\pi G} \int dt d\vec{x} \sqrt{-g} R \quad (16)$$

$$L = \int d\vec{x} \frac{c^3}{16\pi G} \sqrt{-g} R \quad (17)$$

If we substitute into the action the Friedman metric (open case)

$$d^2 s = c^2 dt^2 - a^2(t)(d\chi^2 + \sinh^2 \chi^2 (d\Theta^2 + \sin^2 \Theta d\phi^2)) \quad (18)$$

If we introduce the new variable  $\eta$  which satisfies the differential relation  $ad\eta = cdt$

$$d^2 s = a^2(\eta)(d\eta^2 - d\chi^2 - \sinh^2 \chi (d\Theta^2 + \sin^2 \Theta d\phi^2)) \quad (19)$$

Then the calculation gives

$$R = \frac{6(a - a''(\eta))}{a^3}, g = -a^8 \sinh^4 \chi \sin^2 \Theta \quad (20)$$

Taking the integral of the angles, we have:

$$S = \varpi \int d\eta a(a - a''(\eta)) = \varpi \int d\eta (a^2 + a'^2) = \int d\eta L, \quad (21)$$

where the Lagrangian is

$$L = a'^2 + a^2 = E - U \quad (22)$$

is the Lagrangian of the particle in the accelerating potential

$$U = -a^2 \quad (23)$$

Suppose that quantum mean values obey the Eirenfest theorems which are our case the Euler-Lagrange equations, which give

$$\langle a'' \rangle - \langle a \rangle = 0 \quad (24)$$

Their general solution is

$$a = a_1 e^\eta + a_2 e^{-\eta} \quad (25)$$

I.Case 1(the initial radius of Universe is no-zero and positive: $a_1 > 0, a_2 > 0$ ) Making a shift of  $\eta$ , we can make

$$a_1 = a_2 \quad (26)$$

so that

$$a = a_1 e^\eta + a_1 e^{-\eta} = a_0 \cosh \eta, \quad a' = a_0 \sinh \eta \quad (27)$$

The neighboring geodesics equations

$$\frac{D^2 \Delta x^\gamma}{D\tau^2} = R_{\alpha\beta\gamma}^\gamma \nu^\alpha \nu^\beta \Delta x^\gamma \quad (28)$$

In the case of zero spatial velocities we have

$$\frac{D^2 \Delta x^\gamma}{D\tau^2} = R_{00\gamma}^\gamma (\nu^0)^2 \Delta x^\gamma \quad (29)$$

The sign of  $R_{00\gamma}^\gamma$  shows if the universe accelerates

$$R_{00\eta}^\eta = -\left(\frac{a'}{a}\right)^2 + \frac{a''}{a} = -\left(\frac{a'}{a}\right)^2 + 1 = -\tanh^2 \eta + 1 > 0 \quad (30)$$

We observe the positive acceleration of the universe, which has not ever been given by any theory.

I.Case 2(the initial radius of Universe is zero: $a_1 > 0, a_2 < 0$ ) Making a shift of  $\eta$ , we can make

$$a_1 = a_2 \quad (31)$$

so that

$$a = a_1 e^\eta - a_1 e^{-\eta} = a_0 \sinh \eta, \quad a' = a_0 \cosh \eta \quad (32)$$

The neighboring geodesics equations

$$\frac{D^2 \Delta x^\gamma}{D\tau^2} = R_{\alpha\beta\gamma}^\gamma \nu^\alpha \nu^\beta \Delta x^\gamma \quad (33)$$

In the case of zero spatial velocities we have

$$\frac{D^2 \Delta x^\gamma}{D\tau^2} = R_{00\gamma}^\gamma (\nu^0)^2 \Delta x^\gamma \quad (34)$$

The sign of  $R_{00\gamma}^\gamma$  shows if the universe accelerates

$$R_{00\eta}^\eta = -\left(\frac{a'}{a}\right)^2 + \frac{a''}{a} = -\left(\frac{a'}{a}\right)^2 + 1 = -\csc^2 \eta + 1 < 0 \quad (35)$$

We see that if at the starting time the universe was at one point then we would not see the acceleration of Universe. We can see the accelerating Universe only if the initial radius of Universe is not zero(see case 1).This is the argument against Big Bang.

### 3. The Cosmological Constant

In Case 1 of the accelerating Universe

$$a = a_0 \cosh \eta \quad (36)$$

Remembering that  $ad\eta = cdt$  and integrating we obtain

$$ct = a_0 \sinh \eta \quad (37)$$

The Hubble constant

$$H = \frac{da/dt}{a} = \frac{cda/d\eta}{a^2} = \frac{ca'}{a^2} \quad (38)$$

Substituting there again (31) and using (32), we have:

$$H = \frac{ca_0 \sinh \eta}{a_0^2 \cosh^2 \eta} = \frac{\sinh^2 \eta}{t \cosh^2 \eta} = \frac{\tanh^2 \eta}{t} \quad (39)$$

From where we have:

$$\eta = \operatorname{arctanh}(Ht)^{1/2} \quad (40)$$

If we suppose  $Ht=2/3$  we have

$$\eta = \operatorname{arctanh}(2/3)^{1/2} \approx 1.15 \quad (41)$$

Using neighboring geodesics equation (28) and (29) we see that the sign

$$\frac{D^2(\Delta x^\gamma)}{D\tau^2} / \Delta x^\gamma = R^\gamma_{00\gamma} (v^0)^2 = \frac{c^2}{a^2} \left( \left( -\frac{at'}{a} \right)^2 + \frac{a''}{a} \right) = -H^2 + \frac{c^2}{a^2} \quad (42)$$

Using equation for Hubble constant, we obtain

$$\frac{D^2(\Delta x^\gamma)}{D\tau^2} / \Delta x^\gamma = -\left( \frac{ca'}{a^2} \right)^2 + \frac{c^2}{a^2} = \frac{c^2(-a'^2 + a^2)}{a^4} = \frac{c^2 a_0^2}{a^4} = \frac{c^2}{a_0^2 \cosh^4 \eta} \quad (43)$$

Using (32), we receive

$$\frac{D^2(\Delta x^\gamma)}{D\tau^2} / \Delta x^\gamma = \frac{(\sinh \eta)^2}{t^2 \cosh^4 \eta} = \frac{(\tanh \eta)^2}{t^2 \cosh^2 \eta} = \frac{(Ht)^2}{t^2 \cosh^2 \eta} = \frac{H^2}{\cosh^2 \eta} \quad (44)$$

Substituting the values of  $\eta$  and  $H$ , we have according to the theory

$$\frac{D^2(\Delta x^\gamma)}{D\tau^2} / \Delta x^\gamma = \frac{H^2}{3} \approx 1.33 * 10^{-36} \frac{1}{s^2} \quad (45)$$

Experiment gives the following value of Universe Expansion acceleration

$$\frac{D^2(\Delta x^\gamma)}{D\tau^2} / \Delta x^\gamma = 2 \frac{2\pi G \rho_{cr}}{3} \approx 2 * 10^{-36} \frac{1}{s^2} \quad (46)$$

## Conclusion

The construction of quantum gravity is possible if we stand on very good land of quantum states, this path now has shown the way to calculate the cosmological constant

## References

1. *Moyotl A. , Novales-Sánchez H., Toscano J. J., Tutut E.S.* // Int. J. Mod. Phys. A. 2014. **29**. P. 1450039.
2. *Langacker P.* The Standard Model and Beyond. Boca Raton, 2010.
3. *Olive K.A.* et al. (Particle Data Group) // Chin. Phys. C. 2014. **38**. P. 090001.
4. *Kostelecký V.A., Samuel S.* // Phys. Rev. D. 1989. **39**. P. 683.
5. *Carroll S. M., Harvey J.A., Kostelecký V.A., Lane C.D., Okamoto T.* // Phys. Rev. Lett. 2001. **87**. P. 141601.
6. *Weinberg S.* The Quantum Theory of Fields. Vol. I. Foundations. Cambridge, 1995.
7. Православный Христианский священник призывает Путина и Медведева на исповедь
8. *Colladay D., Kostelecký V.A.* // Phys. Rev. D. 1998. **58**. P. 116002.
9. *Kostelecký V.A. (Editor).* Proceedings of the Fifth Meeting on CPT and Lorentz Symmetry. Singapore, 2011.
10. *Jackiw R., Kostelecký V.A.* // Phys. Rev. Lett. 1999. **82**. P. 3572.
11. *Carroll S, Field G., Jackiw R.* // Phys. Rev. D. 1990. **41**. P. 1231.
12. *Zhukovsky V.Ch., Lobanov A.E., Murchikova E.M.* // Phys. Rev. D. 2006. **73**. P. 065016.
13. *Жуковский В. Ч., Лобанов А.Е., Мурчи́кова Е.М.* // Ядерная физика. 2007. **70**. С. 1289.
14. *Frolov I.E., Zhukovsky V.Ch.* // J. Phys. A. 2007. **40**. P. 10625.
15. *Heisenberg W., Euler H.* // Z. Phys. 1936. **98**. S. 714.
16. *Тернов И.М., Жуковский В. Ч., Борисов А.В.* Квантовые процессы в сильном внешнем поле. М., 1989.

17. *Берестецкий В.Б., Лифшиц Е.М., Питаевский Л.П.* Квантовая электродинамика. 4-е изд., испр. М., 2002.
18. *Borodulin V.I., Rogalyov R.N., Slabospitsky S.R.* CORE (COmpendium of RElations). Version 2.1. ИИЕР-95-90. Protvino, 1995; arXiv:hep-ph/9507456.
19. *Рунус В.И.* // Проблемы квантовой электродинамики интенсивного поля (Тр. ФИАН. Т. 168). М., 1986. — С. 52.
20. *Тернов И.М., Багров В.Г., Бордовицын В.А., Дорофеев О.Ф.* // ЖЭТФ. 1968. **55**. С. 2273.
21. *Tsai W.-y.* // Phys. Rev. D. 1973. **8**. P. 3446.
22. *Chyi T.-K., Hwang C.-W., Kao W. F., Lin G.-L., Ng K.-W., Tseng J.-J.* // Phys. Rev. D. 2000. **62**. P. 105014.
23. *Соколов А.А., Тернов И.М.* Релятивистский электрон. М., 1982.
24. *Schwinger J.* // Phys. Rev. 1948. **73**. P. 416; 1949. **76**. P. 790.
25. *Kostecký V.A., Russell N.* // Rev. Mod. Phys. 2011. **83**. P. 11; arXiv:0801.0287v9 (26 Feb 2016).
26. *Colladay D., Kostecký V.A.* // Phys. Lett. B. 2001. **511**. P. 209.

