

# Grand Unification from a Classical Interpretation of “Spin” Angular Momentum

Brent Lee Jarvis  
Embry Riddle Aeronautical University  
[JarvisB@my.erau.edu](mailto:JarvisB@my.erau.edu)

**Abstract:** It is shown from the laws of conservation of energy and angular momentum that the gravitational energy of a bound state particle is dependent upon its geometric mean angular frequency. A classical interpretation for the intrinsic “spin” angular momentum of a particle is submitted that can be tested macroscopically with the Rarita–Schwinger equation for the orbit of Mercury. Librations in the relative obliquity of planetary bodies are shown to be analogous to the Larmor precession of atomic particles. Schrodinger's concept of a bound state electron's zitterbewegung is hypothesized to be the librations in the electron's relative obliquity.

**Keywords:** Gravitational Energy, Spin, Obliquity, Larmor Precession, Zitterbewegung

## INTRODUCTION

It is well known that the angular momentum  $L$  of a bound state particle in an eccentric orbit is equivalent at the extreme points of its orbit:

$$(1) \quad L = m\mathbf{v}_{\max} \times \mathbf{r}_{\min} = m\mathbf{v}_{\min} \times \mathbf{r}_{\max},$$

where  $m$  is a particle's mass,  $v$  is its tangential velocity, and  $r$  is its orbital radius. We also know from the law of conservation of energy that

$$(2) \quad 0 = \left(\frac{1}{2}m\mathbf{v}_{\max}^2 - \frac{GMm}{r_{\min}}\right) - \left(\frac{1}{2}m\mathbf{v}_{\min}^2 - \frac{GMm}{r_{\max}}\right),$$

where  $G$  is Newton's gravitational constant and  $M$  is the mass of a primary body.

## A PARTICLE'S NON-RELATIVISTIC GRAVITATIONAL ENERGY

Rearranging Eq. 2 and removing the common factor  $m$ ,

$$(3) \quad \frac{1}{2}(\mathbf{v}_{\max}^2 - \mathbf{v}_{\min}^2) = GM \left( \frac{1}{r_{\min}} - \frac{1}{r_{\max}} \right).$$

Substituting the values of  $v_{\min}$  and  $v_{\max}$  from Eq. (1) in Eq. (3),

$$(4) \quad \frac{L^2}{2m^2} \left( \frac{1}{r_{\min}} + \frac{1}{r_{\max}} \right) \left( \frac{1}{r_{\min}} - \frac{1}{r_{\max}} \right) = GM \left( \frac{1}{r_{\min}} - \frac{1}{r_{\max}} \right),$$

and upon rearrangement of the radii,

$$(5) \frac{L^2}{2m^2} = \frac{GM}{\left(\frac{1}{r_{\min}} + \frac{1}{r_{\max}}\right)}.$$

From orbital mechanics we know the semi-major axis  $a$  and semi-minor axis  $b$  of a particle's orbit are equivalent to its arithmetic and geometric mean distances from the focus respectively. The sum of the inverse extreme radii in Eq. (5) is therefore,

$$(6) \left(\frac{1}{r_{\min}} + \frac{1}{r_{\max}}\right) = \frac{r_{\min} + r_{\max}}{r_{\min} r_{\max}} = \frac{2a}{b^2}.$$

Combining Eqs. (5 & 6),

$$(7) \frac{L^2}{2m^2} = \frac{GMb^2}{2a},$$

from which a particle's angular momentum  $L$  is,

$$(8) L = mb\sqrt{\frac{GM}{a}}.$$

Since  $L$  is a conserved quantity and a particle's geometric mean distance  $b = \sqrt{r_{\min}r_{\max}}$ , a particle's geometric mean tangential velocity  $v_b$  is,


$$(9) v_b = \sqrt{v_{\min} v_{\max}} = \frac{L}{mb} = \sqrt{\frac{GM}{a}}.$$

A particle's non-relativistic gravitational energy  $U(b, t)$  can then be defined from Eqs. (7 & 9) as,

$$(10) U(b, t) = -\frac{GMm}{a} = -\frac{L^2}{mb^2} = -mv_b^2 = -L \cdot \frac{v_b}{b} = -L \cdot \omega_b,$$


where  $\omega_b$  is a particle's geometric mean angular frequency.

## A CLASSICAL INTERPRETATION OF “SPIN”



We can see from the equation of time graph to the left that the frequency of Earth's obliquity (mauve dashed curve) is roughly twice the frequency of its eccentricity (blue dash-dot curve). The superposition of the orthogonal temporal waves (red curve) can be graphed parametrically as a Lissajous figure that can be observed from the Earth's surface as the solar analemma:

$$(11) x(t) = A \sin(st + \delta), y(t) = B \sin t =$$



where  $A$  and  $B$  are the amplitudes of the eccentric and oblique temporal waves respectively,  $\delta$  is the observational angle of the Lissajous figure, and the “spin” ratio  $s = f_E : f_O \approx 1:2$  (for Earth's orbit), where  $f_E$  and  $f_O$  are the frequencies of the Earth's eccentricity and obliquity respectively.

The visual appearance of a Lissajous figure is dependent upon the “height-to-width” ratio  $A : B$  and the spin ratio  $s$  (a circle resulting when  $A = B$ ,  $s = 1:1$ , and  $\delta = \pi:2$ ). We can see from the visual appearance of the solar analemmas of other planets in our system,




that a planet's spin ratio  $s$  can vary between  $\approx 1:1$  to  $1:2$ , analogous to the spin of bosons and fermions in electrodynamics. It can also be deduced from Eq. (11) that  $s = 2:1$ ,  $\delta = \pi:2$  results in,

$$(12) \quad (A \cos(2t), \sin t) = (A(1 - 2 \sin^2 t), \sin t) = (A - 2A \sin^2 t, \sin t),$$

which is a horizontally offset version of the parametric equation for a parabola. This could explain why particles with  $s \geq 2$  have not been observed in atomic nuclei. The planet Mercury, with  $s \approx 3:2$ , would be analogous to a composite gravitino; theoretically governed by the Rarita–Schwinger equation<sup>[1]</sup>,

$$(13) \quad (\epsilon^{\mu\nu\rho\sigma} \gamma_\sigma \gamma_\rho \partial_\mu - m \frac{i^2}{2} [\gamma^\mu, \gamma^\nu]) \psi_\nu = 0,$$

where  $\epsilon^{\mu\nu\rho\sigma}$  is the alternating symbol,  $\gamma_\sigma$  and  $\gamma_\rho$  are Dirac matrices,  $m$  is the gravitino's mass, and  $\psi_\nu$  is a vector-valued spinor that has additional components compared to the Dirac equation spinor<sup>[2]</sup>. With the values in Eq. (11) set to  $A = B$ ,  $s = 3:2$ ,  $\delta = \pi:2$ , the Lissajous figure for a gravitino would be:



## THE ORIGIN OF A PLANET'S OBLIQUE ANGULAR FREQUENCY

It is proposed that a planet's oblique angular frequency  $\tilde{\omega} = 2\pi f_O$  (Larmor precession)<sup>[3]</sup> is caused by the torque  $\tau$  induced on its magnetic moment  $m_P$  by the total magnetic field  $\mathbf{B}$  of the system<sup>[4,5,6,7]</sup>,

$$(14) \quad \tau = m_P \times \sum_{i=1}^n \mathbf{B}_i = \gamma \mathbf{J} \times \sum_{i=1}^n \mathbf{B}_i = \mathbf{J} \times \tilde{\omega} = 1 + \frac{M_{orb}}{(M_{spin} + M_{orb})} \mathbf{J} \times \sum_{i=1}^n \mathbf{B}_i,$$

where the last term describes the Einstein–de Haas<sup>[4]</sup> and Barnett<sup>[5]</sup> effects,  $\gamma$  is a planet's gyromagnetic ratio,  $\mathbf{J}$  is its total (gyromagnetic) angular momentum<sup>[3]</sup>,  $M_{orb}$  is its orbital magnetization<sup>[6,7]</sup> and  $M_{spin}$  is its spin magnetization<sup>[5]</sup>. Isolating the oblique angular frequency  $\tilde{\omega}$ <sup>[3]</sup> to one side in Eq. (14) yields,

$$(15) \quad \tilde{\omega} = \frac{|\boldsymbol{\tau}|}{\mathbf{J} \cdot \sin \theta} = \gamma \sum_{i=1}^n \mathbf{B}_i = \left( 1 + \frac{M_{\text{orb}}}{(M_{\text{spin}} + M_{\text{orb}})} \right) \sum_{i=1}^n \frac{\mu_0 \mathbf{m}_i}{4\pi r_i^3} \sqrt{1 + 3 \cos^2 \theta_i},$$

where  $\mu_0$  is the magnetic constant and  $\theta_i$  is the azimuthal angle from the polar axis of each of the magnetic dipole moments  $\mathbf{m}_i$  within the system.

The oblique angular frequency  $\tilde{\omega}$  (Larmor precession) for planets with  $s \approx 1:2$  (analogous to composite fermions) could theoretically be described with the Bargmann–Michel–Telegdi (BMT) equation<sup>[8]</sup>,

$$(16) \quad \tilde{\omega} = \frac{|\boldsymbol{\tau}|}{\mathbf{J} \cdot \sin \theta} = \gamma \sum_{i=1}^n \mathbf{B}_i = 2 \mathbf{m}_p (F^{\alpha\lambda} - u^\alpha u_\sigma F^{\sigma\lambda}) a_\lambda,$$

where  $\mathbf{m}_p$  is the planet's magnetic moment,  $u^\alpha$  is its four-velocity, and  $F^{\alpha\sigma}$  is the “gravitomagnetic” field-strength tensor. Assuming the coefficient 2 in the BMT equation is a particle's inverse spin ratio  $\xi = 2 = 1:s$ , the coefficient would be  $\xi = 1$  for bosons and  $\xi = 2/3$  for gravitinos.

## CONCLUSION

It was discovered by Schrodinger<sup>[9]</sup> from analysis of solutions to the Dirac equation<sup>[2]</sup> that the energy of a bound state electron oscillates with an angular frequency  $\tilde{\omega}_e$ <sup>[9,10]</sup> equivalent to,

$$(17) \quad \tilde{\omega}_e = \frac{2m_e c^2}{\hbar},$$

where  $c$  is the speed of light in a vacuum,  $m_e$  is an electron's mass, and  $\hbar$  is Planck's reduced constant. Schrodinger referred to this as an electron's “zitterbewegung” (wiggling motion). It is hypothesized that an electron's zitterbewegung is the angular frequency of its obliquity  $\tilde{\omega}$ <sup>[3,10]</sup>, and the reason  $\tilde{\omega}$  differs by a factor of 2 from the Planck–Einstein relation is because fermions are  $s = 1:\xi = 1:2$  particles.

Since an electron's angular momentum is quantized in units of  $\hbar$ , and we know from Einstein's special theory of relativity that  $E = mc^2$ , the energy of a bound state electron's orbit could be interpreted from Eqs. (10 & 17) as being its gravitational (potential) energy,

$$(18) \quad E = \frac{\hbar \cdot \tilde{\omega}_e}{2} = \frac{\mathbf{L} \cdot \tilde{\boldsymbol{\omega}}}{\xi} = \mathbf{L} \cdot \omega_b = U(b, t).$$

Since a particle's angular momentum  $\mathbf{L}$  is dependent upon its geometric mean distance  $b$  (Eq. 8), the oscillations in an electron's energy states could be interpreted as oscillations in  $\omega_b$  relative to  $b$  (the gravitational energy  $U(b, t)$  radius is normalized to  $b$  instead of  $\infty$ ).

With this interpretation of Eq. (18), the relative consistency observed in the stellar orbital speeds of spiral galaxies could be explained if the geometric mean angular frequency  $\omega_b$  of the stars increase proportionately with their distance from the galactic nucleus. Geological records<sup>[11]</sup> indicate our Sun oscillates relative to the plane of our galaxy in  $\approx 31 \pm 1$  Myr cycles during its  $\approx 225 - 250$  Myr orbit, which could provide physical evidence for this hypothesis<sup>[12]</sup>.

## REFERENCES

- [1] S. Weinberg, "*The quantum theory of fields*", Vol. 3, Cambridge p. 335
- [2] P.A.M. Dirac, (1982) [1958]. "*Principles of Quantum Mechanics*". International Series of Monographs on Physics (4th ed.). Oxford University Press. p. 255. ISBN 978-0-19-852011-5
- [3] Malcolm H. Levitt, Wiley, (2001). "*Spin Dynamics*".
- [4] A. Einstein, W. J. de Haas, (1915). "*Experimental proof of the existence of Ampère's molecular currents*", Koninklijke Akademie van Wetenschappen te Amsterdam, Proceedings, **18 I**, pp. 696–711
- [5] B. T. Draine, (2003). "§7.3 *Barnett effect*". In Andrew W. Blain; F. Combes; Bruce T. Draine; D. Pfenniger; Yves Revaz. *The Cold Universe*. Springer. p. 276. ISBN 3-540-40838-X.
- [6] R. Resta, (2010). "*Electrical polarization and orbital magnetization: the modern theories*", *Journal of Physics: Condensed Matter*, **22** (12), p. 123201, Bibcode:2010JPCM...22i3201R, doi:10.1088/0953-8984/22/12/123201
- [7] J. Shi; G. Vignale; Q. Niu, Qian, (2007). "*Quantum Theory of Orbital Magnetization and Its Generalization to Interacting Systems*", *Phys. Rev. Lett.*, American Physical Society, **99** (19): 197202, arXiv:0704.3824, Bibcode:2007PhRvL..99s7202S, doi:10.1103/PhysRevLett.99.197202
- [8] V. Bargmann; L. Michel; V. L. Telegdi, (1959). "*Precession of the Polarization of Particles Moving in a Homogeneous Electromagnetic Field*", *Phys. Rev. Lett.* 2, 435
- [9] Huang, (1949). "*On the zitterbewegung of the electron*", *Am. J. Phys.* 47, 797.
- [10] D. Hestenes, (1990). "*The Zitterbewegung Interpretation of Quantum Mechanics*", *Found. Physics.*, Vol. 20, No. 10, 1213–1232. [http://geocalc.clas.asu.edu/pdf/ZBW\\_I\\_QM.pdf](http://geocalc.clas.asu.edu/pdf/ZBW_I_QM.pdf)
- [11] M.R. Rampino, R.B. Stothers, (1984). "*Terrestrial mass extinctions, cometary impacts, and the Sun's motion perpendicular to the galactic plane*", *Nature* 308, 709 – 712.
- [12] B. Jarvis, (2017). "*On the Anomalous Oscillation of Newton's Gravitational Constant*", <https://thewinnower.com/papers/6067-on-the-anomalous-oscillation-of-newton-s-gravitational-constant>