

An Holomorphic Study Of Smarandache Automorphic and Cross Inverse Property Loops^{*†}

Tèmítópé Gbóláhàn Jaíyéolá[‡]

Department of Mathematics,

Obafemi Awolowo University, Ile Ife, Nigeria.

jaiyeolatemitope@yahoo.com, tjayeola@oauife.edu.ng

Abstract

By studying the holomorphic structure of automorphic inverse property quasigroups and loops[AIPQ and (AIPL)] and cross inverse property quasigroups and loops[CIPQ and (CIPL)], it is established that the holomorph of a loop is a Smarandache; AIPL, CIPL, K-loop, Bruck-loop or Kikkawa-loop if and only if its Smarandache automorphism group is trivial and the loop is itself is a Smarandache; AIPL, CIPL, K-loop, Bruck-loop or Kikkawa-loop.

1 Introduction

1.1 Quasigroups And Loops

Let L be a non-empty set. Define a binary operation (\cdot) on L : If $x \cdot y \in L$ for all $x, y \in L$, (L, \cdot) is called a groupoid. If the system of equations ;

$$a \cdot x = b \quad \text{and} \quad y \cdot a = b$$

have unique solutions for x and y respectively, then (L, \cdot) is called a quasigroup. For each $x \in L$, the elements $x^\rho = xJ_\rho, x^\lambda = xJ_\lambda \in L$ such that $xx^\rho = e^\rho$ and $x^\lambda x = e^\lambda$ are called the right, left inverses of x respectively. Now, if there exists a unique element $e \in L$ called the identity element such that for all $x \in L, x \cdot e = e \cdot x = x$, (L, \cdot) is called a loop. To every loop (L, \cdot) with automorphism group $AUM(L, \cdot)$, there corresponds another loop. Let the set $H = (L, \cdot) \times AUM(L, \cdot)$. If we define 'o' on H such that $(\alpha, x) \circ (\beta, y) = (\alpha\beta, x\beta \cdot y)$ for all $(\alpha, x), (\beta, y) \in H$, then $H(L, \cdot) = (H, \circ)$ is a loop as shown in Bruck [7] and is called the Holomorph of (L, \cdot) .

^{*}2000 Mathematics Subject Classification. Primary 20N05 ; Secondary 08A05

[†]**Keywords and Phrases** : Smarandache loop, holomorph of loop, automorphic inverse property loop(AIPL), cross inverse property loop(CIPL), K-loop, Bruck-loop, Kikkawa-loop

[‡]All correspondence to be addressed to this author.

A loop(quasigroup) is a weak inverse property loop (quasigroup)[WIPL(WIPQ)] if and only if it obeys the identity

$$x(yx)^\rho = y^\rho \quad \text{or} \quad (xy)^\lambda x = y^\lambda.$$

A loop(quasigroup) is a cross inverse property loop(quasigroup)[CIPL(CIPQ)] if and only if it obeys the identity

$$xy \cdot x^\rho = y \quad \text{or} \quad x \cdot yx^\rho = y \quad \text{or} \quad x^\lambda \cdot (yx) = y \quad \text{or} \quad x^\lambda y \cdot x = y.$$

A loop(quasigroup) is an automorphic inverse property loop(quasigroup)[AIPL(AIPQ)] if and only if it obeys the identity

$$(xy)^\rho = x^\rho y^\rho \text{ or } (xy)^\lambda = x^\lambda y^\lambda$$

Consider (G, \cdot) and (H, \circ) being two distinct groupoids(quasigroups, loops). Let A, B and C be three distinct non-equal bijective mappings, that maps G onto H . The triple $\alpha = (A, B, C)$ is called an isotopism of (G, \cdot) onto (H, \circ) if and only if

$$xA \circ yB = (x \cdot y)C \quad \forall x, y \in G.$$

The set $SYM(G, \cdot) = SYM(G)$ of all bijections in a groupoid (G, \cdot) forms a group called the permutation(symmetric) group of the groupoid (G, \cdot) . If $(G, \cdot) = (H, \circ)$, then the triple $\alpha = (A, B, C)$ of bijections on (G, \cdot) is called an autotopism of the groupoid(quasigroup, loop) (G, \cdot) . Such triples form a group $AUT(G, \cdot)$ called the autotopism group of (G, \cdot) . Furthermore, if $A = B = C$, then A is called an automorphism of the groupoid(quasigroup, loop) (G, \cdot) . Such bijections form a group $AUM(G, \cdot)$ called the automorphism group of (G, \cdot) .

The left nucleus of L denoted by $N_\lambda(L, \cdot) = \{a \in L : ax \cdot y = a \cdot xy \quad \forall x, y \in L\}$. The right nucleus of L denoted by $N_\rho(L, \cdot) = \{a \in L : y \cdot xa = yx \cdot a \quad \forall x, y \in L\}$. The middle nucleus of L denoted by $N_\mu(L, \cdot) = \{a \in L : ya \cdot x = y \cdot ax \quad \forall x, y \in L\}$. The nucleus of L denoted by $N(L, \cdot) = N_\lambda(L, \cdot) \cap N_\rho(L, \cdot) \cap N_\mu(L, \cdot)$. The centrum of L denoted by $C(L, \cdot) = \{a \in L : ax = xa \quad \forall x \in L\}$. The center of L denoted by $Z(L, \cdot) = N(L, \cdot) \cap C(L, \cdot)$.

As observed by Osborn [22], a loop is a WIPL and an AIPL if and only if it is a CIPL. The past efforts of Artzy [2, 3, 4, 5], Belousov and Tzurkan [6] and recent studies of Keedwell [17], Keedwell and Shcherbacov [18, 19, 20] are of great significance in the study of WIPLs, AIPLs, CIPQs and CIPLs, their generalizations(i.e m-inverse loops and quasigroups, (r,s,t)-inverse quasigroups) and applications to cryptography. For more on loops and their properties, readers should check [8],[10], [12], [13], [27] and [24].

Interestingly, Adeniran [1] and Robinson [25], Oyebo and Adeniran [23], Chiboka and Solarin [11], Bruck [7], Bruck and Paige [9], Robinson [26], Huthnance [14] and Adeniran [1] have respectively studied the holomorphs of Bol loops, central loops, conjugacy closed loops, inverse property loops, A-loops, extra loops, weak inverse property loops, Osborn loops and Bruck loops. Huthnance [14] showed that if (L, \cdot) is a loop with holomorph (H, \circ) , (L, \cdot) is a WIPL if and only if (H, \circ) is a WIPL. The holomorphs of an AIPL and a CIPL are yet to be studied.

For the definitions of inverse property loop (IPL), Bol loop and A-loop readers can check earlier references on loop theory.

Here ; a K-loop is an A-loop with the AIP, a Bruck loop is a Bol loop with the AIP and a Kikkawa loop is an A-loop with the IP and AIP.

1.2 Smarandache Quasigroups And Loops

The study of Smarandache loops was initiated by W. B. Vasantha Kandasamy in 2002. In her book [27], she defined a Smarandache loop (S-loop) as a loop with at least a subloop which forms a subgroup under the binary operation of the loop. In [16], the present author defined a Smarandache quasigroup (S-quasigroup) to be a quasigroup with at least a non-trivial associative subquasigroup called a Smarandache subsemigroup (S-subsemigroup). Examples of Smarandache quasigroups are given in Muktibodh [21]. In her book, she introduced over 75 Smarandache concepts on loops. In her first paper [28], on the study of Smarandache notions in algebraic structures, she introduced Smarandache : left(right) alternative loops, Bol loops, Moufang loops, and Bruck loops. But in [15], the present author introduced Smarandache : inverse property loops (IPL), weak inverse property loops (WIPL), G-loops, conjugacy closed loops (CC-loop), central loops, extra loops, A-loops, K-loops, Bruck loops, Kikkawa loops, Burn loops and homogeneous loops.

A loop is called a Smarandache A-loop(SAL) if it has at least a non-trivial subloop that is a A-loop.

A loop is called a Smarandache K-loop(SKL) if it has at least a non-trivial subloop that is a K-loop.

A loop is called a Smarandache Bruck-loop(SBRL) if it has at least a non-trivial subloop that is a Bruck-loop.

A loop is called a Smarandache Kikkawa-loop(SKWL) if it has at least a non-trivial subloop that is a Kikkawa-loop.

If L is a S-groupoid with a S-subsemigroup H , then the set $SSYM(L, \cdot) = SSYM(L)$ of all bijections A in L such that $A : H \rightarrow H$ forms a group called the Smarandache permutation(symmetric) group of the S-groupoid. In fact, $SSYM(L) \leq SYM(L)$.

The left Smarandache nucleus of L denoted by $SN_\lambda(L, \cdot) = N_\lambda(L, \cdot) \cap H$. The right Smarandache nucleus of L denoted by $SN_\rho(L, \cdot) = N_\rho(L, \cdot) \cap H$. The middle Smarandache nucleus of L denoted by $SN_\mu(L, \cdot) = N_\mu(L, \cdot) \cap H$. The Smarandache nucleus of L denoted by $SN(L, \cdot) = N(L, \cdot) \cap H$. The Smarandache centrum of L denoted by $SC(L, \cdot) = C(L, \cdot) \cap H$. The Smarandache center of L denoted by $SZ(L, \cdot) = Z(L, \cdot) \cap H$.

Definition 1.1 *Let (L, \cdot) and (G, \circ) be two distinct groupoids that are isotopic under a triple (U, V, W) . Now, if (L, \cdot) and (G, \circ) are S-groupoids with S-subsemigroups L' and G' respectively such that $A : L' \rightarrow G'$, where $A \in \{U, V, W\}$, then the isotopism $(U, V, W) : (L, \cdot) \rightarrow (G, \circ)$ is called a Smarandache isotopism(S-isotopism).*

Thus, if $U = V = W$, then U is called a Smarandache isomorphism, hence we write $(L, \cdot) \simeq (G, \circ)$.

But if $(L, \cdot) = (G, \circ)$, then the autotopism (U, V, W) is called a Smarandache autotopism (S-autotopism) and they form a group SAUT(L, \cdot) which will be called the Smarandache

autotopism group of (L, \cdot) . Observe that $SAUT(L, \cdot) \leq AUT(L, \cdot)$. Furthermore, if $U = V = W$, then U is called a Smarandache automorphism of (L, \cdot) . Such Smarandache permutations form a group $SAUM(L, \cdot)$ called the Smarandache automorphism group (SAG) of (L, \cdot) .

Let L be a S-quasigroup with a S-subgroup G . Now, set $H_S = (G, \cdot) \times SAUM(L, \cdot)$. If we define 'o' on H_S such that $(\alpha, x) \circ (\beta, y) = (\alpha\beta, x\beta \cdot y)$ for all $(\alpha, x), (\beta, y) \in H_S$, then $H_S(L, \cdot) = (H_S, \circ)$ is a quasigroup.

If in L , $s^\lambda \cdot s\alpha \in SN(L)$ or $s\alpha \cdot s^\rho \in SN(L) \forall s \in G$ and $\alpha \in SAUM(L, \cdot)$, (H_S, \circ) is called a Smarandache Nuclear-holomorph of L , if $s^\lambda \cdot s\alpha \in SC(L)$ or $s\alpha \cdot s^\rho \in SC(L) \forall s \in G$ and $\alpha \in SAUM(L, \cdot)$, (H_S, \circ) is called a Smarandache Centrum-holomorph of L hence a Smarandache Central-holomorph if $s^\lambda \cdot s\alpha \in SZ(L)$ or $s\alpha \cdot s^\rho \in SZ(L) \forall s \in G$ and $\alpha \in SAUM(L, \cdot)$.

The aim of the present study is to investigate the holomorphic structure of Smarandache AIPLs and CIPLs (SCIPLs and SAIPLs) and use the results to draw conclusions for Smarandache K-loops (SKLs), Smarandache Bruck-loops (SBRLs) and Smarandache Kikkawa-loops (SKWLs). This is done as follows.

1. The holomorphic structure of AIPQs (AIPLs) and CIPQs (CIPLs) are investigated. Necessary and sufficient conditions for the holomorph of a quasigroup(loop) to be an AIPQ (AIPL) or CIPQ (CIPL) are established. It is shown that if the holomorph of a quasigroup(loop) is a AIPQ (AIPL) or CIPQ (CIPL), then the holomorph is isomorphic to the quasigroup(loop). Hence, the holomorph of a quasigroup(loop) is an AIPQ (AIPL) or CIPQ (CIPL) if and only if its automorphism group is trivial and the quasigroup(loop) is a AIPQ (AIPL) or CIPQ (CIPL). Furthermore, it is discovered that if the holomorph of a quasigroup(loop) is a CIPQ (CIPL), then the quasigroup(loop) is a flexible unipotent CIPQ (flexible CIPL of exponent 2).
2. The holomorph of a loop is shown to be a SAIPL, SCIPL, SKL, SBRL or SKWL respectively if and only if its SAG is trivial and the loop is a SAIPL, SCIPL, SKL, SBRL, SKWL respectively.

2 Main Results

Theorem 2.1 *Let (L, \cdot) be a quasigroup(loop) with holomorph $H(L)$. $H(L)$ is an AIPQ (AIPL) if and only if*

1. $AUM(L)$ is an abelian group,
2. $(\beta^{-1}, \alpha, I) \in AUT(L) \forall \alpha, \beta \in AUM(L)$ and
3. L is a AIPQ (AIPL).

Proof

A quasigroup(loop) is an automorphic inverse property loop (AIPL) if and only if it obeys the identity

Using either of the definitions of an AIPQ(AIPL), it can be shown that $H(L)$ is a AIPQ(AIPL) if and only if $AUM(L)$ is an abelian group and $(\beta^{-1}J_\rho, \alpha J_\rho, J_\rho) \in AUT(L) \forall \alpha, \beta \in AUM(L)$. L is isomorphic to a subquasigroup(subloop) of $H(L)$, so L is a AIPQ(AIPL) which implies $(J_\rho, J_\rho, J_\rho) \in AUT(L)$. So, $(\beta^{-1}, \alpha, I) \in AUT(L) \forall \alpha, \beta \in AUM(L)$.

Corollary 2.1 *Let (L, \cdot) be a quasigroup(loop) with holomorph $H(L)$. $H(L)$ is a CIPQ(CIPL) if and only if*

1. $AUM(L)$ is an abelian group,
2. $(\beta^{-1}, \alpha, I) \in AUT(L) \forall \alpha, \beta \in AUM(L)$ and
3. L is a CIPQ(CIPL).

Proof

A quasigroup(loop) is a CIPQ(CIPL) if and only if it is a WIPQ(WIPL) and an AIPQ(AIPL). L is a WIPQ(WIPL) if and only if $H(L)$ is a WIPQ(WIPL).

If $H(L)$ is a CIPQ(CIPL), then $H(L)$ is both a WIPQ(WIPL) and a AIPQ(AIPL) which implies 1., 2., and 3. of Theorem 2.1. Hence, L is a CIPQ(CIPL). The converse follows by just doing the reverse.

Corollary 2.2 *Let (L, \cdot) be a quasigroup(loop) with holomorph $H(L)$. If $H(L)$ is an AIPQ(AIPL) or CIPQ(CIPL), then $H(L) \cong L$.*

Proof

By 2. of Theorem 2.1, $(\beta^{-1}, \alpha, I) \in AUT(L) \forall \alpha, \beta \in AUM(L)$ implies $x\beta^{-1} \cdot y\alpha = x \cdot y$ which means $\alpha = \beta = I$ by substituting $x = e$ and $y = e$. Thus, $AUM(L) = \{I\}$ and so $H(L) \cong L$.

Theorem 2.2 *The holomorph of a quasigroup(loop) L is a AIPQ(AIPL) or CIPQ(CIPL) if and only if $AUM(L) = \{I\}$ and L is a AIPQ(AIPL) or CIPQ(CIPL).*

Proof

This is established using Theorem 2.1, Corollary 2.1 and Corollary 2.2.

Theorem 2.3 *Let (L, \cdot) be a quasigroups(loop) with holomorph $H(L)$. $H(L)$ is a CIPQ(CIPL) if and only if $AUM(L)$ is an abelian group and any of the following is true for all $x, y \in L$ and $\alpha, \beta \in AUM(L)$:*

1. $(x\beta \cdot y)x^\rho = y\alpha$.
2. $x\beta \cdot yx^\rho = y\alpha$.
3. $(x^\lambda \alpha^{-1} \beta \alpha \cdot y\alpha) \cdot x = y$.
4. $x^\lambda \alpha^{-1} \beta \alpha \cdot (y\alpha \cdot x) = y$.

Proof

This is achieved by simply using the four equivalent identities that define a CIPQ(CIPL):

Corollary 2.3 *Let (L, \cdot) be a quasigroup(loop) with holomorph $H(L)$. If $H(L)$ is a CIPQ(CIPL) then, the following are equivalent to each other*

1. $(\beta^{-1}J_\rho, \alpha J_\rho, J_\rho) \in AUT(L) \forall \alpha, \beta \in AUM(L)$.
2. $(\beta^{-1}J_\lambda, \alpha J_\lambda, J_\lambda) \in AUT(L) \forall \alpha, \beta \in AUM(L)$.
3. $(x\beta \cdot y)x^\rho = y\alpha$.
4. $x\beta \cdot yx^\rho = y\alpha$.
5. $(x^\lambda \alpha^{-1} \beta \alpha \cdot y\alpha) \cdot x = y$.
6. $x^\lambda \alpha^{-1} \beta \alpha \cdot (y\alpha \cdot x) = y$.

Hence,

$$(\beta, \alpha, I), (\alpha, \beta, I), (\beta, I, \alpha), (I, \alpha, \beta) \in AUT(L) \forall \alpha, \beta \in AUM(L).$$

Proof

The equivalence of the six conditions follows from Theorem 2.3 and the proof of Theorem 2.1. The last part is simple.

Corollary 2.4 *Let (L, \cdot) be a quasigroup(loop) with holomorph $H(L)$. If $H(L)$ is a CIPQ(CIPL) then, L is a flexible unipotent CIPQ(flexible CIPL of exponent 2).*

Proof

It is observed that $J_\rho = J_\lambda = I$. Hence, the conclusion follows.

Remark 2.1 *The holomorphic structure of loops such as extra loop, Bol-loop, C-loop, CC-loop and A-loop have been found to be characterized by some special types of automorphisms such as*

1. Nuclear automorphism(in the case of Bol-,CC- and extra loops),
2. central automorphism(in the case of central and A-loops).

By Theorem 2.1 and Corollary 2.1, the holomorphic structure of AIPLs and CIPLs is characterized by commutative automorphisms.

Theorem 2.4 *The holomorph $H(L)$ of a quasigroup(loop) L is a Smarandache AIPQ(AIPL) or CIPQ(CIPL) if and only if $SAUM(L) = \{I\}$ and L is a Smarandache AIPQ(AIPL) or CIPQ(CIPL).*

Proof

Let L be a quasigroup with holomorph $H(L)$. If $H(L)$ is a SAIPQ(SCIPQ), then there exists a S-subquasigroup $H_S(L) \subset H(L)$ such that $H_S(L)$ is a AIPQ(CIPQ). Let $H_S(L) = G \times SAUM(L)$ where G is the S-subquasigroup of L . From Theorem 2.2, it can be seen that $H_S(L)$ is a AIPQ(CIPQ) if and only if $SAUM(L) = \{I\}$ and G is a AIPQ(CIPQ). So the conclusion follows.

Corollary 2.5 *The holomorph $H(L)$ of a loop L is a SKL or SBRL or SKWL if and only if $SAUM(L) = \{I\}$ and L is a SKL or SBRL or SKWL.*

Proof

Let L be a loop with holomorph $H(L)$. Consider the subloop $H_S(L)$ of $H(L)$ such that $H_S(L) = G \times SAUM(L)$ where G is the subloop of L .

1. Recall that by [Theorem 5.3, [9]], $H_S(L)$ is an A-loop if and only if it is a Smarandache Central-holomorph of L and G is an A-loop. Combing this fact with Theorem 2.4, it can be concluded that: the holomorph $H(L)$ of a loop L is a SKL if and only if $SAUM(L) = \{I\}$ and L is a SKL.
2. Recall that by [25] and [1], $H_S(L)$ is a Bol loop if and only if it is a Smarandache Nuclear-holomorph of L and G is a Bol-loop. Combing this fact with Theorem 2.4, it can be concluded that: the holomorph $H(L)$ of a loop L is a SBRL if and only if $SAUM(L) = \{I\}$ and L is a SBRL.
3. Following the first reason in 1., and using Theorem 2.4, it can be concluded that: the holomorph $H(L)$ of a loop L is a SKWL if and only if $SAUM(L) = \{I\}$ and L is a SKWL.

References

- [1] J. O. Adeniran (2005), *On holomorphic theory of a class of left Bol loops*, Al.I.Cuza 51, 1, 23–28.
- [2] R. Artzy (1955), *On loops with special property*, Proc. Amer. Math. Soc. 6, 448–453.
- [3] R. Artzy (1959), *Crossed inverse and related loops*, Trans. Amer. Math. Soc. 91, 3, 480–492.
- [4] R. Artzy (1959), *On Automorphic-Inverse Properties in Loops*, Proc. Amer. Math. Soc. 10,4, 588–591.
- [5] R. Artzy (1978), *Inverse-Cycles in Weak-Inverse Loops*, Proc. Amer. Math. Soc. 68, 2, 132–134.
- [6] V. D. Belousov (1969), *Crossed inverse quasigroups(CI-quasigroups)*, Izv. Vyss. Ucebn; Zaved. Matematika 82, 21–27.

- [7] R. H. Bruck (1944), *Contributions to the theory of loops*, Trans. Amer. Math. Soc. 55, 245–354.
- [8] R. H. Bruck (1966), *A survey of binary systems*, Springer-Verlag, Berlin-Göttingen-Heidelberg, 185pp.
- [9] R. H. Bruck and L. J. Paige (1956), *Loops whose inner mappings are automorphisms*, The annuals of Mathematics, 63, 2, 308–323.
- [10] O. Chein, H. O. Pflugfelder and J. D. H. Smith (1990), *Quasigroups and Loops : Theory and Applications*, Heldermann Verlag, 568pp.
- [11] V. O. Chiboka and A. R. T. Solarin (1991), *Holomorphs of conjugacy closed loops*, Scientific Annals of A.I.I.Cuza. Univ. 37, 3, 277–284.
- [12] J. Dene and A. D. Keedwell (1974), *Latin squares and their applications*, the English University press Lts, 549pp.
- [13] E. G. Goodaire, E. Jespers and C. P. Milies (1996), *Alternative Loop Rings*, NHMS(184), Elsevier, 387pp.
- [14] E. D. Huthnance Jr.(1968), *A theory of generalised Moufang loops*, Ph.D. thesis, Georgia Institute of Technology.
- [15] T. G. Jaíyéolá (2006), *An holomorphic study of the Smarandache concept in loops*, Scientia Magna Journal, 2, 1, 1–8.
- [16] T. G. Jaíyéolá (2006), *Parastrophic invariance of Smarandache quasigroups*, Scientia Magna Journal, 2, 3, 48–53.
- [17] A. D. Keedwell (1999), *Crossed-inverse quasigroups with long inverse cycles and applications to cryptography*, Australas. J. Combin. 20, 241–250.
- [18] A. D. Keedwell and V. A. Shcherbacov (2002), *On m -inverse loops and quasigroups with a long inverse cycle*, Australas. J. Combin. 26, 99–119.
- [19] A. D. Keedwell and V. A. Shcherbacov (2003), *Construction and properties of (r, s, t) -inverse quasigroups I*, Discrete Math. 266, 275–291.
- [20] A. D. Keedwell and V. A. Shcherbacov, *Construction and properties of (r, s, t) -inverse quasigroups II*, Discrete Math. 288 (2004), 61–71.
- [21] A. S. Muktibodh (2006), *Smarandache Quasigroups*, Scientia Magna Journal, 2, 1, 13–19.
- [22] J. M. Osborn (1961), *Loops with the weak inverse property*, Pac. J. Math. 10, 295–304.
- [23] Y. T. Oyebo and O. J. Adeniran, *On the holomorph of central loops*, Pre-print.

- [24] H. O. Pflugfelder (1990), *Quasigroups and Loops : Introduction*, Sigma series in Pure Math. 7, Heldermann Verlag, Berlin, 147pp.
- [25] D. A. Robinson (1964), *Bol loops*, Ph. D thesis, University of Wisconsin, Madison, Wisconsin.
- [26] D. A. Robinson (1971), *Holomorphic theory of extra loops*, Publ. Math. Debrecen 18, 59–64.
- [27] W. B. Vasantha Kandasamy (2002), *Smarandache Loops*, Department of Mathematics, Indian Institute of Technology, Madras, India, 128pp.
- [28] W. B. Vasantha Kandasamy (2002), *Smarandache Loops*, Smarandache notions journal, 13, 252–258.