

# Pulsus Journals

## How many points are there in a line segment? - A new answer from discrete-cellular space viewpoint --Manuscript Draft--

<b>Manuscript Number:</b>	Pulsus Journals-17-599R1
<b>Full Title:</b>	How many points are there in a line segment? - A new answer from discrete-cellular space viewpoint
<b>Short Title:</b>	How many points are there in a line segment?
<b>Article Type:</b>	Letter to the Editor
<b>Section/Category:</b>	JOURNAL OF PURE AND APPLIED MATHEMATICS
<b>Keywords:</b>	Euclid's five axioms; Euclidean geometry; discrete-cellular space, discrete cosmology models, lattice universe, cellular automata, cell biology, cellular computing
<b>Corresponding Author:</b>	Victor Christianto, DDiv. Institut Pertanian Malang INDONESIA
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	Institut Pertanian Malang
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Victor Christianto, DDiv.
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	Victor Christianto, DDiv. Florentin Smarandache, Dr.
<b>Order of Authors Secondary Information:</b>	
<b>Manuscript Region of Origin:</b>	INDONESIA
<b>Abstract:</b>	While it is known that Euclid's five axioms include a proposition that a line consists at least of two points, modern geometry avoid consistently any discussion on the precise definition of point, line, etc. It is our aim to clarify one of notorious question in Euclidean geometry: how many points are there in a line segment? - from discrete space viewpoint. In retrospect, it may offer an alternative of quantum gravity, i.e. by exploring discrete gravitational theories. To elucidate our propositions, in the last section we will discuss some implications of discrete cellular-space model in several areas of interest: (a) cell biology, (b) cellular computing, (c) Maxwell equations, (d) low energy fusion, and (e) cosmology modelling.
<b>Suggested Reviewers:</b>	<p>Thee Houw Liong, Dr. Prof. emeritus, Institut Teknologi Bandung thehl007@gmail.com He is a well-known professor of physics in Indonesia, with some advanced knowledge on computational physics</p> <p>Sir Roger Penrose, Prof. Mathematics professor, University of Oxford rouse@maths.ox.ac.uk He is leading and renowned professor in mathematical-physics in the world</p> <p>David Tong, Dr. Professor of Theoretical Physics, University of Cambridge d.tong@damtp.cam.ac.uk He is a renowned theoretical physicist</p> <p>Wolfgang Sproessig, Dr.</p>

	<p>Prof. in mathematical physics, Technische Universitat Bergakademie Freiberg  sproessig@math.tu-freiberg.de  He is a renowned mathematical physicist</p>
	<p>Hendra Gunawan, Dr.  Professor of Mathematics, Institut Teknologi Bandung  hgunawan82@gmail.com  he is a renowned mathematics professor in Indonesia</p>
<p><b>Opposed Reviewers:</b></p>	<p>Stephen Hawking, Dr.  professor, University of Cambridge  Prof.Hawking@damtp.cam.ac.uk  He seems to be too involved in some variations of quantum gravity, blackhole etc.</p>

# How Many Points are there in a Line Segment? – A new answer from Discrete-Cellular Space viewpoint

Victor Christianto\*<sup>1</sup>, Florentin Smarandache<sup>2</sup>

<sup>1</sup>Malang Institute of Agriculture (IPM), Malang, Indonesia. Founder of [www.ketindo.com](http://www.ketindo.com)

\*Email: [victorchristianto@gmail.com](mailto:victorchristianto@gmail.com). URL: [http://researchgate.net/profile/Victor\\_Christianto](http://researchgate.net/profile/Victor_Christianto)

<sup>2</sup>Dept. Mathematics and Sciences, University of New Mexico, Gallup – USA. Email:  
[florentin.smarandache@laposte.net](mailto:florentin.smarandache@laposte.net)

## ABSTRACT

While it is known that Euclid's five axioms include a proposition that a line consists at least of two points, modern geometry avoid consistently any discussion on the precise definition of point, line, etc. It is our aim to clarify one of notorious question in Euclidean geometry: how many points are there in a line segment? – from discrete-cellular space (DCS) viewpoint. In retrospect, it may offer an alternative of quantum gravity, i.e. by exploring discrete gravitational theories. To elucidate our propositions, in the last section we will discuss some implications of discrete cellular-space model in several areas of interest: (a) cell biology, (b) cellular computing, (c) Maxwell equations, (d) low energy fusion, and (e) cosmology modelling.

Keywords: discrete-cellular space, discrete cosmology models, lattice universe, cellular automata, cell biology, cellular computing

*“...I consider it quite possible that physics cannot be based on the field concept, i.e., on continuous structures. In that case nothing remains of my entire castle in the air gravitation theory included, -and of- the rest of modern physics.” A. Einstein*

## 1. Introduction

So many students from all ages have asked this question: *how many points are there in a line segment?* And a good math teacher will answer politely: in the circumference of a circle there are infinite number of points[1]. Similarly one can also ask: how many lines

are there in a rectangular? The answer again is known: there are infinite number of lines in given rectangular.

But a careful student will ask again: but what is the definition of point and line? Teacher will answer again: a point is a circle with zero diameter, and line is composed of infinite points.

If our beloved student persists, he/she will continue to ask: but teacher, if a circle has zero diameter, then an infinite number of zeroes will not make a finite line, isn't it?

At this time, there is fair chance that the teacher feel upset and say: "*shut up and calculate!*"

That is what usually happens in most primary school mathematics classroom, and the situation is not getting better in undergraduate classroom. Only in graduate math class, then the students are allowed to ask numerous questions, such as foundations of mathematics etc. A more serious debate among mathematicians over this notorious *continuum problem* has been recorded in ref. [11].

Here we will offer a simpler solution of the above posed question from a discrete-cellular space (DCS) viewpoint, with wide implications, including more clarity over distinction between quantization and discretization.

## **2. Solution: the space consists of circles with finite diameter (discrete cellular model)**

The obvious paradox that we set in the introduction section can be simplified as follows:

$$0+0+0+\dots ad\ infinitum = 0$$

Therefore the basic postulate that a line segment consists of circles with zero diameter is contradictory by itself.

Our proposed solution is to assume that the space consists of circles with small but finite diameter ( $z$ ), therefore if a line segment consists of circles like that, we have:

$$z+z+z+ \dots ad\ infinitum = \text{finite line}$$

One implication of this proposition is that we should better consider the geometry of space not as continuum, but as a discrete-cellular space. And we must remember that discretization of space is much more fundamental than quantization.

Moreover, we can consider the following:

- a. It can be shown that similar indeterminacy problem plagues the very definition of differential calculus, as no one knows that actual size of  $dx$ . See H.J.M. Bos [2]:

*2.15. I turn now to a difficulty which necessarily arises in any attempt to set up an infinitesimal calculus which takes the differential as fundamental concept, namely the indeterminacy of differentials.*

*The first differential  $dx$  of the variable  $x$  is infinitely small with respect to  $x$ , and it has the same dimension as  $x$ . These are the only conditions it has to satisfy, and they do not determine a unique  $dx$ , for if  $dx$  satisfies the conditions then clearly so do  $2dx$  and  $\frac{1}{2}dx$  and in general all  $adx$  for finite numbers  $a$ . That is, all quantities that have the same dimension and the same order of infinity as  $dx$  might serve as  $dx$ .*

*Moreover, there are elements not from this class which satisfy the conditions for  $dx$ ; for instance  $dx^2/a$  and  $\sqrt{adx}$ , for finite positive  $a$  of the same dimension as  $x$ .  $dx^2/a$  is infinitely small with respect to  $dx$  and  $\sqrt{adx}$  is infinitely large with respect to  $dx$ , so that there is even not a privileged class of infinite smallness from which  $dx$  has to be chosen; there is no "first" class of infinite smallness adjacent to finiteness. Thus first-order differentials involve a fundamental indeterminacy.*

- b. Boyer has shown that Planck blackbody radiation can be derived from discrete charge assumption (without partition as assumed by Planck). See [3].
- c. Lee Smolin has described three approaches to quantum gravity in his book[4]. But considering our proposition above, it seems that the notion of quantum gravity may be not necessary. Instead, we should consider discrete gravity theories.
- d. Gary W. Gibbons and George F.R. Ellis have considered a discrete Newtonian cosmology. That is a good start [5].

- e. Gerard 't Hooft has proposed a discrete deterministic interpretation of QM.[6] But it seems the use of both discrete and quantum language are superfluous. We need to let go the quantum terminology with its own excess baggage.
- f. At astronomical scale, Conrad Ranzan has proposed a cellular universe, which is essentially a Newtonian Steady-State model but with a discrete cellular space model.[7] In our view, such an approach needs to be explored and investigated further. See also our recent paper, where we suggest an ultradiscrete KdV as model of cosmology [8]. See also Lindquist-Wheeler model [9][10]. We discuss this approach in the last example of last section of this paper.
- g. It may be possible for certain conditions, to consider a partially continuum and partially discrete space. In other words, we may have a hybrid space. But we do not investigate it yet.

### **3. A bit of philosophical considerations**

According to Miguel Lorente, in order to understand better these models it would be useful to consider three levels of human knowledge in the comprehension of the physical world[12]:

Level 1: Physical magnitudes, such as distances, intervals, force, mass, charge, that are given by our sensation and perceptions.

Level 2: Mathematical structures, that are the result of metrical properties given by measurements and numerical relations among them.

Level 3: Fundamental concepts, representing the ontological properties of physical world given by our intelligence in an attempt to know the reality. This

level of knowledge is not accepted by some philosophical positions like logical positivism, conventionalism, neokantismus.

There must be some connections between the three levels. In QM the theoretical models of microphysics in level 2 are related to observable magnitudes in level 1 by correspondence laws. If we accept level 3 it should be connected to level 2 and to level 1 (through level 2). In fact, the rules governing the constructions of theoretical models in level 2 must be grounded in some fundamental (ontological) properties of the physical world.

It is also worth noting that there are different interpretations of the concepts of space and time. They are usually divided in three classes, as follows [12]:

(a) Dualistic theories: Space is a container where the particles and waves are moving.

Time is also a separated entity with respect to which the motion takes place. Therefore space and time are absolute and can be thought of in the absence of particles (Newton).

(b) Monistic theories: Spacetime is identified with some properties of matter and can not be conceivable without the existence of the later. The field of forces and also the sources are nothing more that geometrical deformations of the Spacetime (Einstein, Kaluza-Klein, Wheeler).

(c) Relational theories: Spacetime consists of the set of relations among some fundamental objects: monads (Leibniz), units (Penrose), processes (Weisaecker, Finkelstein), preparticles (Bunge, Garc'ia Sucre), objects (Hilbert).

In the present paper, following our argument in the previous section, we assert that the space consists of discrete cells with finite dimension, which is the most realistic model to

the best of our knowledge. Next we will discuss some implications in different areas of interest.

#### **4. Proof of concept: A few implications of discrete-cellular space**

To elucidate our propositions, in this section we will discuss some implications of discrete cellular-space model in several areas of interest: (a) cell biology, (b) cellular computing, (c) Maxwell equations, (d) low energy fusion, and (e) cosmology modelling.

(a) **Cell biology.** The mathematical modeling of cell populations can be, broadly speaking, split into two categories: continuum and discrete models. Discrete models treat cells as individual entities and hence provide a natural framework within which to make use of an increasing amount of experimental data available at the cellular and subcellular scales. There are now many different types of discrete cell-level models used to describe cell populations, e.g., cellular automata, cellular Potts models, cell-vertex, and off-lattice cell based model. [19]

While continuum models have their own advantages, they also have certain limitations, as follows [20]: Continuum models of the cell aim at capturing its passive dynamics. In addition to the limitations mentioned above, current models do not yet typically account for active biology: deformations and stresses experienced as a direct consequence of biochemical responses of the cell to mechanical load cannot be predicted by current continuum models. However, by contrasting the predicted purely mechanical cell response to experimental observations, one could isolate phenomena involving active biology, such as cell contraction or migration, from the passive mechanical response of the cell. Alternatively, continuum models might be

envisioned that account for active processes through time-dependent properties or residual strains that are linked to biological processes. Another limitation of continuum models stems from lack of description of cytoskeletal fibers. As such, they are not applicable for micromanipulations of the cell with a probe of the same size or smaller than the cytoskeletal mesh ( $\sim 0.1\text{--}1.0\ \mu\text{m}$ ). This includes most AFM experiments. In addition, the continuum models exclude small Brownian motions due to thermal fluctuations of the cytoskeleton, which would correspond to fluctuations of the network nodes in a continuum model and have been shown to play a key role in cell motility (Mogilner and Oster, 1996). Finally, continuum models have so far employed a limited number of time constants to characterize the cell's behavior. However, cells have recently been shown to exhibit behaviors with power-law rheology implying a continuous spectrum of time scales (Fabry et al., 2001; Desprat et al., 2004). In the meantime, models involving a finite number of time constants consistent with the time scale of the experimental technique can be used, recognizing their limitations.[20]

- (b) **Cellular computer.** Around 18 years ago, Sipper described a number of interesting features of cellular computer. He began his article by noting that von Neumann's architecture—which is based upon the principle of one complex processor that sequentially performs a single complex task at a given moment—has dominated computing technology for the past 50 years. Recently, however, researchers have begun exploring alternative computational systems based on entirely different principles. Although emerging from disparate domains, the work behind these systems shares a common computational philosophy, which can be called as cellular

computing [21]. Those cellular computers are supposed to have three principles in common. Combining these three principles results in the definition cellular computing = simplicity + vast parallelism + locality. Because the three principles are highly interrelated, attaining vast parallelism, for example, is facilitated by the cells' simplicity and local connectivity. Changing any single term in the equation results in a different computational paradigm. So, for example, foregoing the simplicity property results in the distributed computing paradigm. Cellular computing has been placed further along the parallelism axis to emphasize the "vastness" aspect.[21]

What specific application areas invite a cellular computing approach? Research has raised several possibilities: (1) Image processing. Applying cellular computers to perform image-processing tasks arises as a natural consequence of their architecture. For example, in a two-dimensional grid, a cell (or group of cells) can correspond to an image pixel, with the machine's dynamics designed to perform a desired image-processing task. Research has shown that cellular image processors can attain high performance and exhibit fast operation times for several problems. (2) Fast solutions to NP-complete problems. Even if only a few such problems can be dealt with, doing so may still prove highly worthwhile. NP-completeness implies that a large number of hard problems can be efficiently solved, given an efficient solution to any one of them. The list of NP-complete problems includes hundreds of cases from several domains, such as graph theory, network design, logic, program optimization, and scheduling, to mention but a few. (3) Generating long sequences of high-quality random numbers. This capability is of prime import in domains such as computational physics and computational chemistry. Cellular computers may prove a

good solution to this problem. (4) Nanoscale calculating machines. Cellular computing's ability to perform arithmetic operations raises the possibility of implementing rapid calculating machines on an incredibly small scale. These devices could exceed current models' speed and memory capacity by many orders of magnitude. (5) Novel implementation platforms. Such platforms include reconfigurable digital and analog processors, molecular devices, and nanomachines.[21]

- (c) **Maxwell equations.** While X.S. Wang[23] was able to derive the above mentioned Maxwell's equations in vacuum based on a continuum mechanics model of vacuum and a singularity model of electric charges, in the meantime Krasnoholovets has managed to show quite remarkably that the very definition of charge can be modelled from the viewpoint of tessellate space.[22] He argued that the Maxwell Equations are the manifestation of hidden dynamics of surface fractals. He also concludes that James Clerk Maxwell was right when he involved imaginary *cogwheels* constructing the equations of motion of the electromagnetic field [22, p. 128].
- (d) **Low energy fusion.** Since the early years of Condensed Matter Nuclear Science (aka. LENR/Cold fusion), Robert W. Bussard from Energy/Matter Conversion Corporation has argued in favor of internal nuclear fusion in metal lattice to explain the low energy reaction as reported by Pons and Fleischmann.[24] Subsequently, there are a number of researchers who have explored the implications of lattice vibration and lattice structure models from solid state physics in order to explain CMNS process.[25][26][27]. Such approaches seem to be quite promising and they are

worthy to continue further. For a recent discussion on discrete and continuum modelling, see for instance [28].

- (e) **Cosmology modelling.** Many physicists and philosophers alike have debated a long standing puzzle: whether the space is continuous or discrete. It has been known for long time that most of the existing cosmology models rely on pseudo-Riemannian metric as the cornerstone of Einsteinian universe. But the metric itself is based on continuum theory. It is known that such models have led us to too many (monster) problems, including dark matter and dark energy etc. Now what if the universe is discrete? Then perhaps we can solve these problems naturally. Philosophically speaking, the notion of discrete space can be regarded as basic question in definition of differential calculus and limit. If it is supposed that space is continuous then we can use standard differential calculus, but if we assume it is finite and discrete, then we should use difference equation or finite difference theories. This problem is particularly acute when we want to compute our mathematical models in computers, because all computers are based on discrete mathematics. Then we can ask: is it possible that the discrete mathematics can inspire cosmology theorizing too? Despite majority of cosmologists rely on such a Standard Model which is called Lambda CDM theory, we will explore here the redshift theory based on a few of lattice-cellular models, including Lindquist-Wheeler theory and beyond it. We will also touch briefly some peculiar models such as Voronoi tessellattice and also Conrad Ranzan's cellular model of the Universe.

- a. Lindquist-Wheeler's theory:

In this model, the matter content is assumed to be discrete; identical spherically symmetric islands uniformly distributed in a regular lattice. This attempt was first introduced in 1957 by Lindquist and Wheeler (LW) in a seminal paper. While LW suggested that their global dynamics is similar to Friedmann universe for closed dust dominated universe, Shalaby has shown that LW-model can be extended to yield a redshift equation, as follows:[9]

$$1 + z = 1 + \langle \gamma \rangle \ln \left( \frac{a_r}{a_e} \right) = 1 + \langle \gamma \rangle \ln(1 + z_{FRW}) \cong (1 + z_{FRW})^{\langle \gamma \rangle} \quad (1)$$

It can be shown, that the value of  $\langle \gamma \rangle$  approximates geometrically to be 2/3, however, numerically its value was estimated to be 7/10.[9] Liu also analyzed LW model, and he concludes that the LW redshifts can differ from their FLRW counterparts by as much as 30%, even though they increase linearly with FLRW redshifts, and they exhibit a non-zero integrated Sachs-Wolfe effect, something which would not be possible in matter-dominated FLRW universes without cosmological constant. [10]

b. Voronoi Tessellation model:

Rien van de Weygaert describes a novel model based on Voronoi tessellation. The spatial cosmic matter distribution on scales of a few up to more than a hundred Megaparsec displays a salient and pervasive foam-like pattern.[14] Voronoi tessellations are a versatile and flexible mathematical model for such weblike spatial patterns. Cellular patterns may be the source of an intrinsic geometrically biased clustering. However, so far we do not find a redshift equation from this model.[13]

c. Non-expanding cellular universe:

Conrad Ranzan suggests a DSSU cellular cosmology (dynamics steady state universe), which he claims to be problem-free. The cosmic redshift is shown to be a velocity-differential effect caused by a flow differential of the space medium. He obtains the cosmic redshift equation in its basic form:[7]

$$z = (1 + z_{GC})^N - 1 \quad (2)$$

There are of course other cellular cosmology models, some of them have been reviewed by Marmet, but this paper is not intended for such an exhaustive list of redshift models. See Marmet [18].

## 5. Concluding Remarks

An old question and paradox in Euclidean geometry may be resolved consistently, once we accept and assume a discrete space instead of continuum model which is full of indeterminacies.

Many implications and further developments can be expected both in particle physics realm and also in cosmology theorizing. More observation and experiments are recommended to verify whether the space is discrete, continuous, or hybrid.

In retrospect, it may offer an alternative of quantum gravity, i.e. by exploring discrete gravitational theories. To elucidate our propositions, in the last section we discuss some implications of discrete cellular-space model in several areas of interest: (a) cell biology, (b) cellular computing, (c) Maxwell equations, (d) low energy fusion, and (e) cosmology modelling.

## Acknowledgement

The first author (VC) would like to thank to participants of 3<sup>rd</sup> *Conference on Theoretical Physics and Nonlinear Phenomena*, held in Bandung at 1<sup>st</sup> August 2017, whose discussions have inspired this paper. And special thanks to Prof. (Em.) Thee Houw Liong from Bandung Institute of Technology (ITB) for encouraging advises, and also to anonymous reviewer for suggesting improvement of this paper. We also gratefully appreciate Dr. Volodymyr Krasnoholovets for sending his new book describing tessellation model of space and its physical implications.

## Document history:

- version 1.0: 29<sup>th</sup> August 2017, pk. 16:14
- version 1.1: 31<sup>st</sup> August 2017, pk. 16:18
- version 1.2: 29<sup>th</sup> January 2018, pk. 22:11

## References:

- [1] <https://math.stackexchange.com/questions/1799125/how-many-points-in-a-line-segment>
- [2] H.J.M. Bos. Differentials, Higher Order Differentials, and the Derivative in the Leibnizian calculus. *Arch. Hist. Exact. Sci.* vol. 14, p. 24
- [3] Timothy H. Boyer. Connecting blackbody radiation, relativity, and discrete charge in classical electrodynamics. arXiv: physics/0605003 (2006)
- [4] Lee Smolin. *Three roads to quantum gravity*. Basic Books, 2001.
- [5] Gary Gibbons and George Ellis . Discrete Newtonian Cosmology. *Classical and Quantum Gravity* vol. 31, no. 2, (2013), see also arXiv: 1308.1852 (2013). url: <http://iopscience.iop.org/article/10.1088/0264-9381/31/2/025003/meta>
- [6] Gerard 't Hooft. A Deterministic Interpretation of Quantum Mechanics. url: [www.rug.nl/research/vsi/events/groenewold/thooft.pdf](http://www.rug.nl/research/vsi/events/groenewold/thooft.pdf)
- [7] Conrad Ranzan. Large scale structure of the Dynamic Steady State Universe. *American Journal of Astronomy and Astrophysics* 2016, 4(6): 65-77. <http://www.sciencepublishinggroup.com>. See also [www.cellularuniverse.org](http://www.cellularuniverse.org)
- [8] Victor Christianto & Florentin Smarandache & Yunita Umniyati. An outline of Cellular Automaton Universe via Cosmological KdV equation. Published at *Prespacetime Journal*, and this paper was also presented at CTPNP 3<sup>rd</sup>, 1<sup>st</sup> Aug 2017, held in Bandung, Indonesia.
- [9] Mohamad Shalaby. Dynamics and Light Propagation in a Universe with Discrete-Matter Content. arXiv: 1205.68773 [gr-qc]
- [10] Rex G. Liu. The Lindquist-Wheeler formulation of lattice universes. arXiv: 1501.05169 [gr-qc]

- [11] <https://www.ias.edu/ideas/2011/kennedy-continuum-hypothesis>
- [12] Miguel Lorente. A realistic interpretation of lattice gauge theories. arXiv: hep-lat/0312044v1 (2003)
- [13] Rien van de Weygaert. Voronoi Tessellations and the Cosmic Web: spatial patterns and clustering across the Universe. Proc. of the 4th International Symposium on Voronoi Diagrams in Science and Engineering, p.230-239, July 09-11, 2007 [doi>10.1109/ISVD.2007.48]. url: <http://www.astro.rug.nl/~weygaert/tim1publication/vorwey.isvd07.pdf>
- [14] A.A. Kirillov & D. Turaev. Foam-like structure of the Universe. *Phys. Lett. B* 656 (2007) 1-9.
- [15] Conrad Ranzan. Cosmic redshift in the non-expanding cellular universe: velocity-differential theory of cosmic redshift. *American Journal of Astronomy and Astrophysics* 2014, 2(5): 47-60
- [16] \_\_\_\_\_. Large scale structure of the Dynamic Steady State Universe. *American Journal of Astronomy and Astrophysics* 2016, 4(6): 65-77. <http://www.sciencepublishinggroup.com>
- [17] Chris Hand. Simple cellular automata on a spreadsheet. *CHEER*, vol. 17, 2005. url: <http://economicsnetwork.ac.uk/cheer/ch17/hand.htm>
- [18] Louis Marmet. On the Interpretation of Red-Shifts: A Quantitative Comparison of Red-Shift Mechanisms II. <http://www.marmet.org>
- [19] Philip J. Murray *et al.* From a discrete to a continuum model of cell dynamics in one dimension. *Phys. Rev. E* 80 (2009).
- [20] Mohammad R. K. Mofrad, Helene Karcher, and Roger D. Kamm. Continuum elastic or viscoelastic models of the cell. Chapter 4 in *Cytoskeletal Mechanics: Models and Measurements in Cell Mechanics*. <https://books.google.co.id/books?isbn=1139458108>
- [21] M. Sipper, The Emergence of Cellular Computing, *IEEE Computer*, vol. 32, no. 7, pp. 18-26, July 1999. url: <https://www.cs.bgu.ac.il/~sipper/publications/cellcomp.pdf>
- [22] Volodymyr Krasnoholovets. Structure of Space and the Submicroscopic Deterministic Concept of Physics. USA: Apple Academic Press & CRC, 2017. P. 118-145
- [23] Xiao-Song Wang. Derivation of the Maxwell's Equations Based on a Continuum Mechanical Model of Vacuum and a Singularity Model of Electric Charges. arXiv: physics/0609027 (2006)
- [24] Robert W. Bussard. Virtual state internal nuclear fusion in metal lattice. *Fusion Technology*, Volume 16, September 1989.
- [25] Steven E. Jones, E.P. Palmer, J.B. Czirr, D.L. Decker, G.L. Jensen, J.M. Thorne, S.F. Taylor, and J. Rafelski; "Observations of Cold Nuclear Fusion in Condensed Matter," *Nature* 338: 737, 740, April 1989
- [26] Martin Fleischmann and Stanley Pons; "Electrochemically Induced Nuclear Fusion of Deuterium," *Journal Electroanalytical Chemistry*, 261, 301, 1989.

[26] V.B. Belyaev et al. Nuclear fusion induced by X-rays in a crystal. arXiv: 1603.01723 (2016)

[27] Yoshiaki Arata & Yue-Chang Zhang. Formation of condensed metallic deuterium lattice and nuclear fusion. *Proc. Japan Acad.*, 78, Ser. B (2002), 57-60.

[28] V. A. Kuzkin, A. M. Krivtsov, E. A. Podolskaya & M. L. Kachanov (2016): Lattice with vacancies: elastic fields and effective properties in frameworks of discrete and continuum models, *Philosophical Magazine*, DOI: 10.1080/14786435.2016.1167979