

# Visualizing $\zeta(n \geq 2)$ and Proving Its Irrationality

Timothy W. Jones

December 21, 2017

## 1 Introduction

Apery's proof that

$$\zeta(3) = \sum_{k=1}^{\infty} \frac{1}{k^3}$$

is irrational is difficult [1]. It doesn't generalize to show  $\zeta(2n+1)$  is irrational for  $n > 1$ . Here we develop a much simpler proof that does so generalize.

The current state of affairs with proving  $z_n$ ,  $n$  odd, is irrational is quite limited. It is known that there are infinitely many odd  $n > 3$  that are irrational [11] and that at least one of 5, 7, 9, and 11 are irrational [16]. The proofs of these result uses group theory and complex analysis. Zudilin gives a literature review and develops both results in [15]. The even case follows easily from the transcendence of  $\pi$  [5, 10] and Bernoulli's famous formula:

$$\zeta(2n) = (-1)^{k+1} \frac{(2\pi)^{2k} B_{2k}}{2(2k)!}.$$

This formula is derived from a trigonometric series expansion [2].

The attempts on the part of Zudilin and others reflect the combinatorial problem of the general case. One certainly senses that showing  $\zeta(2)$  is irrational using Apery's ideas [4] is easier than showing  $\zeta(3)$  and, closely reading [15] one sees Apery's ideas are generating an ever increasing combinatorial puzzle. A way to *see* the similarities of all cases is our theme.

The visualization part of this article starts by exploring decimal representations. If some real number requires an infinite number of unambiguous

digits in all basis, then it must be irrational. Hardy shows that all decimal representations of a rational number  $a/b$  in a given base  $d$  are finite, repeating, or mixed depending on the relationship between  $b$  and  $d$  [9]. The ambiguous case of say  $.4\bar{9} = .5$  in base ten is not included as an infinite representation of the finite decimal  $.5$ , base 10. An irrational number in all bases is an infinite non-repeating decimal. The idea of the first part of the paper is to suggest that

$$\zeta(n) - 1 = z_n = \sum_{k=2}^{\infty} \frac{1}{k^n} \tag{1}$$

can't be represented by a finite decimal in any base.

Our visualization involves a simple geometric construction that allows the terms of (1) to be given as sector areas and to be added. There is then some connection with circles on the plain, but our plain is not the complex plain, nor even the Cartesian plain – just concentric circles with sectors designated by a radius. If the radius goes through a point (we call it a dot) on a circle the sector area is given by a single decimal in a base associated with the circle. The construction allows for a clear visualization of the decimal representations of all terms, Section 2, and partial sums, Section 3, of (1) in all bases  $k^n$ , where  $k$  is a natural number greater than 1.<sup>1</sup>

In Section 4 we consider the limit of partials using  $.\bar{1}$ , base 4. The circles associated with this series generate finite decimals base 4, but no single circle or finite addition (finite decimal, base 4) can give the convergence value of  $1/3$ . If this is generally true, then the convergence point of  $z_n$  must not reside as a dot on any of its term's circles, but its term's circles give all finite decimal representations in bases  $k^n$ . But this is all rational numbers between 0 and 1, so  $z_n$  must be irrational. We have some grounds to suspect the irrationality of  $z_n$ , all  $n > 1$ .<sup>2</sup>

The proof part of the paper consists of two Sections: Section 5 (Bertrand) gives a Theorem that uses Bertrand's postulate and yields a corollary used in Section 6 (Cantor). In Section 5 we give two lemmas and a theorem with a corollary. We show that partials of all  $z_n$  can't be expressed as finite decimals in any base  $d$  where  $d$  is the denominator of one of the partial's terms. The limiting case, then, is the rub [12]. Section 6 gives a proof that  $z_2$  is irrational

---

<sup>1</sup>Henceforth, just bases  $k^n$ .

<sup>2</sup>Henceforth, just  $z_n$ .

that uses the corollary from Section 5 and is based on Cantor's classic proof that the real numbers are not countable [6].

Cantor's diagonal method consists of modifying a list of decimal numbers, supposed to be all reals in a fixed base with values between 0 and 1. Each digit down the diagonal of the list is modified, yielding a number that is not in the list, contradicting all reals have been enumerated. We first give a variation of this proof to show how it can be used to construct an irrational number. We then give a bolder modification of Cantor's technique. We list all rational numbers between 0 and 1 using all bases  $k^2$ , referencing the visualization sections' circles and the corollary of Section 5. We construct a number not associated with any rational number using partial sums of  $z_2$ , one after the other. The resultant infinite series, limit of the partials, we claim must be irrational.

In the conclusion, Section 7, we mention other series and argue that the result developed to show  $z_2$  is irrational applies to other  $z_n$ , mutatis mudantis.

## 2 Term Visualization

The series  $z_2$  is referenced in what follows, but any  $z_n$ ,  $n \geq 2$  can similarly be referenced.




Figure 1: A circle with radius  $1/\sqrt{\pi}$  has an area of 1.

We can visualize the first term,  $1/4$ , of  $z_2$  using a circle. In Figure 1 we

have a circle of radius  $\sqrt{1/\pi}$ . The area of this circle is

$$\pi r^2 = \pi \cdot (\sqrt{1/\pi})^2 = 1.$$

In Figure 2, four equally spaced dots are placed around the circle, giving four equal sector areas. Each area must be  $1/4$  of the area of the circle or  $1/4$ . Sector areas corresponding to these dots, between 0 and 1, are given by  $x/4$ ,  $x = 1, 2, 3$  or a single, non-zero decimal base 4. If a radius on the circle doesn't go through one of the dots, the sector area formed will require more than a single decimal in base 4: Figure 3. We will designate this circle with  $C_4$ .




Figure 2: A circle with area 1 is divided up using  $2^2$ . The area of the shaded sector is  $1/4$ .

The next term is  $1/9$ . The circle in Figure 4 has radius  $\sqrt{2/\pi}$  with 9 equally spaced dots around it. Its area is 2:

$$\pi r^2 = \pi \cdot (\sqrt{2/\pi})^2 = 2.$$

We will designate this circle with  $C_9$ .

By making  $C_4$  and  $C_9$  concentric circles, Figure 5, the area of the annulus formed is  $2 - 1$ . If a radius is drawn through a dot on  $C_9$ , it will generate a sector area of  $x/9$  on  $C_4$ . If a radius misses dots on both circles, then the sector area formed is not equal to a single decimal in either base 4 or base 9. It will require more than a single digit in either of these bases.

Figure 6 shows the first three terms of  $z_2$  rendered with  $C_4$ ,  $C_9$ , and  $C_{16}$ . Clearly, we can continue this process using equally spaced  $k^2$  dots on circles




Figure 3: A radius that does not go through any dot generates a sector area that requires more than one decimal, base 4.




Figure 4: Nine equally spaced dots on a circle of radius  $\sqrt{2/\pi}$ :  $C_9$ .

of radius  $\sqrt{(k-1)/\pi}$ . If a given radius misses all dots on all such circles, the sector area associated with it must be irrational. This follows as the sector areas generated by radii through a given dot, say  $x$ , on the  $C_{k^2}$  circle will be given by  $.x$  base  $k^2$ , a single decimal digit, and all rational numbers can be so designated;  $km/k^2 = m/k$  with  $m < k$ .




Figure 5:  $C_4$  and  $C_9$  as concentric circles. The area of the blue sector is  $1/9$ .




Figure 6: The shaded sector area is not a single decimal in base 4, 9, or 16.

This is a visualization of the terms of  $z_2$ . Next we will visualize adding these terms.

### 3 Visualization of Partial Sums

Two sector areas can be added.




Figure 7: The addition of  $1/4 + 1/9$  using  $C_4$  and  $C_9$  with  $C_9$  offset.

In Figure 7,  $1/4$  is added to  $1/9$  by rotating  $C_9$  in a counter-clockwise direction to line up with the  $1/4$  dot on  $C_4$ . This addition is somewhat analogous to the head to toe (here 1 to 0) method of vector addition. In Figure 8,  $1/16$  is added to  $1/4 + 1/9$  using the same 1 (head) to 0 (toe) method. The resulting radius generates an area on all annuli and  $C_4$ 's circle that corresponds to  $1/4 + 1/9 + 1/16$ . Clearly these additions can be used to form such radii for all partial sums of  $z_2$ .




Figure 8: The addition of  $1/4 + 1/9 + 1/16$  using  $C_4$ ,  $C_9$ , and  $C_{16}$  with the offset method. The area of the shaded sector is the sum.






Figure 9: The radius associated with the sum  $1/4 + 1/9 + 1/16$  misses all dots on  $C_4$ ,  $C_9$ , and  $C_{16}$ .

Figure 7 and Figure 8 show rotations of  $C_9$  and  $C_{16}$  to effect fraction additions. Figure 9 shows the resulting radius with the un-rotated versions of these circles. Designating partial sums,

$$s_k^m = \frac{1}{2^m} + \frac{1}{3^m} + \cdots + \frac{1}{k^m},$$

Figure 9 accurately shows that the partial

$$s_4^2 = \sum_{k=2}^4 \frac{1}{k^2}$$

generates a radius that does not go through any of its first few term's dotted circles. We can infer that the sum is not expressible as a single decimal digit in base 4, 9, or 16.

The denominators of  $z_2$  are just all decimal bases squared. So if a radius misses all dots on all  $C_{k^2}$  circles then its associated sector area value must require more than one decimal in all  $k^2$  bases. It must be irrational.

We can now visualize the problem of proving  $z_2$  is irrational. We need to show that the limit radius generated by adding the terms of  $z_2$  does not

go through any of the dots on any of the circles defined by its terms. The difficulty is that radii can converge to a dot on a circle without a radius going through the dot. The geometric series gives guidance. We will analyze it next.

## 4 Geometric series

Infinite repeating decimals are really geometric series. For example, in base 4,

$$.\bar{1} = \sum_{k=1}^{\infty} \frac{1}{4^k}.$$

This geometric series has a convergence point of  $1/3$ . All its terms occur in  $z_2$ , so we can use our dotted concentric circles to understand the relationship between the rotated  $C_{4^k}$ ,  $k > 1$ , circles used to construct this sum and  $C_3$ , the un-rotated circle having a dot the sum converges to: that is, the unique radius for this convergence point.

Here are some observations. Given any radius, representing a sector area's value, we can read from a system of dotted circles the decimal expansion in a given base, like base 4; conversions via the modulus operator may be necessary to adjust the digits of the expansion. Also all convergent infinite series with terms of the form  $1/a_k$  with  $a_k$  strictly increasing natural numbers have partial radii that rotate counter-clockwise around the circle and go through points on concentric circles farther and farther from the center. This forces series that converge to a rational number to have their convergence radius given by a radius going through an un-rotated *earlier* dot. We can see these patterns in Figure 10. As  $z_2$  and generally  $z_k$  require rotations of all circles giving all rational numbers, there is no such earlier un-rotated circle having a rational point for these series to converge to. The additions of the terms perpetually offsets the radius formed from all rational numbers. This suggests that all  $z_k$  are irrational.

Another observation: there is only one radius for every area, rational and irrational. Unlike decimal representations where  $.\overline{49} = .5$ , there is no ambiguity with reduced fractions and areas. For an irrational number, we can read the decimals from our figure and as the, note *the*, radius never goes through a point in all bases, it never terminates in all bases. If the number were rational, then its denominator would have prime factors that are not

shared by any natural number, a contradiction.

Also note that we observe trajectories, the radii of Figure 10, and how additions build new trajectories. It seems plausible that adjusting a trajectory with additions could cause new trajectories to miss all previous dots as well as the last term added's dots. That is we can perpetually adjust a trajectory to have it miss all dots. Think of a spaceship avoiding equally spaced meteorites arranged in concentric rings as the dots of  $C_{k^2}$  in front of us. We can avoid them all and we know, per convergence of  $z_n$ , that a single radius will emerge and be an irrational number.




Figure 10: Circles  $C_4$  with  $\overline{C_{16}}$  and  $\overline{C_{64}}$  rotated (indicated with overline) to generate the radius associated with  $.111$  base 4.  $C_3$ , unrotated, has the convergence point for  $.\overline{1}$ :  $1/3$ .

Showing the radius for  $z_k$  never goes through a dot on the  $n^k$  system of concentric circles, shows that it must be irrational. In the next two sections we prove that the limit radius for  $z_2$  does not go through any  $C_{k^2}$  dot.

## 5 Bertrand

Our aim in this section is to show that the reduced fractions that give the partial sums of  $z_n$  require a denominator greater than that of the last term defining the partial sum. Restated this says that partial sums can't be expressed as a finite decimal using for a base the denominators of any of the partial sum's terms.

The first lemma is a little more difficult than an exercise in Apostol's Introduction to Analytic Number Theory [2, p. 23, problem 30], its inspiration.

**Lemma 1.** *The reduced fraction,  $r/s$  giving*

$$s_k^m = \sum_{j=2}^k \frac{1}{j^m} = \frac{r}{s} \quad (2)$$

*is such that  $2^m$  divides  $s$ .*

*Proof.* The set  $\{2, 3, \dots, k\}$  will have a greatest power of 2 in it,  $a$ ; the set  $\{2^m, 3^m, \dots, k^m\}$  will have a greatest power of 2,  $ma$ . Also  $k!$  will have a powers of 2 divisor with exponent  $b$ ; and  $(k!)^m$  will have a greatest power of 2 exponent of  $mb$ . Consider

$$\frac{(k!)^m}{(k!)^m} \sum_{j=2}^k \frac{1}{j^m} = \frac{(k!)^m/2^m + (k!)^m/3^m + \dots + (k!)^m/k^m}{(k!)^m}. \quad (3)$$

The term  $(k!)^m/2^m$  will pull out the most 2 powers of any term, leaving a term with an exponent of  $mb - ma$  for 2. As all other terms but this term will have more than an exponent of  $2^{mb-ma}$  in their prime factorization, we have the numerator of (3) has the form

$$2^{mb-ma}(2A + B),$$

where  $2 \nmid B$  and  $A$  is some positive integer. This follows as all the terms in the factored numerator have powers of 2 in them except the factored term  $(k!)^m/2^m$ . The denominator, meanwhile, has the factored form

$$2^{mb}C,$$

where  $2 \nmid C$ . This leaves  $2^{ma}$  as a factor in the denominator with no powers of 2 in the numerator, as needed.  $\square$

**Lemma 2.** *If  $p$  is a prime such that  $k > p > k/2$ , then  $p^m$  divides  $s$  in (2).*

*Proof.* First note that  $(k, p) = 1$ . If  $p|k$  then there would have to exist  $r$  such that  $rp = k$ , but by  $k > p > k/2$ ,  $2p > k$  making the existence of a natural number  $r > 1$  impossible.

The reasoning is much the same as in Lemma 1. Consider

$$\frac{(k!)^m}{(k!)^m} \sum_{j=2}^k \frac{1}{j^m} = \frac{(k!)^m/2^m + \dots + (k!)^m/p^m + \dots + (k!)^m/k^m}{(k!)^m}. \quad (4)$$

As  $(k, p) = 1$ , only the term  $(k!)^m/p^m$  will not have  $p$  in it. The sum of all such terms will not be divisible by  $p$ , otherwise  $p$  would divide  $(k!)^m/p^m$ . As  $p < k$ ,  $p^m$  divides  $(k!)^m$ , the denominator of  $r/s$ , as needed.  $\square$

**Theorem 1.** *If*

$$s_k^m = \frac{1}{2^m} + \frac{1}{3^m} + \cdots + \frac{1}{k^m} = \frac{r}{s}, \quad (5)$$

*with  $r/s$  reduced, then  $s > k^m$ .*

*Proof.* Bertrand's postulate states that for any  $k \geq 2$ , there exists a prime  $p$  such that  $k < p < 2k$  [9]. If  $k$  of (5) is even we are assured that there exists a prime  $p$  such that  $k > p > k/2$ . If  $k$  is odd  $k - 1$  is even and we are assured of the existence of prime  $p$  such that  $k - 1 > p > (k - 1)/2$ . As  $k - 1$  is even,  $p \neq k - 1$  and  $p > (k - 1)/2$  assures us that  $2p > k$ , as  $2p = k$  implies  $k$  is even, a contradiction.

For both odd and even  $k$ , using Bertrand's postulate, we have assurance of the existence of a  $p$  that satisfies Lemma 2. Using Lemmas 1 and 2, we have  $2^m p^m$  divides the denominator of (5) and as  $2^m p^m > k^m$ , the proof is completed.  $\square$

So, for  $z_2$ , we have the following.

**Definition 1.**

$$D_{k^2} = \{0, 1/k^2, \dots, (k^2 - 1)/k^2\} = \{0, .1, \dots, .(k^2 - 1)\} \text{ base } k^2$$

**Corollary 1.**

$$s_n^2 \notin \bigcup_{k=2}^n D_{k^2}$$

*Proof.* Immediate.  $\square$

Corollary 1 shows that we have made progress in our goal of finding what properties are shared by all  $z_n$ . Referring to Figure 9, the partials for all  $z_n$  generate radii that miss all the dots on the circles associated with their terms. Next we use an application of Cantor to show the limit radius for these partials,  $z_n$  itself, misses all such dots.

## 6 Cantor

Cantor's diagonal method, mentioned in the introduction, is a technique that systematically eliminates numbers as it builds a convergence point. It involves an infinite procedure. Here's an example of its use to show the existence of an irrational number. List all rational numbers between 0 and 1. They are countable, so this can be done. Use base 10.

$$\begin{aligned}
 &.d_{11}d_{12}d_{13}\dots \\
 &.d_{21}d_{22}d_{23}\dots \\
 &.d_{31}d_{32}d_{33}\dots \\
 &.d_{41}d_{42}d_{43}\dots \\
 &.d_{51}d_{52}d_{53}\dots \\
 &\vdots
 \end{aligned}$$

Go down the diagonal and change the value of the decimal to 3, if it is not 3 and 7, if it is: Table 1. Record the changes following a decimal point.

row	new	original
1	$.c_1d_{12}d_{13}\dots$	$.d_{11}d_{12}\dots$
2	$.d_{21}c_2d_{23}\dots$	$.d_{21}d_{22}\dots$
3	$.d_{31}d_{32}c_3\dots$	$.d_{31}d_{32}\dots$
4	$.d_{41}d_{42}d_{43}c_4\dots$	$.d_{41}d_{42}\dots$

Table 1: Cantor's diagonal method building an irrational number:  $.c_1c_2\dots$

We notice that  $.c_1$  of row 1 is different  $.d_{11}d_{12}\dots$  and  $.c_1c_2$  is different than  $.d_{21}d_{22}\dots$  of row 2, as well as  $.d_{11}d_{12}\dots$  of row 1.

We have reduced the infinite construction of  $.c_1c_2\dots$  to finite considerations and we can conclude that the infinite decimal  $.c_1c_2c_3\dots$  is not in the list. As it is also between 0 and 1, it must be irrational. Think of the space ship trajectory given by the radius of earlier sections. We our building our trajectory by small increments and decrements avoiding the dots ahead. The result  $.c_1c_2\dots$  is a sum of discrete steering wheel corrections. It avoids all of the rational trajectories.

We have *constructed* an irrational number. In this application of Cantor's diagonal method, we can verify we have succeeded in constructing a number

not in our list: subtract  $.c_1c_2\dots$  from a number on the list and the result is not zero.

## 6.1 Modification of Cantor

Here's a modification of Cantor's method that will show  $z_2$  is irrational. List all the rational numbers between 0 and 1 using  $D_{k^2}$ . These are arranged down a diagonal in Table 2. Our mission is to create a number that isn't in the first row, then isn't the first or second row, and then repeat this process for all rows.

$D_4$							
	$D_9$						
		$D_{16}$					
			$D_{25}$				
				$D_{49}$			
					$D_{64}$		
						$D_{81}$	
							$\ddots$

Table 2: A list of all rational numbers between 0 and 1.

The process is to add the numbers above each  $D_{k^2}$ , for all  $k \geq 2$ , as given in Table 3. The result is not in  $D_{k^2}$ . This is Corollary 1. So, for example,  $1/4 + 1/9$  is not in  $D_4$ ,  $1/4 + 1/9$  is not in  $D_4$  or  $D_9$ ,  $1/4 + 1/9 + 1/16$  is not in  $D_4$ ,  $D_9$ , or  $D_{16}$ , etc.. Just like Cantor allows us to conclude a number we construct is not in a list, we can conclude the number we construct,  $z_2$  is not in the list. As our list consists of all rational numbers between 0 and 1,  $z_2$  must be irrational.

## 6.2 Observations

There are differences from Cantor's diagonal method. Cantor needs to be careful with his *if 3, 7 else 3* program. If he replaced everything with 9's or 0's then an ambiguity of  $.1\bar{9} = .2$  might arise; he might not have assurance that a number is excluded from the list. Working with fractions (or all bases



1/4							
1/9	1/4	1/4	1/4	1/4			
$D_4$	1/9	1/9	1/9	1/9			
	$D_9$	1/16	1/16	1/16			
		$D_{16}$	1/25	1/25			
			$D_{25}$	1/36			
				$D_{36}$			
					$D_{49}$		
						$D_{64}$	
							$\dots$

Table 3: A list of all rational numbers between 0 and 1 modified to exclude them all via partial sums of  $z_2$ .

or radii), as we are, and not a single number base, this problem does not arise. Also, worth noting is the absence of a notational verification that the convergence point  $z_2$  is not equal to some decimal expansion version of a rational number. As was mentioned, using Cantor to construct an irrational number using a single base, this is obtained.

In this regard, you might call our use of Cantor as strong Cantor (playing on strong induction): it is strictly eliminative; all rational possibilities are eliminated. The reasoning is like the following strong Cantor proof that the sum of all natural numbers is not a natural number. Using the sum of natural number from 1 to  $n$  is  $n(n + 1)/2$ , we can construct Table 4 and conclude that the infinite sum is not a natural number. We don't know what it is, only what it's not. For the  $z_k$  case, knowing the series converges to a real number and, having eliminated all rational numbers, only an irrational number is left.

1	1	1	...	1	
+2	+2	2	...	2	
$\notin \{1\}$	$\notin \{2\}$	+3	...	$\vdots$	
		$\notin \{3\}$	...	$\vdots$	
				$\vdots$	
				$n - 1$	
				$+n$	
				$\notin \{n\}$	
					$\ddots$

Table 4: Strong Cantor example showing the sum of all natural numbers is not a natural number.

## 7 Conclusion

### 7.1 Other series

The telescoping series

$$\sum_{k=2}^{\infty} \left( \frac{1}{n} - \frac{1}{n+1} \right) = 1/2 - 1/3 + 1/3 - 1/4 + \dots = 1/2$$

or

$$\sum_{k=2}^{\infty} \frac{1}{n(n+1)} = \frac{1}{6} + \frac{1}{12} + \frac{1}{20} + \dots = 1/2$$

shows the necessity of partials escaping terms. For example, the sum of the first three terms is 3/10 which can be expressed with 6/20 in  $D_{20}$ . Partial sums backtrack to earlier denominators thus preventing Cantor's diagonal process from being valid. The geometric series has partials that sum to fractions with denominators from the last term of the partial, but the term's denominators don't cover all pertinent rational numbers.

For both examples, placing them in Cantor's diagonal of Table 3 shows the necessity of partials escaping their terms and the terms *covering* the rationals.

## 7.2 General $z_n$

Do the ideas given here give a proof that  $\zeta(n \geq 2)$ ? As all bases  $k^n$  have the same prime factors as  $k$ , the answer is yes: Table 3, in conjunction with Section 5, works when these other series are used.

If one assumes Corollary 1, does the proof distill to a geometric proof? Note that the denominator of the partial sums of a  $z_k$  series with upper bound  $n$  will be much larger than  $n^k$ , more like  $(n!)^k$ , so this theorem is highly plausible. Also simple number theory proofs show that  $(n-1, n) = (n, n+1) = 1$ , that the natural numbers are consecutively relatively prime. So one suspects such partial sums will have denominators that have chaotically occurring prime factors. This points to the central intuition about this series; the fractions added have denominators growing by one (to a power) and this marks how the series differs from the “spaciness” of the geometric and telescoping series. If one grants Theorem 1 as intuitively plausible is Figure 9 of Section 3 enough: the nudging of a trajectory by the terms (the addition of terms) of any  $\zeta(n)$  builds a trajectory that never “hits” a rational dot; all rational sector areas are perpetually offset yielding a sector area that must be irrational – is that enough? For more on geometric proofs see Sondow’s proof of the irrationality of  $e$  [13]. His proof can be translated into a concentric circle version.

Finally, this paper suggests that one suspects a geometric self-similarity argument could be made: if one  $z_n$  is irrational, they all are. Hence, if this were true, via Apéry or Bernoulli, all odd  $z_n$  are irrational by similarity with these known cases.

## References

- [1] R. Apéry, Irrationalité de  $\zeta(2)$  et  $\zeta(3)$ , *Astérisque* **61** (1979), 11-13.
- [2] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer, New York, 1976.
- [3] J. Bertrand, *Memoire sur le nombre de valeurs que peut prendre une fonction quand on y permute les lettres qu’elle renferme*, J. Ec. Polyt., 30 (1845) 123-140.
- [4] F. Beukers, A Note on the Irrationality of  $\zeta(2)$  and  $\zeta(3)$ , *Bull. London Math. Soc.*, **11**, (1979), 268–272.

- [5] L. Berggren, J. Borwein, and P. Borwein, *Pi: A Source Book*, 3rd ed., Springer, New York, 2004.
- [6] R. Courant, H. Robbins, *What is Mathematics*, Oxford University Press, London, 1948.
- [7] P. Erdős, *Beweiss eines Satzes von Tschebyschef*, Acta Litt. Sci. Reg. Univ. Hungar, Fr.-Jos., Sect. Sci. Math., 5 (1932) 194-198.
- [8] P. Eymard and J.-P. Lafon, *The Number  $\pi$* , American Mathematical Society, Providence, RI, 2004.
- [9] G. H. Hardy, E. M. Wright, R. Heath-Brown, J. Silverman, and A. Wiles, *An Introduction to the Theory of Numbers*, 6th ed., Oxford University Press, London, 2008.
- [10] F. Lindemann, Über die Zahl  $\pi$ , *Math. Ann.* **20** (1882) 213–225.
- [11] Rivoal, T., La fonction zeta de Riemann prend une infinité de valeurs irrationnelles aux entiers impairs, *Comptes Rendus de l'Académie des Sciences, Série I. Mathématique* 331, (2000) 267-270.
- [12] W. Shakespeare, *The Riverside Shakespeare*, Oxford University Press, 1973.
- [13] J. Sondow, A geometric proof that  $e$  is irrational and a new measure of its irrationality, *Amer. Math. Monthly*, **113**, (2007), 637–641.
- [14] P. L. Tchebychef, *Memoire sur les nombres premiers*, St. Pet. Ac. Mm., VII (1854) 17-33.
- [15] W. W. Zudilin, Arithmetic of linear forms involving odd zeta values, *J. Théorie Nombres Bordeaux*, **16(1)**, (2004) 251–291
- [16] W. W. Zudilin, One of the numbers  $\zeta(5)$ ,  $\zeta(7)$ ,  $\zeta(9)$ ,  $\zeta(11)$  is irrational, *Russian Mathematical Surveys*, **56(4)**, (2001) 747–776.