

# The Incompatibility of the Planck Acceleration and Modern Physics? and the Solution in Mathematical Atomism

Espen Gaarder Haug\*  
Norwegian University of Life Sciences

November 2, 2017

## Abstract

The Planck second is likely the shortest relevant time interval. If the Planck acceleration lasts for one Planck second, one will reach the speed of light. Yet, according to Einstein, no particle with rest-mass can travel at the speed of light as this would require an infinite amount of energy. Modern physics is incompatible with the Planck acceleration in many ways. However in atomism we see that the Planck acceleration happens for the building blocks of the Planck mass and that the Planck mass is dissolved into energy within one Planck second. Further, the Planck mass stands absolutely still as observed from any reference frame. Atomism is fully consistent with the Planck acceleration. The relativistic Planck acceleration is unique among accelerations because it can only happen from absolute rest; it is therefore the same as the Planck acceleration. In other words, atomism predicts breaks in Lorentz invariance at the Planck scale, something several quantum gravity theories address as well. Atomism seems to solve a series of challenges in modern physics and this paper is one of a series in pointing this out.

**Key words:** Planck acceleration, relativistic Planck acceleration, Planck mass, mass gap, atomism.

## Planck Acceleration

The Planck acceleration is known to be

$$a_p = \frac{m_p c^3}{\hbar} = \frac{c}{t_p} = \frac{c^2}{l_p} \approx 5.5609210^{51} \quad (1)$$

where  $l_p$  is the Planck length and  $t_p$  is the Planck second, see [1]. In 1984, Scarpetta had already predicted this as the maximum acceleration possible, [2], something also suggested by [3]:

*“the ‘Planck acceleration’ is both the maximum acceleration for an elementary particle in free space and also the surface gravity of a black hole with minimum mass  $m_p$ ” – Falla and Landsberg 1994*

However, as pointed out by [4], for example, this enormous acceleration means that one will reach the speed of light after one Planck second,  $a_p t_p = c$ . Modern physicists do not really know exactly what the Planck length, the Planck mass, or the Planck second represent. They also do not know precisely how to incorporate them into the framework of modern physics. However, many physicists assume that the Planck second must be the shortest time interval possible. This lead to an interesting paradox. Nothing with rest-mass can undergo Planck acceleration, even for one Planck second. This because no object with rest-mass can travel at the speed of light, since this would require an infinite amount of energy, as first pointed out by Einstein, see [5]. So does this mean the Planck acceleration is meaningless, or merely fiction?

This is where mathematical atomism comes in. Here two indivisible particles, when they are colliding, are the very foundation of mass. When two indivisible particles collide, they become the Planck mass, see [6, 7]. This mass only lasts for one Planck second before the two colliding particles leave each other and turn into pure energy (light) once again. The indivisible particles have no rest-mass when they are not colliding; then they are massless. However, they have what we can call potential mass – this mass show up as rest-mass in the Planck second of collision, a moment when they are at absolute rest.

The smallest building blocks of the universe have a rest-mass equal to half of the Planck mass. Whenever we are working with rest-mass, we are always working with two indivisible particles. Thus the smallest rest-mass is the Planck mass, which lasts for one Planck second, and as applied to standard particles in modern physics it is more correct to claim that the rest-mass of an indivisible particle is  $5.86685 \times 10^{-52}$  kg (when colliding).

---

\*e-mail espenhaug@mac.com.

The atomism model is fully consistent with Planck acceleration and holds that we can go from a velocity of zero to the speed of light in one Planck second, as hypothetically measured with Einstein-Poincaré synchronized clocks. In contrast, no particle in the modern physics model can ever undergo Planck acceleration, even for one Planck second. So one has three choices: 1) introduce a shorter time interval than the Planck second, 2) claim that the Planck acceleration is not relevant for anything with rest-mass (so it is a purely fictitious acceleration), or 3) come up with yet another mathematical trick (fudge) to save the model. Naturally, one could claim that the Planck mass spontaneously radiates into energy within a Planck second. This is basically correct, but modern physics has no model for what would trigger this event. In fact, we (modern physics) do not even have a clear explanation for what energy actually is, as stated by Richard Feynman:

*It is important to realize that in physics today, we have no knowledge what energy is.*

## Relativistic Planck Acceleration

Assume a rocket (or particle) is accelerating with acceleration  $a$ , as measured from the rocket (the proper acceleration). The well-known relativistic acceleration is given by (see for example [8] and [9])

$$\alpha = \frac{a}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{3}{2}}} \quad (2)$$

Assume now that the acceleration as observed from the rest frame is the Planck acceleration,  $a = a_p$ . If one assumes that  $v > 0$ , then this will lead to a relativistic acceleration greater than then Planck acceleration. According to atomism, the Planck acceleration can only start to happen from a Planck mass (remember the Planck mass only lasts for one Planck second in this model), and the maximum velocity of any fundamental particle is (as described by Haug in a series of working papers as well as published papers [10, 11])

$$v_{max} = \sqrt{1 - \frac{l_p^2}{\lambda^2}} \quad (3)$$

for a Planck mass we have  $\bar{\lambda} = l_p$ , which leads to

$$v_{max} = \sqrt{1 - \frac{l_p^2}{l_p^2}} = 0 \quad (4)$$

In other words, the Planck mass is at rest, as observed from any reference frame. The Planck length, the Planck mass, and the Planck time are invariant as observed from any reference frame, in strong contrast to special relativity theory. This also means that the relativistic Planck acceleration is equal to the Planck acceleration

$$\alpha = \frac{a_p}{\left(1 - \frac{v_{max}^2}{c^2}\right)^{\frac{3}{2}}} = a_p \quad (5)$$

Actually, all Planck units are invariant across all reference frames. That is to say, Lorentz symmetry is broken at the very moment we reach Planck energy (the Planck scale). This is consistent with what is predicted by several quantum gravity theories, see for example [12]. In other words, we need a modification of Einstein's special relativity theory. All of Einstein's formulas are valid when using Einstein-Poincaré synchronized clocks, but atomism leads to the incorporation of key mathematical concepts introduced by Max Planck that are intuitive and very easy to understand.

## References

- [1] M. Planck. *The Theory of Radiation*. Dover 1959 translation, 1906.
- [2] G. Scarpetta. *Letter Nuovo Cimento*, 51, 1984.
- [3] D. F. Falla and P. T. Landsberg. Black holes and limits on some physical quantities. *European Journal of Physics*, 15, 1994.
- [4] E. G. Haug. Charged particle radiation power at the Planck scale. <http://vixra.org/pdf/1604.0228v2.pdf>, 2016.
- [5] A. Einstein. *Relativity: The Special and the General Theory*. Translation by Robert Lawson (1931), Crown Publishers, 1916.

- [6] E. G. Haug. The Planck mass particle finally discovered! The true God particle! Good bye to the point particle hypothesis! <http://vixra.org/abs/1607.0496>, 2016.
- [7] E. G. Haug. The mass gap, kg, the Planck constant and the gravity gap. <http://vixra.org>, 2017.
- [8] W. Pauli. Die relativittstheorie. *Encyclopdie der mathematischen Wissenschaften*, 5, 1921.
- [9] L. Marder. *Time and the Space-Traveler*. George Allen & Unwin, London, 1971.
- [10] E. G. Haug. The gravitational constant and the Planck units: A simplification of the quantum realm. *Physics Essays Vol 29, No 4*, 2016.
- [11] E. G. Haug. The ultimate limits of the relativistic rocket equation: The Planck photon rocket. *Acta Astronautica*, 136, 2017.
- [12] F. Kislak and H. Krawczynski. Planck-scale constraints on anisotropic Lorentz and CPT invariance violations from optical polarization measurements. *Physical Review D*, 95, 2017.