

# **Gravity and Light Speed**

## **A new Law of stellar Aberration**

by

Florian Michael Schmitt

January 2018, Berlin

### **Abstract**

All relevant experiments that disagree with static or entrained ether and led to the historical breakthrough of Special Relativity [1] will be revised from the scratch. It will be shown that an alternative model based on rightly understood gravitational dragging of light is able to explain the notorious phenomena. Invariance of light speed, time dilation and Lorentz contraction [2] will become obsolete. Focus is laid on the most problematic classic subjects as there are: Michelson/Morley experiment [3], Sagnac effect [4] and Michelson/Gale/Pearson [5] experiment, stellar and terrestrial aberration as well as anomaly of Mercury orbit shift. A series of fateful and fundamental misinterpretations will be disclosed. Chapter 4 will develop a revolutionary new law for stellar aberration based on wave nature of light.

### **1. Introduction**

The historic dispute about ether theories was circling around the experimental evidence, most importantly the Sagnac and Michelson/Gale experiment and the Michelson/Morley experiment. The Sagnac effect [4] as well as the Michelson/Gale/Pearson [5] experiment are esteemed to be disproving entrained ether but being in accordance with static ether, the Michelson/Morley [3] experiment attests the opposite. A similar picture is given by the problem of stellar and terrestrial aberration. Generally spoken, static ether concepts [2] were explaining stellar aberration but failed on terrestrial aberration, entrained ether concepts [6] [7] vice versa. Special Relativity [1] solved all the contradictions by postulating invariance of light speed, but at the expense of logical reason, and understanding of the nature of light was never brought to an end, when Special Relativity [1] prematurely terminated any further investigation on this topic. This paper is aiming to positively test a gravity entrained ether theory, whereby light is fully dragged by gravity, on many of the applicable experiments and on aberration.

## 2. The Michelson/Morley Experiment, tested on static and entrained ether

First of all there is the famous Michelson/Morley interferometer experiment, which was interpreted to be the mightiest of all arguments for Special Relativity.

The setup of this experiment simplified was as follows:




Fig. 1: Schematic setup of the Michelson/Morley experiment

By means of a 45° beam splitting mirror light was sent through two orthogonal pathes, joined together by the same mirror and projected on a screen or telescope, showing interference fringes once properly adjusted. Since one of the rays would be in line with earth's motion und therefore should have to overcome a longer or shorter distance, the other orthogonal ray would not be affected by earth's movement at all. The expected difference of travel distance should have been according to Michelson's well known formula:

$$\Delta l = 2 \cdot L \cdot \frac{v^2}{c^2} \quad (1)$$

Whereby L is the interferometer arm length, thus representing the travel length of light at rest. During its path to the right, the horizontal ray will have to catch up with the receding right hand mirror, on its way back the 45° mirror will move towards the ray, and for this reason the difference will have to be second order, i.e. depending on  $v^2/c^2$ . On an assumed interferometer arm length of 1m, movement velocity of earth around sun of approx. 30 km/s and light speed of approx.. 300.000 km/s, this would result to  $2 \times 10^{-8}$  m, equivalent to 0,04 times a wavelength (500nm), hence 0,04 fringes on the screen (all basically according to Michelson).

By turning the whole setup at 90° the fringes should shift now by 0,04, indicating the difference of travel distances of both rays. But the experiment gave a null result and was interpreted favouring the light speed to be invariant from the observer's movement.

It was never quite discussed to the end whether earth's orbital speed, earth's rotational speed, the solar system's speed around the galaxy or the total speed against CMB (cosmic microwave background) has to be called upon for calculation. Michelson obviously decided for the first. If he had opted (if he had known of) for the total speed against CMB, the difference would have been even 4 fringes, and the null result becomes even more distinct.

To put things in order, it is necessary to compare the circumstances and all velocities that are relevant for the experiment setup, and have a model for clarification. We now try to test the experiment on both ether theories, starting with static ether. The maximum and minimum available speeds have to be considered, i.e. speed against CMB and earth's rotational speed, since the experiment only deals with utmost diurnal period. For simplification and to draw an even clearer picture we assume the following rounded speeds: Light speed 300.000 km/s, CMB speed 300 km/s, earth's rotational speed 0,5 km/s. Furtheron we assume all speeds to be in one line, the orthogonal ray is unaffected and we point our focus only on the ray that is fully affected, i.e. the horizontal (Earth being represented by the circle):

- $v$  is the velocity of earth's rotation on equator
- $V$  is the total velocity of earth against CMB




Fig. 2: Relevant velocities on Michelson/Morley experiment in static ether

We obtain the following situation regarding the relevant speed that the mirrors move with or against the lightray:

First ray in direction of CMB:

$$+V+v$$

Second ray against direction of CMB:

$-V-v$

The difference of both rays now is:

$$+V+v-(-V)-(-v) = 2V+2v$$

Obviously, the total speed against CMB is adding up and earth's rotation as well to:

$$300 \text{ km/s} + 0,5 \text{ km/s} = 300,5 \text{ km/s}$$

Therefore the result of the experiment, according to (1), would have to be expected to be 4,013 times a fringe shift. **The experimentally obtained null result therefore is evidence that static ether is in contradiction with the Michelson/Morley experiment.** Many attempts have been done to find corrective processes such as deviated deflection on moving mirrors [8], deviations due to beam width, Lorentzian length contraction [2] etc. to explain the issue, according to the author's overview none of them being satisfactorily.

Now we come to the gravity entrained ether, where the picture becomes different:




Fig. 3: Relevant velocities on Michelson/Morley experiment in fully entrained ether

The speed against CMB does not account, because light is being fully entrained by earth's movement. Also the orbital speed around sun is irrelevant for the same reason.

Someone might say now that sun's gravitational influence must have to be taken into account, but at any place on earth's surface the effect is smaller by  $6 \times 10^{-4}$  than earth's gravity. Let alone earth's rotational speed will be relevant. Someone might say now, why of all the smallest speed? And if light is fully entrained by earth, should not no speed at all be relevant? And for both cases, would not actually centrifugal force partly

or fully cancel out the gravitational effects? And even further, would then light not have been accelerated to earth's rotational speed since long ago? At this point it is necessary to put right a couple of fundamental misunderstandings on the whole issue and do a brief Gedankenexperiment.

If one person is standing on earth, of course it will be already accelerated to the earth's rotational speed on surface, indeed we all become born already accelerated. But nevertheless there would be no such acceleration if there was not a single bonding of the rotating body (earth) and the body to be accelerated (person). If someone would be hovering on top of a rotating carousel, he would not at all acquire the carousel's speed, though he would still fall down to earth due to gravity! And of course it must be the same with light, travelling in or against earth's rotation, earth would just move away under its feet! And even centrifugal force, being a furious force, can grip and cancel out gravity only if a physical bonding is present. But gravity does not need such bonding, therefore, assuming light being dragged by gravity, light will be dragged along with earth's movement around the sun and even against the CMB, but not earth's rotation!

Insofar the assumption that only rotational speed remains relevant is plausible. And the Michelson/Morley experiment has to deal solely with the tiny rotational speed of earth, and also only on the second order effect thereof, giving an even tinier result. According to (1) the difference then should be only  $1 \times 10^{-5}$  of a fringe shift, equivalent to a difference of light speed of 0,4 mm/s, and the most accurate interferometer experiments ever done give no less than  $4 \times 10^{-4}$  of a fringe! **Therefore the gravity entrained ether holds good for explaining the Michelson/Morley experiment result quite well.**

### 3. The Sagnac effect and Michelson/Gale/Pearson Experiment, tested on static and entrained ether

First we will test the Sagnac experiment on the basis of a static ether. The setup of the Sagnac experiment simplified was as follows:




Fig. 4: Schematic setup of the Sagnac experiment

Contrarily to the Michelson/Morley experiment the two rays were forced into a full roundtrip rather than a back and forth path. Additionally the interferometer was mounted on a spinning disc and the angular speed would have to cause the movement of the mirrors with or against the light rays. The expected difference of distance should have been, deriving from Sagnac's well known formula:

$$\Delta t = 4 \cdot A \cdot \frac{\omega}{c^2} \quad (2)$$

Whereby  $\Delta t$  is the time difference,  $A$  the area enclosed by the light ray's roundtrip and  $\omega$  the angular speed.

Also we have:  $\omega = \frac{v}{r}$  and  $A = r^2 \cdot \pi$  thus:

$$\Delta t = 4 \cdot A \cdot \frac{\omega}{c^2} = 8 \cdot A \cdot \frac{v}{r \cdot c^2} = 8 \cdot \frac{A}{r} \cdot \frac{v}{c^2} = 8 \cdot r \cdot \pi \cdot \frac{v}{c^2}$$

and  $\Delta l = \Delta t \cdot c$  thus:

$$\Delta l = 8 \cdot r \cdot \pi \cdot \frac{v}{c} \quad (3)$$

So the difference of traveling distances should be amounting to a first order relation, i.e.  $v/c$ . Usually the Sagnac effect is interpreted to deal with disc's rotational speed only, which is not reasonable, since this speed is the smallest of all involved. We have to put this right, as per following image:

- $v$  is the rotational velocity of disc on disc's rim
- $V$  is the total velocity of earth against CMB




Fig. 5: Relevant velocities on Sagnac experiment in static ether

Assuming one light ray circling around the disc clockwise (red), we obtain for one cycle:

On the upper section:

$-V+v$

On the lower section:

$$+V+v$$

Adding up to  $+2v$ . All additions of speed perpendicular to CMB speed (i.e. orthogonal direction) are cancelling each other. Most important:  $V$  (speed against CMB) is shown to be cancelling out in general.

For the second ray (counterclockwise, green):

On the upper section:

$$+V-v$$

On the lower section:

$$-V-v$$

Adding up to  $-2v$ .  $V$  again is cancelling out.

The difference for both rays now is:

$$+2v - (-2v) = 4v$$

Indeed the Sagnac experiment gave a positive result with a difference as to be expected by (1) and (3). **The effect is in accordance with static ether** and shows the difference of light travel distances due to the movement of the observer towards one or the opposite direction on the disc against the light ray. The CMB speed is always cancelling out, a roundtrip route of the two light rays is required for this cancellation and consequently both rays will have to enclose an area. Commonly though it is surmised due to a somewhat automatic reflex that the effect must be in contradiction with entrained ether.

Now we draw the equivalent picture based on the assumption that light is being entrained by gravity:




Fig. 6: Relevant velocities on Sagnac experiment in fully entrained ether

Contrarily to the previous speculation we obtain the same situation as per static ether, i.e. difference amounts to  $4v$ , but for different reason. Total speed against CMB is not cancelling out, but does not account at all since light is fully dragged by earth's gravity, and only the speed on the disc's rim is playing a role. Therefore the difference of light traveling distance due to the movement of the observer on the rotating disc towards one or the opposite direction is still existing and in accordance with the experiment.

One might say now that in case of the Sagnac effect, centrifugal force on the rotating disc could become easily stronger than gravity, both light rays would be accelerated and glued towards the disc's rim, and the Sagnac experiment would have to give a null result. We have already pinpointed this argument before, there is no reason why light should be accelerated by a rotating disc unless having a physical bonding to it. **Therefore the Sagnac effect is also in accordance with gravity entrained ether.**

Finally we have to look at the Michelson/Gale/Pearson experiment:




Fig. 7: Schematic setup of the Michelson/Gale/Pearson experiment

The difference to the Sagnac effect is none but the following facts:

- Size of the setup is in the range of a kilometer
- The rotating disc is earth itself
- The lengths of the horizontally and vertically interferometer arms vary decisively in order to obtain as much of a result as possible.

Accepting the Gedankenexperiment we made before, i.e. that light is entrained by earth's gravity but not its rotation, it becomes clear **that also the Michelson/Gale/Pearson experiment is in accordance with gravity entrained ether.**

#### 4. Stellar and terrestrial aberration

Now there is one thing left to clarify, obviously the most difficult of all. Aberration was leading the discussion on ether theories from the beginning to the end. Several attempts have been successful [9] to explain the interferometer problems with entrained ether. But stellar aberration is still esteemed to be fundamentally



incompatible with entrained ether. We will show that this is by far not the case, but we will also reveal another basic misunderstanding, that might shudder our whole cosmic coordinate system.

First we will assume a static ether and purely wave nature of light in order to clarify the aberrational phenomena, i.e. movement of light source does not affect light propagation.

The classic explanation of stellar aberration [10] was, that similar to the falling rain drop, the telescope would have to be twisted in order to follow the light ray since the telescope itself was moving sidewise by earth rotation or earth orbiting respectively. If the ether on the other side was fully dragged by earth, no such aberration could occur at all, because the light ray would always follow earth's movement. On the other hand it was found difficult to explain aberration at all assuming a pure wave nature of light. Therefore the explanation was tacitly based on the idea that light is always coming as a directed beam rather than an undefined series of concentric spheres.




Fig. 8: Stellar aberration by classic explanation

The aberration angle was then calculated upon the distance that the telescope would move whilst the light ray is traveling from the telescope's lens to its mirror. The great misunderstanding is, that stellar aberration in truth does not have anything to do with the telescope having to follow the light path within the short distance inside the telescope nor the still short distance within earth's atmosphere nor even the short distance within the gravitational influence of earth or even the solar system, but the whole distance that light travels from its source, i.e. the distance from stars being billions of lightyears away.

The following images show, how a spherical wave of light will be emitted by its source deliberately long ago, whereby the observer is travelling by deliberate speed, and the task will be to find the point, where the circle, not the beam, meets the observer. From above we also must realize that the true position of the star is unknown and we have to start from a hypothetical middling angle of all observed angles, though this middling angle would show a position of the star that is never visible, because observation angles are always circling and ellipsing around this point. We will see later, that this interpretation makes up an important difference. For convenience the following model values were chosen:

- Light speed  $c$ : 1,5 km/sec
- Earth movement speed  $v$ : 0,6 km/s (on a range of 0,4 to 0,6 km/s) against static ether on orbital path
- Distance of earth path to light source: 1,5 km

- Middling angle of observation at 0,5 km/sec towards source: 60° degree (the angle between the true position of star and the observer at the time of observation, 46,10° in this case, is actually unknown)




Fig. 9: Light propagation from source with 0,3 sec steps, earth's speed 0,6 km/sec

It can be seen, that the observer is moving sidewise during the complete period that the light wave front travels from the source to meet the observer. It is important to mention that at this instant the light wave front hits the observer as a wave normal, and all subsequent wave fronts do as well. Only now we have established the angle, under which the light ray meets the observer, and we add a telescope for better understanding:




Fig. 10 Telescope directed to source.

In this model the observation angle will amount to 62,8542°, as shown per calculation later. All distances, angles and relations of speeds are on scale at the model, verified by means of cad.

Now the same procedure with 0,4 km/sec earth's movement speed:




Fig. 11: Light propagation from source with 0,3 sec steps, earth's speed 0,4 km/sec

Of course, as before, again the meeting point represents a series of wave normals. But as can be seen by adding the telescope, the observation angle this time is 57,1806°. Now the two images of both earth's movement speeds will be overlaid:




Fig. 12: Overlay both situations 0,6 and 0,4 km/sec

The angle between the two light rays now, in this case 5,6736° is deemed to be the common aberration angle. It is most important that this angle is deriving from the difference of the total earth's speed against the static ether e.g. CMB (Cosmic Microwave Background), but not necessarily the earth's speed on the orbit (being 0,2 km/sec in this model).

From the overlay it can be seen also, that both rays do not meet at the same time, since wave fronts do not have equal diameters.

Now the task will be done to show what happens if the middling observation angle is  $90^\circ$ , i.e. the object's / source's position is on the zenith. Only the final overlay is being shown, again the ray turns out to be defined by consecutive wave front normals:




Fig. 13: Overlay 0,6 and 0,4 km/sec but on  $90^\circ$  middling observation angle

Obviously even with the small relation of values for  $c$  and  $v$ , the diameters of both wave fronts are very close and no more visible on this scale image.

Now it might be also interesting, how the concept behaves when the light source is moving:




Fig. 14: Light propagation from source speed 0,6 km/sec with 0,3 sec steps, earth's speed 0,6 km/sec

It becomes clear that the aberration still exists the same way as if the source was not moving. The observer still receives only wave normals, but in this case from ever different source's position. The Doppler effect [11] behaves strictly in the classical way.

The determination of the aberration angle is done geometrically upon the aforementioned scale model. First the angle between observer at time of emission and the source will be calculated:




Fig. 15: Geometric model of 60° observation angle at mean speed

$$(1) \tan(\alpha) = \frac{h}{s+x}$$

$$(2) \tan(\beta) = \frac{h}{x} \Rightarrow h = \tan(\beta) \cdot x$$

$$(3) \cos(\beta) = \frac{x}{l} \Rightarrow x = \cos(\beta) \cdot c$$

Now insert (3) in (2)

$$(4) h = \tan(\beta) \cdot \cos(\beta) \cdot c = \sin(\beta) \cdot c$$

Now insert (4) and (3) in one:

$$\tan(\alpha) = \frac{\tan(\beta) \cdot \cos(\beta) \cdot c}{s + \cos(\beta) \cdot c} \Rightarrow \tan(\alpha) = \frac{\sin(\beta) \cdot l}{s + \cos(\beta) \cdot l} \Rightarrow \tan(\alpha) = \frac{\sin(\beta)}{\frac{s}{l} + \cos(\beta)} \Rightarrow \boxed{\tan(\alpha) = \frac{\sin(\beta)}{\frac{v}{c} + \cos(\beta)}}$$

Now it is important to acknowledge that the angle of aberration at maximum speed against the mean speed is different from the angle at minimum speed against mean speed, i.e. the full aberration angle is not simply double of one of the angles. First we calculate one of the angles:




Fig. 16: Geometric model of 60° observation angle at maximum speed

$$(1) \sin(\gamma_1) = \frac{s_1}{l_1} \cdot \sin(\alpha) = \frac{v_1}{c} \cdot \sin(\alpha)$$

$$(2) \tan(\alpha) = \frac{\sin(\beta)}{v/c + \cos(\beta)} \Rightarrow \alpha = \arctan\left(\frac{\sin(\beta)}{v/c + \cos(\beta)}\right) \text{ (from above)}$$

Now insert (2) in (1):

arbitrary observation angle  $\beta$  at speed  $v_1$

$$\sin(\gamma_1) = \frac{v_1}{c} \cdot \sin\left(\arctan\left(\frac{\sin(\beta)}{v/c + \cos(\beta)}\right)\right)$$

arbitrary observation angle  $\beta$  at speed  $v_2$

$$\sin(\gamma_2) = \frac{v_2}{c} \cdot \sin\left(\arctan\left(\frac{\sin(\beta)}{v/c + \cos(\beta)}\right)\right)$$

For 90° observation angle  $\sin(\beta)=1$  und  $\cos(\beta)=0$

$$\sin(\gamma_1) = \frac{v_1}{c} \cdot \sin\left(\arctan\left(\frac{c}{v}\right)\right)$$

$$\sin(\gamma_1) = \frac{v_1}{c} \cdot \frac{\frac{c}{v}}{\sqrt{1+c^2/v^2}} = \frac{v_1}{c} \cdot \frac{c}{\sqrt{1+c^2/v^2}}$$

90° observation angle at speed  $v_1$

$$\sin(\gamma_1) = \frac{v_1}{v} \cdot \frac{1}{\sqrt{1+c^2/v^2}}$$

90° observation angle at speed  $v_2$

$$\sin(\gamma_2) = \frac{v_2}{v} \cdot \frac{1}{\sqrt{1+c^2/v^2}}$$

And the full aberration angle is the difference of the above angles. Any similarity to the Lorentzian length contraction factor  $\gamma$  [2] is coincidence but might also be of relevance for further investigations.

On the basis of above formulae it is convenient to produce an excel sheet to play with different speeds, distances and angles. The following values were set:

Light speed c: 299.792 km/sec

Earth movement speed v: 300 km/sec +/- 29,78 km/sec on orbital path

Distance of earth path to light source: 20.000 lightyears

Middling angle of observation at 300 km/sec towards source: 90° degree

|  | higher speed     | mean speed         | lower speed      |   |                                       |
|--|------------------|--------------------|------------------|---|---------------------------------------|
|  | km/s, Grad       | km/s, Grad         | km/s, Grad       |   |                                       |
| c  | 299.792,00000    | 299.792,00000      | 299.792,00000    |   |                                       |
| v Earth 368 km/s (+/- 29,78 km/s on orbit)       | 397,78000        | 368,00000000000000 | 338,22000        |   |                                       |
| Incident angle light ray degree                  | 90,00569         | 90,00000           | 89,99431         | 0,011383014335 Difference low/high values is aberration angle x 2 |                                       |
|  | Radian           | 1,57090            | 1,57080          | 1,57070   | 0,005691507167320 aberration angle    |
|  |                  |                    |                  |   | 20,4894258024 Aberration angle arcsec |
| Distance earth path to source km                 | 1,8908481E+20    | 1,8908481E+20      | 1,8908481E+20    |   |                                       |
| lightyears                                       |                  | 20.000.000,00000   |                  |   |                                       |
|  |                  |                    |                  |   |                                       |
| angle difference observer-star to incident angle | 0,07602305829156 | 0,07033155077724   | 0,06464004395692 | 0,0113830143346   |                                       |
|  |                  |                    |                  |   |                                       |
| Angle observer to star at emission degree        | 89,92967         | 89,92967           | 89,92967         |   |                                       |
|  | Radian           | 1,56957            | 1,56957          | 1,56957   |                                       |
|  |                  |                    |                  |   |                                       |
| pathlength earth                                 | 2,508878E+17     | 2,321050E+17       | 2,133221E+17     | 3,756568E+16 Difference low/high values                           |                                       |
| pathlength light                                 | 1,890848E+20     | 1,890848E+20       | 1,890848E+20     | 3,080192E+06 Difference low/high values                           |                                       |
| part   |                  |                    |                  | 1,629000E-14  |                                       |

Fig. 17: Calculation sheet with realistic values

The resulting aberration angle is 20,4894", properly matching the observations. Interestingly there is still a time lack between both wave front spheres of approx. one part of a trillion at 90°, amounting to a distance deviation of approx. 35.000 km in this case that could be responsible for observed irregularities of planet's orbits. The deviation is progressively increasing on flat observation angles. **On the scale of mercury, observed under 60° the deviation would be approx. 10.000 km, well explaining the anomaly of Mercury orbit deviation:**

|  | higher speed      | mean speed        | lower speed       |  |                                  |
|--|-------------------|-------------------|-------------------|--|----------------------------------|
|  | km/s, Grad        | km/s, Grad        | km/s, Grad        |  |                                  |
| c  | 299.792,00000     | 299.792,00000     | 299.792,00000     |  |                                  |
| v Earth 368 km/s (+/- 29,78 km/s on orbit)       | 397,78000         | 368,00000         | 338,22000         |  |                                  |
| Incident angle light ray degree                  | 60,00493          | 60,00000          | 59,99507          | 0,00985 Difference low/high values is aberration angle x 2 |                                  |
|  | Radian            | 1,04728           | 1,04720           | 1,04711  | 0,00493 aberration angle         |
|  |                   |                   |                   |  | 17,73348 Aberration angle arcsec |
| Distance earth path to source km                 | 92.000.000,00000  | 92.000.000,00000  | 92.000.000,00000  |  |                                  |
| lightyears                                       |                   | 0,00001           |                   |  |                                  |
|  |                   |                   |                   |  |                                  |
| angle difference observer-star to incident angle | 0,06580           | 0,06087           | 0,05595           |  |                                  |
|  |                   |                   |                   |  |                                  |
| Angle observer to star at emission degree        | 59,93913          | 59,93913          | 59,93913          |  |                                  |
|  | Radian            | 1,04614           | 1,04614           | 1,04614  |                                  |
|  |                   |                   |                   |  |                                  |
| pathlength earth                                 | 140.947,87888     | 130.402,21696     | 119.855,50904     | 21.092,36983 Difference low/high values                    |                                  |
| pathlength light                                 | 106.227.177,09255 | 106.232.449,53094 | 106.237.723,27750 | -10.546,18495 Difference low/high values                   |                                  |
| part   |                   |                   |                   | -0,0000992697  |                                  |

Fig. 18: Results on mercury perihelion shift

Now for checkup the distance is set to 1.000 km and again 90°, resulting in again 20,4894''

|  | higher speed         | mean speed         | lower speed          |   |
|--|----------------------|--------------------|----------------------|---|
|  | km/s, Grad           | km/s, Grad         | km/s, Grad           |   |
| c  | 299.792,00000        | 299.792,00000      | 299.792,00000        |   |
| v Earth 368 km/s (+/- 29,78 km/s on orbit)       | 397.78000            | 368.0000000000000  | 338.22000            |   |
| Incident angle light ray degree                  | 90,00569             | 90,00000           | 89,99431             | 0,011383014335 Difference low/high values is aberration angle x 2 |
| Radian   | 1,57090              | 1,57080            | 1,57070              | 0,005691507167320 aberration angle                                |
|  |                      |                    |                      | 20,4894258024 Aberration angle arcsec                             |
| Distance earth path to source km                 | 1,0000000E+03        | 1,0000000E+03      | 1,0000000E+03        |   |
| lightyears                                       |                      | 0,00000            |                      |   |
|  |                      |                    |                      |   |
| angle difference observer-star to incident angle | 0,07602305829156     | 0,07033155077724   | 0,06464004395692     | 0,0113830143346   |
| Angle observer to star at emission degree        | 89,92967             | 89,92967           | 89,92967             |   |
| Radian   | 1,56957              | 1,56957            | 1,56957              |   |
|  |                      |                    |                      |   |
| pathlength earth                                 | 1,32685329149042     | 1,22751774563691   | 1,12818221189572     | 0,1986710795946920000 Difference low/high values                  |
| pathlength light                                 | 1.000,00000493362000 | 999,99999999995000 | 1.000,00000493360000 | 0,000000000162572178 Difference low/high values                   |
| part   |                      |                    |                      | 0,0000000000000162572   |

Fig. 19: Calculation sheet with realistic values but unrealistic short distance

Obviously distance is irrelevant for the aberration angle, as it should be.

The fact that it has been herewith proved that the aberration is resulting from the whole distance between source and observer alone, makes it almost irrelevant if the light ray is entrained on the short piece in close distance of source or observer. Since the influencing distance is vanishingly short against the distance between observer and source, the aberration must have already happened on its way. Also experiments with water filled telescopes (by George Bidell Airy, [12]) or the like therefore cannot but have a null result.

The same principle applies for the source. As for any wave, movement of source is irrelevant for the wave front that was emitted at one time. If emitted waves were dragged by the source star, the influence would be again vanishingly because of the comparably very short distance that light might be dragged by gravity of source. **The reverse argument though is, that light entrained by gravity still causes stellar aberration and entrained ether remains fully suitable to explain stellar aberration.**

As well the lack of any observable terrestrial aberration is self- explanatory based on the gravity entrained ether concept. Though light speed does not depend on source's movement, light will be entrained by gravity and the ether will not have any relative speed against the observer. Therefore no differing light speed will be measurable by the observer on earth, nor terrestrial aberration can exist.

#### 4. More empiric evidence esteemed to disprove entrained ether

The Hammar [13] experiment with a setup consisting of differing length interferometer arms partially cladded with heavy lead blocks also gave a null result, although, under the terms of entrained ether, a positive result was expected due to gravitational attraction of light by the blocks. The obtained null result is everything but significant. It is completely implausible why any lightray that is already fully entrained by gravity, should be even more than fully entrained by additional gravity. Also arguments of the sort that mass and gravity of the rotating disc of Sagnac type experiments could influence the light propagation are irrelevant for the same reason, and additionally gravity is confused with centrifugal force, as if gravity would be somehow rotating together with the disc's motion. Setups with glass fibre laser gyroscopes can brilliantly show the Sagnac effect, though both cannot reveal any second order effect that would be overrun by the first order effect. GPS technology is functioning only with earth as the inertial reference frame, calculations solely use the



Sagnac effect and relativistic second order effects play no role. If sun was used as a reference frame, computations based upon Relativity fail to produce correct results, thus favoring the gravity entraining ether concept outlined in this paper.

## 5. Conclusion and Perspective

We have seen that understanding of the nature of light propagation is until today underlying some fundamental misinterpretations that we brought into order with this paper. In fact there is no reason why light should not be entrained by gravity in general, at least on the foundation of the above discussed experiments. Further investigations will have to be accomplished on experiments dealing with laser resonator setups and frequency comparison and have to be revised under terms of gravity entrained ether.

References and Acknowledgments (in order of appearance):

- [1] Einstein, Albert, Zur Elektrodynamik bewegter Körper, Annalen der Physik und Chemie. 17, 1905, S. 891–921, 1905.
- [2] H. A. Lorentz, Versuch einer Theorie der electrischen und optischen Erscheinungen in bewegten Körpern. E.J. Brill, Leiden 1895..
- [3] Michelson, A. A., Morley, E.W, On the Relative Motion of the Earth and the Luminiferous Ether, Bde. 1 von 2, 1887, 34 (203): 333–345 1881, American Journal of Science, 1887.
- [4] G. Sagnac, „L'éther lumineux démontré par l'effet du vent relatif d'éther dans un interféromètre en rotation uniforme.,“ In: Comptes Rendus. 157, S. 708–710., 1913.
- [5] A. A. Michelson und H. G. Gale, „The Effect of the Earth's Rotation on the Velocity of Light, II,“ Astrophysical Journal. 61: 140. Bibcode:1925ApJ....61..140M. doi:10.1086/142879., 1925.
- [6] Fresnel, Augustin, Lettre d'Augustin Fresnel à François Arago sur l'influence du mouvement terrestre dans quelques phénomènes d'optique, Bde. 1 von 2 Oeuvres complètes d'Augustin Fresnel / publiées par MM. Henri de Senarmont, <http://visualiseur.bnf.fr/CadresFenetre?O=NUMM-91937&l=633&M=tdm>, Gallica, 1818.
- [7] Stokes, George, mathematical and physical papers, Vol.1, Cambridge University Press, 1880.
- [8] F. M. Schmitt, „On Aberration of Light and Reflection from Moving Mirrors,“ viXra:1303.0110, Berlin, 2013.
- [9] A. Kelly, Challenging Modern Physics, Boca Raton: Brown Walker Press, 2005.
- [10] Bradley, James, A Letter from the Reverend Mr. James Bradley Savilian Professor of Astronomy at Oxford, and F.R.S. to Dr. Edmond Halley Astronom. Reg. &c. Giving an Account of a New

Discovered Motion of the Fix'd Stars, Bde. %1 von %2Philosophical Transactions 35, 1727-1728, doi:10.1098/rstl.1727.0064, The Royal Society, 1727.

[11] Doppler, Christian, Über das farbige Licht der Doppelsterne und einiger anderer Gestirne des Himmels, Böhmische Gesellschaft der Wissenschaften, 1842.

[12] Airy, George Bidell, On a Supposed Alteration in the Amount of Astronomical Aberration of Light, Produced by the Passage of the Light through a Considerable Thickness of Refracting Medium, Bde. %1 von %2Proceedings of the Royal Society of London, Volume 20, January 1, 1871 20 35-39; doi:10.1098/rspl.1871.0011, The Royal Society, 1871.

[13] G. W. Hammar, The Velocity of Light Within a Massive Enclosure, Physical Review, 1935.