

Planck Mass Measured Totally Independent of Big G Utilizing McCulloch-Heisenberg Newtonian Equivalent Gravity! Is Newton's Gravitational Constant a Composite Constant?

Espen Gaarder Haug*
Norwegian University of Life Sciences

March 31, 2018

Abstract

In 2014, McCulloch showed, in a new and interesting way, how to derive a gravity theory from Heisenberg's uncertainty principle that is equivalent to Newton gravity. McCulloch utilizes the Planck mass in his derivation and gets a gravitational constant of $\frac{\hbar c}{m_p^2}$. This is a composite constant, which is equivalent in value to Newton's gravitational constant. However, McCulloch has pointed that his approach requires an assumption on the value of G , and that this involves some circular reasoning. This is in line with the view that the Planck mass is a derived constant from Newton's gravitational constant, while big G is a universal fundamental constant. Here we will show that we can go straight from the McCulloch derivation to measuring the Planck mass without any knowledge of the gravitational constant. Therefore, from this perspective, there are no circular problems with his method. This means that we can measure the Planck mass totally independent of Newton's gravitational constant, and shows that the McCulloch derivation is a theory of quantum gravity that stands on its own, with no need for input from Newton. This also strongly points towards the idea that Newton's gravitational constant should be seen as a composite constant. Based on this perspective, for the first time since it was theoretically introduced 119 years ago, the Planck units can be measured completely independent of any knowledge of Newton's gravitational constant. This could be an important step towards the development of a full theory of quantum gravity.

Key words: Heisenberg, Planck mass, McCulloch gravity, Newton, gravitational constant, Cavendish apparatus.

1 McCulloch-Heisenberg Newton Equivalent Gravity

In 2014, McCulloch [1] derived an equivalent gravity to that of Newton [2] directly from the Heisenberg uncertainty and gets the following equation for the gravity force (See Appendix A for a short review of his derivation.)

$$F = \frac{\hbar c}{m_p^2} \frac{Mm}{r^2} \quad (1)$$

Where $\frac{\hbar c}{m_p^2} \approx 6.67384 \times 10^{-11} m^3 \cdot kg^{-1} \cdot s^{-2}$. That is basically identical to the empirically-measured gravitational constant value, even if there is large uncertainty in the Newtonian constant [3, 4, 5, 6, 7]; this is something we will get back to later. Still, we cannot know its value without knowing the speed of light, the Planck constant, and the Planck mass. The speed of light is known and can be measured with no knowledge of gravity (and is exact per definition), the Planck constant can be found from the Watt balance [8, 9]. However, the Planck mass is unknown and it is generally assumed that we must know G in order to calculate the Planck mass. This point is mentioned by McCulloch himself in a follow-up paper [10]. In that paper McCulloch states

In the above gravitational derivation, the correct value for the gravitational constant G can only be obtained when it is assumed that the gravitational interaction occurs between whole multiples of the Planck mass, but this last part of the derivation involves some circular reasoning since the Planck mass is defined using the value for G .

*e-mail espenhaug@mac.com. Thanks to Victoria Terces for helping me edit this manuscript.

This is fully in line with modern physics' view that the only way to find the Planck mass is to derive it from big G . The Planck mass, the Planck temperature (energy), the Planck length, and the Planck time were introduced in 1899 by Max Planck [11, 12] himself. Planck derived these units, which he called natural units, from what he considered to be the most fundamental universal constants: Newton's gravitational constant, the speed of light, and the Planck constant. Based on this, we need to know the traditional constant to find the Planck mass from Planck's formula, $m_p = \sqrt{\frac{\hbar c}{G}}$. However, as we will see here, we can build on the McCulloch derivation, complete a few more derivations, and easily design a simple experiment to measure the Planck mass independent of Newton's gravitational constant, or knowledge of any other theories of gravity.

2 The Planck Mass Measured Directly from McCulloch's Derivation and a Cavendish Apparatus

Newton did not measure the gravitational constant himself; this was first done indirectly by Cavendish [13] in 1798. Using a Cavendish apparatus, we can measure the Planck mass without any knowledge of Newton's gravitational constant, or any knowledge of Newtonian gravity. A Cavendish apparatus consist of two small balls and two larger balls, all made of lead, for example. The torque (moment of force) is given by

$$\kappa\theta \tag{2}$$

where κ is the torsion coefficient of the suspending wire and θ is the deflection angle of the balance. We then have the following well-known relationship

$$\kappa\theta = LF \tag{3}$$

where L is the length between the two small balls in the apparatus. Further, F can be set equal to the gravitational force given by McCulloch's Heisenberg-derived formula

$$F = \frac{\hbar c}{m_p^2} \frac{Mm}{r^2} \tag{4}$$

This means we must have

$$\kappa\tau = L \frac{\hbar c}{m_p^2} \frac{Mm}{r^2} \tag{5}$$

We also have that the natural resonant oscillation period of a torsion balance is given by

$$T = 2\pi\sqrt{\frac{I}{\kappa}} \tag{6}$$

Further, the moment of inertia I of the balance is given by

$$I = m \left(\frac{L}{2}\right)^2 + m \left(\frac{L}{2}\right)^2 = 2m \left(\frac{L}{2}\right)^2 \tag{7}$$

this means we have

$$T = 2\pi\sqrt{\frac{mL^2}{2\kappa}} \tag{8}$$

and when solved with respect to κ , this gives

$$\begin{aligned} \frac{T^2}{2^2\pi^2} &= \frac{mL^2}{2\kappa} \\ \kappa &= \frac{mL^2}{2\frac{T^2}{2^2\pi^2}} \\ \kappa &= \frac{mL^2 2\pi^2}{T^2} \end{aligned} \tag{9}$$

Next in equation 5 we are replacing κ with this expression, and solving with respect to the Planck mass

$$\begin{aligned}
\frac{mL^2 2\pi^2}{T^2} \theta &= L \frac{\hbar c}{m_p^2} \frac{Mm}{r^2} \\
\frac{L^2 2\pi^2 r^2}{\hbar c L M T^2} \theta &= \frac{1}{m_p^2} \\
m_p^2 &= \frac{\hbar c M T^2}{L 2\pi^2 r^2 \theta} \\
m_p &= \sqrt{\frac{\hbar c M T^2}{L 2\pi^2 r^2 \theta}}
\end{aligned} \tag{10}$$

The mass M is the mass of each of the two large lead balls in the Cavendish apparatus, not the mass of the Earth. All we need in order to find the mass of the large balls is an accurate weight. The Planck constant can be found from the Watt balance. The angle θ and the time T is what we measure with the Cavendish apparatus. The length L is the distance between the small lead balls and r is the distance between the large lead ball's center to the center of the small lead ball, when the arm is in equilibrium position (mid position).

Today there even exists a small ready-to-use low budget (a few thousand dollars) Cavendish apparatus, where the angle of the arm (and the time) are measured very accurately by fine electronics and plugged directly into a computer with a USB cable. Even by this low budget apparatus we can measure the Planck mass with about 5% inaccuracy from our kitchen table without any knowledge of Newton's gravitational constant.

As soon as we know the Planck mass, we have the the complete composite gravitational constant and the McCulloch formula can then be applied to any standard gravitational predictions, such as finding the mass of the Earth, or predicting the orbital velocity of planets and satellites.

Haug [14] has, in a similar way, shown how the Planck length can be found independent of big G , but his derivation then did not start out with the McCulloch-Heisenberg Newton equivalent gravity theory. Thus, it was not clear that the starting point was not totally independent on already knowledge of Newton gravity, so we think this derivation and discussion in this paper offers additional insight.

3 Why Newton's Gravitational Constant Likely Is a Universal Composite Constant

In our analysis, the first strong indication that the gravitational constant is a composite constant is given by its output units, which are $m^3 \cdot kg^{-1} \cdot s^{-2}$. It would be very strange if something concerning the fundamental nature of reality would be meters cubed, divided by kg and seconds squared. The Planck mass, on the other hand, is somewhat easier to relate to, even if it is somewhat of a mystery at a deeper level. The speed of light is also something we can relate to logically; it is the distance light travels in vacuum during a pre-specified time interval. The Planck constant is more complex (and outside the scope of this paper), but it is related to the view that energy seems to come in quanta. In sum though, the Planck mass, the speed of light, and the Planck constant seem to be more intuitive than the gravitational constant.

In 2016, Haug [15] independently suggested that the gravitational constant of the form $G = \frac{l_p^2 c^3}{\hbar}$, which is basically the same as the McCulloch 2014 constant as the Planck mass, can be written as $m_p = \frac{\hbar}{l_p} \frac{1}{c}$, and we then must have $\frac{\hbar c}{m_p^2} = \frac{l_p^2 c^3}{\hbar}$. Haug has shown that assuming the gravitational constant is a composite will help make all the Planck units more intuitive. For example, the Planck time is given by $t_p = \sqrt{\frac{G\hbar}{c^5}}$; when rewritten based on the idea that the gravitational constant is a composite, this simply gives the (known) $t_p = \frac{l_p}{c}$. The later form is also known from before, but the view that the Newtonian gravitational constant is a composite renders the form $t_p = \sqrt{\frac{G\hbar}{c^5}}$ unnecessary. We might then ask, what is the intuition about c^5 and G ? The answer may not be so clear. On the other hand, the intuition behind $\frac{l_p}{c}$ is very simple; it is simply a very short distance divided by the speed of light, so given a very short time interval, we can see it is coming directly out from the formula. Haug's gravitational constant composit formula has the same challenge in that it seems like we may end up with a circular problem, again, because modern physics typically assumes that we need to know big G before we can find the Planck units. However, as we have shown in this paper this is not the case.

This does not mean big G is wrong; it is just likely to be a composite universal constant rather than a fundamental constant.

We find that many gravitational formulas may be seen in a new perspective when rewritten based on the idea that Newton's gravitational constant is a composite constant; we summarize a selection of such gravitational formulas in Table 1.

	Standard form/way	Planck form	Observed
Gravitational constant	$G \approx 6.67 \times 10^{-11}$	$G = \frac{\hbar c}{m_p^2} = \frac{l_p^2 c^3}{\hbar} \approx 6.67 \times 10^{-11}$	Yes
Gravitational constant	$G = \sqrt{\frac{c^3 2 l_p^2 \pi^2 L r^2 \theta}{\hbar M T^2}}$		Yes
Cavendish Planck mass	Only derived from G	$m_p = \sqrt{\frac{\hbar^3 2 \pi^2 L r^2 \theta}{l_p^2 M T^2 c^5}}$	Easy to do
Newton gravity force	$F = G \frac{mM}{r^2}$	$F = n_1 n_2 \frac{\hbar c}{r^2}$	"Yes" ^a
Gravitational acceleration field	$g = \frac{GM}{r^2}$	$g = N \frac{l_p}{r^2} c^2$	Yes
Mass from acceleration field	$g = \frac{GM}{r^2}$	$M = \frac{gr^2 \hbar}{l_p^2 c^3} = \frac{gr^2 m_p}{l_p c^2}$	Yes
Orbital velocity	$v_o = \sqrt{G \frac{M}{r}}$	$v_o = c \sqrt{N \frac{l_p}{r}}$	Yes
Escape velocity	$v_e = \sqrt{2G \frac{M}{r}}$	$v_e = c \sqrt{2N \frac{l_p}{r}}$	No (?) ^b
Time dilation	$t_2 = t_1 \sqrt{1 - \frac{2GM}{rc^2}}$	$t_2 = t_1 \sqrt{1 - 2N \frac{l_p}{r}}$	Yes
Newton gravitational bending of light	$\delta = \frac{2GM}{rc^2}$	$\delta = 2N \frac{l_p}{r}$	Twice of that
GR gravitational bending of light	$\delta = \frac{4GM}{rc^2}$	$\delta = 4N \frac{l_p}{r}$	Yes
Gravitational red-shift	$\lim_{r \rightarrow +\infty} z(r) = \frac{GM}{r^2}$	$\lim_{r \rightarrow +\infty} z(r) = N \frac{l_p}{r}$	Yes
Schwarzschild radius	$r_s = \frac{2GM}{c^2}$	$r_s = 2N l_p$	No
Einstein field equation	$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \frac{8\pi G}{c^4} T_{\mu\nu}$	$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \frac{8\pi l_p}{m_p c^2} T_{\mu\nu}$	"Yes"
Einstein constant	$\kappa = \frac{8\pi G}{c^2}$	$\kappa = \frac{8\pi l_p}{m_p}$	Yes
Einstein cosmological constant	$\Lambda = \kappa \rho_{vac}$	$\Lambda = \frac{8\pi l_p}{m_p} \rho_{vac}$	Yes
Hawking temperature	$\frac{c^3 \hbar}{8\pi G M k_b}$	$T = \frac{1}{N 8\pi} \frac{m_p c^2}{k_b}$	No ^c
Hawking dissipation time	$t_{ev} = \frac{15360\pi G^2 M^3}{\hbar c^4}$	$T = 15360\pi \frac{l_p}{c}$	No ^d
Bekenstein-Hawking luminosity	$P = \frac{\hbar c^6}{15360\pi G^2 M^2}$	$P = \frac{1}{N^2 15360} \frac{\hbar c^2}{l_p^2}$	No ^e
McCulloch orbital mass	$M = \frac{v^4 \Theta}{2Gc^2(1+Z)}$	$M = \frac{1}{2(1+Z)} m_p \frac{\Theta v^4}{l_p c^4}$	(?) ^f
McCulloch galaxy velocity	$v^4 = \frac{2GMc^2(1+Z)}{\Theta}$	$v = c \left(2N \frac{l_p}{\Theta} (1+Z) \right)^{\frac{1}{4}}$	(?) ^g

Table 1: The table of a series gravity formulas when using the standard Newton gravitational constant and the alternative when arguing that Newton's gravitational constant is a composite constant.

^aActually the Newton gravitational force has never been observed directly, only indirectly though the predictions that comes from mathematically rearranging this formula to develop other predictions, such as orbital velocity.

^bTo my knowledge the escape velocity has not been tested empirically.

^cAt least not directly.

^dAt least not directly.

^eAt least not directly.

^fNeeds further investigation and confirmation, see [16] for more details.

^gNeeds further investigation and confirmation, see [16] for more details.

4 Relative Standard Uncertainty in the Planck Mass versus Newton's Gravitational Constant

Assume we have measured the Planck mass (with a standard uncertainty of 1%) on the kitchen table with Cavendish apparatus plugged into our computer. The standard uncertainty in the gravitational constant must then be

$$\% \text{ Standard relative uncertainty} = \frac{\partial G}{\partial m_p} \times \frac{m_p}{G} = -\frac{2\hbar c}{m_p^3} \times \frac{m_p}{G100} = \frac{1}{50} = 2\% \quad (11)$$

That is to say, the standard uncertainty in the Newton gravitational constant will always be twice that of the standard uncertainty in the Planck mass measurements. This is in line with what is reported by NIST (2014) CODATA, which reports a relative standard uncertainty for the gravitational constant of 4.7×10^{-5} and 2.3×10^{-5} for the Planck mass.

5 Conclusion

We have shown that the Planck mass can be measured with a Cavendish apparatus without any prior knowledge of gravity except the McCulloch gravity derived directly from Heisenberg's uncertainty principle. This no longer posits the Planck mass as simply being a derived constant from big G , but possibly even makes it more important than big G , since the gravitational constant can be written as a composite constant $G = \frac{\hbar c}{m_p}$. This indicates that the Planck units are the truly fundamental units and that the gravitational constant likely is a composite constant. This also indicates that the Planck units plays a very central role in gravity. After years of thinking about the problem, we have come up with this gravity experiment to measure the Planck mass, the Planck length, and Planck time. It is quite remarkable it has taken us about 119 years since Max Planck first suggested the natural Planck units to discovering a way to measure them totally independent of any knowledge of Newton's gravitational constant. This could be an important step toward a theory of quantum gravity that unites the quantum scale and the cosmological scale.

Appendix A: Newton's Gravity from Heisenberg's Uncertainty Principle

In 2014, McCulloch derived Newton's gravitational force from Heisenberg's uncertainty principle, for a more in detailed derivation see the McCulloch papers [1, 10]. Heisenberg's uncertainty principle [17] is given by

$$\Delta p \Delta x \geq \hbar \quad (12)$$

McCulloch goes on to say "Now $E = pc$ so" :

$$\Delta E \Delta x \geq \hbar c \quad (13)$$

This assumption only holds for the Planck momentum $E = pc = m_p c c$. It is implied indirectly in the McCulloch derivation that the Planck mass somehow plays an essential role in gravity. Further, from equation 13, McCulloch goes on to suggest that

$$F = \frac{1}{(\Delta x)^2} \sum_i^n \sum_j^N (\hbar c)_{i,j} \quad (14)$$

where \sum_i^n is the number of Planck masses in a smaller mass m we are working with, \sum_j^N corresponds to the number of Planck masses in the larger mass we are working with. McCulloch from this get the equation

$$F = \frac{\hbar c}{m_p^2} \frac{mM}{(\Delta x)^2} \quad (15)$$

McCulloch also replaces Δx with the radius, something we think is sound, based on extensive analysis. Further, he correctly points out that

$$G = \frac{\hbar c}{m_p^2} \quad (16)$$

which basically means his derivation is equivalent to the Newtonian gravity formula

$$F = G \frac{mM}{r^2} \quad (17)$$

Appendix B: The Planck Time and the Planck Length Directly from McCulloch-Heisenberg Equivalent Newton Gravity

We can also find the Planck time directly from McCulloch-Heisenberg Newton equivalent gravity using a Cavendish experiment by utilizing the derivation below

$$\begin{aligned} \frac{mL^2 2\pi^2}{T^2} \theta &= L \frac{\hbar c}{m_p^2} \frac{Mm}{r^2} \\ \frac{mL^2 2\pi^2}{T^2} \theta &= L \frac{t_p c^5}{\hbar} \frac{Mm}{r^2} \\ t_p^2 &= \frac{\hbar L^2 2\pi^2 r^2}{LMT^2 c^5} \theta \\ t_p &= \sqrt{\frac{\hbar L 2\pi^2 r^2 \theta}{MT^2 c^5}} \end{aligned} \quad (18)$$

Similarly, we can also find the Planck length directly from the McCulloch-Heisenberg Newton equivalent gravity, taking into account that an elementary particle can be written as

$$m = \frac{\hbar}{\lambda} \frac{1}{c} \quad (19)$$

In this case, we know the mass is the Planck mass, so the reduced Compton wavelength is related to the Planck length that we can find directly using a Cavendish apparatus

$$\begin{aligned} \frac{mL^2 2\pi^2}{T^2} \theta &= L \frac{\hbar c}{m_p^2} \frac{Mm}{r^2} \\ l_p^2 &= \frac{\hbar L^2 2\pi^2 r^2 \theta}{LMT^2 c^3} \\ l_p &= \sqrt{\frac{\hbar L 2\pi^2 r^2 \theta}{MT^2 c^3}} \end{aligned} \quad (20)$$

In other words, all of the natural Planck units can be found directly from a quantum-derived Newtonian equivalent gravity theory.

References

- [1] M. E. McCulloch. Gravity from the uncertainty principle. *Astrophysics and Space Science*, 349(2), 2014.
- [2] I. Newton. *Philosophiae Naturalis Principia Mathematica*. London, 1686.
- [3] G. S. Bisnovaty-Kogan. Checking the variability of the gravitational constant with binary pulsars. *International Journal of Modern Physics D*, 15(07), 2006.
- [4] B. Fixler, G. T. Foster, J. M. McGuirk, and M. A. Kasevich. Atom interferometer measurement of the Newtonian constant of gravity. *Science*, 315, 2007.
- [5] S. Galli, A. Melchiorri, G. F. Smoot, and O. Zahn. From Cavendish to Planck: Constraining Newton's gravitational constant with CMB temperature and polarization anisotropy. *Physical Review D*, 80, 2009.
- [6] G. Rosi, F. Sorrentino, L. Cacciapuoti, M. Prevedelli, and G. M. Tino. Precision measurement of the Newtonian gravitational constant using cold atoms. *Nature*, 510, 2014.

- [7] F. Scholkmann. Indications for a diurnal and annual variation in the anisotropy of diffusion patterns: A reanalysis of data presented by J. Dai (2014, nat. sci.). *Progress in Physics*, 10, 2014.
- [8] M. Stock. The Watt balance: determination of the Planck constant and redefinition of the kilogram. *Philosophical Transactions of the Royal Society*, 369:3936–3953, 2011.
- [9] I. A. Robinson and S. Schlamminger. First determination of the Planck constant using the LNE Watt balance. *Forthcoming, Metrologia*, 2016.
- [10] M. E. McCulloch. Quantised inertia from relativity and the uncertainty principle. *EPL (Europhysics Letters)*, 115(6), 2016.
- [11] M. Planck. *Naturlische Masseinheiten*. Der Königlich Preussischen Akademie Der Wissenschaften, p . 479., 1899.
- [12] M. Planck. *Vorlesungen über die Theorie der Wärmestrahlung*. Leipzig: J.A. Barth, p. 163, see also the English translation “The Theory of Radiation” (1959) Dover, 1906.
- [13] H. Cavendish. Experiments to determine the density of the Earth. *Philosophical Transactions of the Royal Society of London, (part II)*, 88, 1798.
- [14] E. G. Haug. Can the Planck length be found independent of big G ? *Applied Physics Research*, 9(6), 2017.
- [15] E. G. Haug. The gravitational constant and the Planck units. A simplification of the quantum realm. *Physics Essays Vol 29, No 4*, 2016.
- [16] M. E. McCulloch. Galaxy rotations from quantised inertia and visible matter only. *Astrophysics and Space Science*, 362, 2017.
- [17] W. Heisenberg. Über den anschaulichen inhalt der quantentheoretischen kinematik und mechanik. *Zeitschrift für Physik*, (43):172–198, 1927.