

Duality transform between black and white psychological profiles.

Johan Noldus*

April 22, 2018

Abstract

It is shown that the Fourier transformation is the appropriate defining characteristic of black and white polarization states in psychological archetypes.

1 Introduction.

In previous work, I have described the necessity for at least a black-white theory of psychological profiles. In principle, psychological archetypes are described by means of a real vectorspace $V = \mathbb{R}^n$ given that no obvious extremal states need to a priori exist. Such assumption would lead to a description by means of convex spaces. Black is an effective charge described by a delta peak distribution on V whereas a pure white state is described by its Fourier transform \mathcal{F} . They are extremal weak distributional states in $L^2(V, \mu)$ the Hilbertspace of square integrable functions on V with respect to the measure μ . Circularly polarized states are then defined as “black content equals white content” which is the space of eigenstates of the Fourier transform.’

2 Elaboration.

In what follows, we take the pure theory and assume $n = 1$. V then is spanned by means of the black states given by $\delta(x - a)$, white states are of the form e^{ika} . As such, no information loss occurs and black-white are just different configurations of the same substance. Hence, given

$$g(x) = \int dyg(y)\delta(y - x)$$

we have that

$$(\mathcal{F}g)(k) = \int dyg(y)e^{iky}.$$

We now look for states for which $|g(x)|^2 \sim |(\mathcal{F}g)(x)|^2$ or, more precisely,

$$(\mathcal{F}g)(x) \sim e^{id_x}g(x)$$

*email: johan.noldus@gmail.com, Relativity group, departement of mathematical analysis, University of Gent, Belgium.

for some d . Taking into account that

$$(\mathcal{F}(g))(k) = \int dx e^{ikx} g(x)$$

the aforementioned class is given by the Gaussian functions

$$e^{-a(x-b)^2}$$

since

$$\int dx e^{ikx} e^{-a(x-b)^2} = \int dx e^{ikb} e^{-a((x-b) - \frac{ik}{2a})^2} e^{-\frac{k^2}{4a}} = \frac{\pi}{\sqrt{a}} e^{-ab^2} e^{-\frac{1}{4a}(k-2iab)^2}$$

and for this function to satisfy our criterion it is necessary and sufficient that $a = \frac{1}{2}$ and b is freely chosen. Hence, the diagonal in the white-black plane is one dimensional and parametrized by b just as the black line interval is. Taking $n > 1$ would entail a definition of whiteness given by $\delta(|x| - a)$ thereby suppressing $n - 1$ dimensions. Taking a to zero and scaling the Gaussian functions appropriately leads to the pure white states whereas taking a to infinity provides for the pure black states. The appropriate duality is therefore $a \rightarrow \frac{1}{4a}$. Hence one has to consider the operators

$$P_{d,\alpha} = e^{-ixd} \circ \mathcal{F} \circ S_\alpha$$

and consider eigenvectors $v_{d,\alpha}$ with the appropriate eigenvalue to be calculated above (hint: $\alpha = \frac{1}{\sqrt{2a}}$). All these functions are eigenvalues of the normal operator

$$T_{2b} \mathcal{F}^2$$

with eigenvalue $2\pi^2$. Here, e^{ixd} is the multiplication operator and S_α the scaling operator defined by

$$(S_\alpha g)(x) = g(\alpha x).$$

Generally, $d =$ There are some interesting commutation relations

$$e^{-ixd} \mathcal{F} = \mathcal{F} T_d$$

where

$$(T_d g)(x) = g(x + d)$$

and

$$(\mathcal{F} S_\alpha g)(x) = (S_{\frac{1}{\alpha}} \mathcal{F} g)(x).$$

Hence,

$$P_{d,\alpha}^\dagger P_{d,\alpha} \sim 1$$

given that e^{idx} is unitary and

$$S_\alpha^\dagger S_\alpha = \frac{1}{\alpha} 1.$$

Notice that

$$x \mathcal{F} x + \partial_x \mathcal{F} \partial_x = i \mathcal{F}^\dagger$$

and therefore, the Heisenberg algebra is equivalent to

$$X^\dagger X - P^\dagger P = i1$$

on inproduct spaces with a complex valued bilinear form gauged by $X^\dagger = P$. The Fourier transform is then recuperated by finding a unitary operator \mathcal{F} on a Hilbert space representation of X, P such that $X^\dagger = \mathcal{F} X^H \mathcal{F}$

with $X^H = X$. In general, the angle α between the mixed states and the “axis” of black extremal states should be given by a self dual function, that is

$$\frac{1}{f(x)} = f\left(\frac{1}{4x}\right)$$

giving rise to

$$f(x) = \frac{1}{\sqrt{2x}}.$$

Therefore,

$$\tan\left(\frac{\pi}{2} - \alpha\right) = f(x)$$

and the magnitude is determined by the shift parameter b . Henceforth, a compactification of the infinite dimensional function space to a real two dimensional Hilbert space with black-white extremal states is given by

$$|a, b\rangle = b(\cos(\alpha)|b\rangle + \sin(\alpha)|w\rangle).$$

There is no information loss but the correspondence is not a linear one for sure given that any linear combination on the right hand side determines a Gaussian but not so for linear combinations of the left hand side. It remains a task to find other suitable, higher, “pictorial” charges such as shapes, including tigers, cats, dogs and sharks, to further specify the psychological mapping to relative moral values.