

34 **2. Terms and definitions.**

35 **2.1. Boundary points and matrices.**

- 36 • Given a point P on $\partial\Delta$ (the boundary of Δ) and given a unitary matrix U
 37 such that $R(U)=P$, we call U a **boundary matrix** of Δ . See (1.3)
 38 • A **regular boundary point** is a point where the boundary is smooth.
 39 • A non-permutation boundary matrix for a regular boundary point is called a
 40 **regular boundary matrix**.

41 **2.2. Properties of unitary matrices given A_0 and B_0 .** In this section, we
 42 define four properties of unitary matrices that will be very useful when examining
 43 boundary matrices of Δ . These properties will be referred to throughout the paper
 44 in relation to a given unitary matrix U .

45 The first three of these properties are matrices related to U . These matrices are
 46 defined in [1], p.27. They provide a language to talk about unitary matrices within
 47 the context of the determinantal conjecture.

48 **B-matrix**

49
$$B = UB_0U^* \quad (2.1)$$

50 **C-matrix**

51
$$C = A_0 + UB_0U^* \quad (2.2)$$

52 Using (1.3), $R(U) = \det(C)$

53 **F-matrix**

54
$$F = BC^{-1} - C^{-1}B$$

55 We can change the F-matrix into a more useful form:

56
$$F = (C - A_0)C^{-1} - C^{-1}(C - A_0)$$

57
$$F = C^{-1}A_0 - A_0C^{-1} \quad (2.3)$$

59 The F-matrix is only defined when C is invertible or equivalently $R(U) \neq 0$.

60 Since A_0 is diagonal, we see that F is a zero-diagonal matrix.

61 As demonstrated in [1], p.27, the F-matrix is 0 if and only if U is a permutation
 62 matrix.

63 The fourth property is conditional. Given a unitary matrix U with $R(U) \neq 0$ and
 64 with F-matrix $F \neq 0$. let $T = tr(ZF)$, where Z is any skew-hermitian matrix. T is a
 65 complex number and can be seen as a vector in the complex plane. If for all possible
 66 skew-hermitian matrices Z , all values of T are either parallel or anti-parallel, then we
 67 say that U is **trace-argument constant**. We take the zero-vector as being parallel
 68 to any vector.

69 **2.3. Additional matrix related definitions.**

- 70 • An **essentially-hermitian** matrix is a matrix that can be written as $e^{i\theta}H +$
 71 λI where θ is real, H is hermitian, λ is complex and I is the identity matrix.
 72 Equivalently an essentially-hermitian matrix is a normal matrix with collinear
 73 eigenvalues. This definition comes from [5].

74 **3. Main Results.**

75 **THEOREM 3.1.** *Every regular boundary matrix U of Δ with $R(U) \neq 0$ is trace-*
 76 *argument constant.*

77 **THEOREM 3.2.** *$\partial\Delta$ is smooth at all non-zero, non-permutation points.*

78 **THEOREM 3.3.** *Given a unitary matrix that is trace-argument constant, its F -*
 79 *matrix is essentially-hermitian with $\lambda = 0$.*

80 **4. Proof of Theorem 3.1.** Our aim is to examine boundary matrices of Δ .
 81 Towards this aim, it is useful to consider smooth unitary matrix functions going
 82 through these boundary matrices and see how they behave under (1.3). For this
 83 reason, we introduce the functional form of (1.3).

$$84 \quad R(t) = \det(A_0 + U(t)B_0U^*(t)) \quad (4.1)$$

85 where t is real and $U(t)$ is some smooth function of unitary matrices.

86 Suppose $U(t)$ goes through a boundary matrix of interest, U_0 at $t = 0$.

87 Every unitary matrix can be written as an exponential of a skew-hermitian matrix.
 88 So we can write:

89 $U(t) = e^{S(t)}U_0$, where $S(t)$ is a smooth function of skew hermitian matrices with
 90 $S(0) = 0$.

91 Every choice of $S(t)$ with $S(0) = 0$, gives us every possible $U(t)$ that passes
 92 through U_0 at $t = 0$.

93 We wish to examine $U(t)$ and $R(t)$ near $t = 0$.

94 For small Δt ,

$$95 \quad U(\Delta t) = (e^{S(\Delta t)})U_0$$

$$96 \quad U(\Delta t) = (e^{S(0)+(\Delta t)S'(0)})U_0$$

$$97 \quad U(\Delta t) = (e^{(\Delta t)S'(0)})U_0$$

98 If we take the above function and plug it into $R(t)$ we'll get $R(\Delta t)$, but it won't
 99 be in a form useful to us. We use a result from [1], p.27 for this purpose. In order to
 100 state this result within the context of this paper, we first need the functional forms
 101 of the B-matrix, C-matrix, F-matrix (these were defined in section 2):

$$102 \quad B(t) = U(t)B_0U^*(t) \quad (4.2)$$

$$103 \quad C(t) = A_0 + B(t) \quad (4.3)$$

$$104 \quad F(t) = C^{-1}(t)A_0 - A_0C^{-1}(t) \quad (4.4)$$

105 Now we can state the result from [1]:

106 When $F(0) \neq 0$,

$$107 \quad R(\Delta t) = R(0) + (\Delta t) \det(C(0)) \text{tr}(S'(0)F(0)) + O((\Delta t)^2) \quad (4.5)$$

$$108 \quad R'(0) = \det(C(0)) \text{tr}(S'(0)F(0)) \quad (4.6)$$

110 If $F(0) = 0$ then U_0 is a permutation matrix and hence not a regular boundary
111 matrix (section 2). Our concern here is with regular boundary matrices so we will
112 assume $F(0) \neq 0$.

113 Note that $C(0)$ is just the C-matrix of U_0 and $F(0)$ is just the F-matrix of U_0 .
114 Also, $F(0)$ is only defined as long as $R(0) \neq 0$.

115 Assume U_0 is a regular boundary matrix with $R(0) \neq 0$. Then the tangent line to
116 the curve $R(t)$ at $t = 0$ must remain the same regardless of our choice of $S(t)$. This
117 is illustrated in Figure 1 where the closed curve indicates $\partial\Delta$. $R'(0)$ can be seen as
118 a vector in the complex plane. So all possible values of $R'(0)$ are either parallel or
119 anti-parallel.

120 $S'(0)$ is a skew hermitian matrix since the difference of skew-hermitian matrices
121 is also skew-hermitian. $S'(0)$ can turn out to be any skew-hermitian matrix.

122 *Proof.* Suppose we choose an arbitrary skew-hermitian matrix and multiply each
123 element of the matrix by t . Then we get a smooth function of skew-hermitian matrices
124 $S(t)$ with $S(0) = 0$ such that $S'(0)$ is the skew-hermitian matrix we initially chose. \square


125 So we can rewrite $R'(0)$ without any reference to the $S(t)$ function:

$$126 \quad R'(0) = \det(C(0)) \text{tr}(ZF(0)) \quad (4.7)$$

127 where Z is a skew-hermitian matrix. Since all values of $R'(0)$ are either parallel
128 or anti-parallel, all values of $\text{tr}(ZF(0))$ are parallel or anti-parallel, regardless of the
129 choice of Z . That gives us Theorem 3.1.

130 **5. Proof of Theorem 3.2.** In [1], p.26, Theorem 4, Bebiano and Queiró prove
131 that if within the neighborhood of a non-zero point $z \in \partial\Delta$, Δ is contained within an
132 angle less than π , then z must be a permutation point. We extend this result to show
133 that at a non-zero, non-permutation point, the boundary is smooth.

134 *Proof.* Given we have a non-zero, non-permutation point $z \in \partial\Delta$. Proof by
135 contradiction. Assume the boundary is not smooth at z . Since we know that Δ is not
136 contained within an angle less than π by Bebiano and Queiró's result, we know that
137 within the neighborhood of z , Δ is contained within some angle greater than π . So we


 FIG. 1. Region Δ with tangents at a boundary point

138 can find two smooth functions $R_1(t) \subseteq \Delta$ and $R_2(t) \subseteq \Delta$ such that $R_1(0) = R_2(0) = z$
 139 and $R'_1(0)$ is not parallel or anti-parallel to $R'_2(0)$.

140 Let U be a boundary matrix for z and let F be the F -matrix of U . Then using
 141 (4.6),

$$142 \quad R'_1(0) = \det(C) \operatorname{tr}(Z_1 F)$$

$$143 \quad R'_2(0) = \det(C) \operatorname{tr}(Z_2 F)$$

144 where Z_1 and Z_2 are two skew-hermitian matrices. But since $R'_1(0)$ and $R'_2(0)$
 145 are not parallel or anti-parallel, they form a basis for all the complex numbers as a
 146 vector space over the real numbers.

147 So $V = a \times \det(C) \operatorname{tr}(Z_1 F) + b \times \det(C) \operatorname{tr}(Z_2 F)$ goes in any direction depending
 148 on the choice of real numbers a and b .

$$149 \quad V = \det(C) (a \times \operatorname{tr}(Z_1 F) + b \times \operatorname{tr}(Z_2 F))$$

$$150 \quad V = \det(C) \operatorname{tr}((a \times Z_1 + b \times Z_2)F)$$

151 $Z_n = a \times Z_1 + b \times Z_2$ is also a skew-hermitian matrix.

152 So given any direction, there exists a skew-hermitian matrix Z_n such that $\det(C) \operatorname{tr}(Z_n F)$ ■
 153 goes in that direction. Hence there exists a smooth function $R_n(t) \subseteq \Delta$ such that
 154 $R_n(0) = z$, and $R'_n(0)$ is parallel or anti-parallel to that direction.

155 So there are functions going through z in all directions, contained within Δ . So
 156 z is not a boundary point. We arrive at a contradiction. □

157 This proves [Theorem 3.2](#).

158 **6. Proof of Theorem 3.3.** For $n = 3$, we define the following 12 skew-hermitian
 159 matrices with zero diagonal:

$$160 \quad Z_{12} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad Z_{13} = \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \quad Z_{23} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}$$

$$161 \quad Z_{21} = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad Z_{31} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix} \quad Z_{32} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}$$

$$162 \quad Z_{12,i} = Z_{21,i} = \begin{bmatrix} 0 & i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad Z_{13,i} = Z_{31,i} = \begin{bmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{bmatrix} \quad Z_{23,i} = Z_{32,i} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & i \\ 0 & i & 0 \end{bmatrix} \quad \blacksquare$$

163 Note that the commas do not indicate tensors. They're just used here as a label
 164 to distinguish imaginary and real matrices.

165 We define Z_{ab} and $Z_{ab,i}$ similarly for all $n > 3$, where $a \neq b$. For a given n we
 166 have $n(n-1)$ real matrices and $n(n-1)$ imaginary matrices.

167 Given a trace-argument constant unitary matrix U with F-matrix F . Suppose
 168 $F_{ab} = F_{ab,r} + iF_{ab,i}$

169 where $F_{ab,r}$ and $F_{ab,i}$ are real numbers.

$$170 \quad \operatorname{tr}(Z_{ab}F) = F_{ab} - F_{ba}$$

$$171 \quad \operatorname{tr}(Z_{ab,i}F) = (F_{ab} + F_{ba})i$$

172 Substitute in for F_{ab} and F_{ba}

$$173 \quad \operatorname{tr}(Z_{ab}F) = (F_{ab,r} - F_{ba,r}) + i(F_{ab,i} - F_{ba,i})$$

$$174 \quad \operatorname{tr}(Z_{ab,i}F) = (-F_{ab,i} - F_{ba,i}) + i(F_{ab,r} + F_{ba,r})$$

175 Since U is trace-argument constant,

$$(F_{ab,i} - F_{ba,i})(-F_{ab,i} - F_{ba,i}) = (F_{ab,r} + F_{ba,r})(F_{ab,r} - F_{ba,r})$$

We can simplify this to get:

$$F_{ab,r}^2 + F_{ab,i}^2 = F_{ba,r}^2 + F_{ba,i}^2$$

$$|F_{ab}| = |F_{ba}|$$

We can write:

$$F_{ab} = |F_{ab}| \angle \theta_{ab}$$

$$F_{ba} = |F_{ab}| \angle \theta_{ba}$$

slope of $tr(Z_{ab}F)$:

$$\frac{\sin(\theta_{ab}) - \sin(\theta_{ba})}{\cos(\theta_{ab}) - \cos(\theta_{ba})} = -\cot\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right)$$

similarly,

$$\text{slope of } tr(Z_{cd}F) = -\cot\left(\frac{\theta_{cd} + \theta_{dc}}{2}\right), \text{ where } c \neq d$$

$$\cot\left(\frac{\theta_{cd} + \theta_{dc}}{2}\right) = \cot\left(\frac{\theta_{ab} + \theta_{ba}}{2}\right)$$

therefore either:

$$\frac{\theta_{cd} + \theta_{dc}}{2} = \frac{\theta_{ab} + \theta_{ba}}{2}$$

or,

$$\frac{\theta_{cd} + \theta_{dc}}{2} = \frac{\theta_{ab} + \theta_{ba}}{2} + \pi$$

For some specific x, y where $x \neq y$

$$\text{let } \beta = \frac{\theta_{xy} + \theta_{yx}}{2}$$

$$\text{let } H = e^{-i\beta} F$$

For any $a \neq b$,

$$H_{ab} = |H_{ab}| \angle \alpha_{ab}$$

$$\frac{\alpha_{ab} + \alpha_{ba}}{2} = 0 \text{ or } \pi$$

Therefore H is zero-diagonal, with transpositional elements of equal magnitude and opposite arguments. Therefore H is hermitian.

We can write F as:

$$F = e^{i\beta} H$$

This completes our proof of [Theorem 3.3](#).

7. Conjectures. Before we state our conjectures we define a region Δ_S which is a restriction of Δ . See [\(1.1\)](#).

$$\Delta_S = \{ \det(A_0 + OB_0O^*) : O \in O(n) \} \quad (7.1)$$

206 where $O(n)$ is the set of $n \times n$ real orthogonal matrices.

207 As proven in [3], p.207, theorem 4.4.7, a matrix is normal and symmetric if and
208 only if it is diagonalizable by a real orthogonal matrix.

209 Therefore Δ_S is the set of determinants of sums of normal, symmetric matrices
210 with prescribed eigenvalues. We know Δ_S contains all the permutation points.

211 CONJECTURE 7.1 (Restricted Marcus-de Oliveira Conjecture).

$$212 \quad \Delta_S \subseteq \text{co}\left\{\prod (a_i + b_{\sigma(i)})\right\}$$

213 The above conjecture is supported by computational experiments.

214 CONJECTURE 7.2 (Boundary Conjecture).

$$215 \quad \partial\Delta \subseteq \partial\Delta_S$$

216 THEOREM 7.3. *If the boundary conjecture is true, the restricted Marcus-de Oliveira*
217 *conjecture implies the full Marcus-de Oliveira conjecture.*

218 *Proof.* The unitary group and the real orthogonal group are compact subsets of
219 the $n \times n$ complex matrices. Since a continuous image of a compact set is compact,
220 Δ and Δ_S are compact subsets of the complex plane. Hence they are both closed by
221 the Heine-Borel theorem.

222 Suppose we know Conjecture 7.1 is true. Then Δ_S along with its boundary is
223 within the convex-hull. Suppose we also know that Conjecture 7.2 is true. Then we
224 know that $\partial\Delta$ is inside the convex-hull. Can we have a unitary matrix U such that
225 $R(U)$ is outside the convex-hull? No, because that would mean we have points of Δ
226 on both the inside and outside of $\partial\Delta$. This is impossible since Δ is a closed set. So
227 Δ is within the convex hull proving Conjecture 1.1. \square

228 **8. Conclusion.** We hope that further analysis on boundary matrices of Δ , either
229 by expanding on the results in this paper, or novel research, leads to a proof of the
230 Boundary Conjecture. Then proving the full Marcus-de Oliveira conjecture would
231 amount to proving the restricted conjecture. Whether the restricted conjecture is any
232 easier to prove is unknown, but it's an avenue worth exploring.

233

REFERENCES

- 234 [1] N. BEBIANO AND J. QUERIO, *The determinant of the sum of two normal matrices with prescribed*
235 *eigenvalues*, Linear Algebra and its Applications, 71 (1985), pp. 23–28.
236 [2] G. N. DE OLIVEIRA, *Research problem: Normal matrices*, Linear and Multilinear Algebra, 12
237 (1982), pp. 153–154.
238 [3] R. HORN AND C. JOHNSON, *Matrix Analysis*, Cambridge University Press, 1990.
239 [4] M. MARCUS, *Derivations, plücker relations and the numerical range*, Indiana University Math
240 Journal, 22 (1973), pp. 1137–1149.
241 [5] A. K. N. BEBIANO AND J. DA PROVIDÊNCIA, *The validity of the marcus-de oliveira conjecture*
242 *for essentially hermitian matrices*, Linear Algebra and its Applications, 197-198 (1994),
243 pp. 411–427.