

On Tensors and Equations of the Electromagnetic Field

Yuriy A. Spirichev

Scientific Department, «Research and Design Institute of Radio-Electronic Engineering», Zarechny, Russia

Email address

yurii.spirichev@mail.ru (Yuriy A. Spirichev)

To cite this article

Yuriy A. Spirichev. On Tensors and Equations of the Electromagnetic Field. *American Journal of Modern Physics and Application*. Vol. 5, No. 3, 2018, pp. 50-54.

Received: May 29, 2018; Accepted: June 18, 2018;

Abstract

It is shown that the electromagnetic field is completely described by an asymmetric tensor of the second rank, which is a four-dimensional derivative of the electromagnetic potential. This tensor can be decomposed into the canonical antisymmetric and the new symmetric EMF tensor. From this tensor, in the form of its complete divergence, the EMF equations follow. One of them is an electromagnetic analog of the Lamé equation for an elastic medium. It is shown that the longitudinal waves of the divergence of the vector potential propagate at a speed greater than the speed of light and do not have a magnetic component.

Keywords

Electromagnetic Field, Asymmetric Tensor, Symmetric Tensor, Maxwell Equations, Longitudinal Waves

1. Introduction

The theoretical basis of the classical theory of the electromagnetic field (EMF) is Maxwell's equations, generalizing the experimental results obtained by the end of the 18th century. The development of the classical theory of EMF led to its description in the form of an antisymmetric tensor of the second rank, from which the Maxwell equations follow. These equations played a key role in the development of theoretical physics and had a strong influence on the creation of a special theory of relativity and other theories. Already at the beginning of the twentieth century, classical electrodynamics was considered to be a complete science, and the theory of EMF was further developed in the form of quantum electrodynamics. Despite this, in the classical theory of EMF there were some vague spots and controversial issues. For example, for about a hundred years there is the Abraham-Minkowski problem, the crux of the problem lay in the absence of a common opinion about the correct energy-momentum tensor of the interaction of EMF with matter, the form of the angular momentum in matter and the existence of the Abraham electromagnetic force [1-13]. This situation leads to the search for new energy-momentum tensors of EMF [14-16]. At present, the bibliography on this problem is about 300 works [17]. Another problem is the

mechanism for transferring the angular momentum by a plane electromagnetic wave [18-22]. The problem is that the canonical EMF wave equations do not describe this process. Until recently, electrodynamics did not even have wave equations for the energy and momentum of EMF. Such equations, following from the new energy-momentum tensor and describing the transfer of the angular momentum of the electromagnetic wave, were obtained by in work [16]. In the classical theory of EMF, Newton's third law is not always satisfied in the interaction of arbitrarily moving electric charges and non-parallel currents. This led to the hypothesis of the existence of a "scalar (potential) magnetic field" [23], the introduction of which into electrodynamics makes it possible to ensure the fulfillment of Newton's third law in all cases. The reality of the scalar magnetic field is confirmed in experiments on the longitudinal interaction of direct currents [23, 24]. There is the problem of longitudinal electromagnetic waves in vacuum [25, 26], which is that Maxwell's equations allow the existence of longitudinal waves of a scalar electromagnetic potential, but it has not been possible to detect such waves experimentally yet. The most obvious incompleteness of classical electrodynamics is manifested in plasma theory. Until now, there is no understanding of what electromagnetic forces hold charged particles in ball lightning, and the problem of prolonged plasma confinement in existing technical installations,

despite half a century of intensive work, is far from being solved. There is no understanding of the cause of the existence of hot spots in Z-pinchs, the phenomenon of the magnetic dynamo and a number of other plasma phenomena. The above mentioned problems require reasonable attention to the basics of the classical EMF theory and to the Maxwell equations themselves.

The purpose of this article is to consider the foundations of the classical theory of EMF with the aim of eliminating the existing discussion issues in it and the convergence of the classical theory with quantum electrodynamics.

One of the reasons for posing this problem was the author's opinion that the description of EMF using the canonical antisymmetric tensor is incomplete and does not ensure the mathematical correctness of the EMF sources introduction into its equations. The reason for this was the following. Maxwell's equations with sources follow from the canonical antisymmetric EMF tensor in the form of a four-dimensional divergence along one of its indices, which is equated to the source of the field in the form of a four-dimensional current density [27]. But the antisymmetric tensor of the second rank has divergences for each of the indices. These two divergences have opposite signs, so the total divergence of the antisymmetric EMF tensor as a four-dimensional rotor, is zero and cannot have a field source. Therefore, equating only one of the divergences of the antisymmetric EMF tensor to the source of the field is mathematically incorrect.

Usually, the antisymmetric EMF tensor is written in the form $F_{[\mu\nu]} = \partial_\mu \mathbf{A}_\nu - \partial_\nu \mathbf{A}_\mu$, where \mathbf{A}_ν is the four-dimensional electromagnetic potential. The first term of this expression is the four-dimensional derivative of the electromagnetic potential and is an asymmetric tensor of the second rank $F_{\mu\nu} = \partial_\mu \mathbf{A}_\nu$. This EMF tensor can be written in the form of its expansion into symmetric and antisymmetric tensors $F_{\mu\nu} = F_{[\mu\nu]}/2 + F_{(\mu\nu)}/2$. The first term of this expansion is the canonical antisymmetric EMF tensor $F_{[\mu\nu]} = \partial_\mu \mathbf{A}_\nu - \partial_\nu \mathbf{A}_\mu$, and the second term represents the new symmetric EMF tensor $F_{(\mu\nu)} = \partial_\mu \mathbf{A}_\nu + \partial_\nu \mathbf{A}_\mu$. Thus, a complete description of the EMF is an asymmetric tensor of

$$F_{(\mu\nu)} = \begin{pmatrix} 2\frac{1}{c^2}\partial_t\varphi & \frac{1}{c}i\cdot(\partial_t A_x - \partial_x\varphi) & \frac{1}{c}i\cdot(\partial_t A_y - \partial_y\varphi) & \frac{1}{c}i\cdot(\partial_t A_z - \partial_z\varphi) \\ \frac{1}{c}i\cdot(\partial_t A_x - \partial_x\varphi) & 2\partial_x A_x & (\partial_x A_y + \partial_y A_x) & (\partial_x A_z + \partial_z A_x) \\ \frac{1}{c}i\cdot(\partial_t A_y - \partial_y\varphi) & (\partial_x A_y + \partial_y A_x) & 2\partial_y A_y & (\partial_y A_z + \partial_z A_y) \\ \frac{1}{c}i\cdot(\partial_t A_z - \partial_z\varphi) & (\partial_x A_z + \partial_z A_x) & (\partial_y A_z + \partial_z A_y) & 2\partial_z A_z \end{pmatrix} \quad (2)$$

The canonical antisymmetric EMF tensor $F_{[\mu\nu]}$ describes the four-dimensional rotation of the EMF. Then, by analogy with a continuous medium, the symmetric tensor $F_{(\mu\nu)}$ describes the four-dimensional deformation of the EMF. The

the second rank $F_{\mu\nu} = \partial_\mu \mathbf{A}_\nu$. From this tensor, in the form of its divergences, the EMF equations follow. Since the total divergence of the canonical antisymmetric tensor, as a four-dimensional rotor, is identically zero, the EMF equations in the form of a full four-dimensional divergence follow from the symmetric tensor $F_{(\mu\nu)} = \partial_\mu \mathbf{A}_\nu + \partial_\nu \mathbf{A}_\mu$. This divergence of the symmetric tensor should be attributed to EMF sources in the form of charges and currents.

In this article EMF and its sources are considered in a vacuum. The geometry of space-time is taken in the form of pseudo-Euclidean Minkowski space in the form (ct, ix, iy, iz) and differentiation between covariant and contravariant indexes is irrelevant. The four-dimensional electromagnetic potential is defined as $\mathbf{A}_\nu(\varphi/c, i\mathbf{A})$, where φ and \mathbf{A} are the scalar and vector potentials of the EMF. The four-dimensional current density is defined as $\mathbf{J}_\nu(\rho \cdot c, i\mathbf{J})$, where ρ and \mathbf{J} are the electric charge density and current density.

2. Asymmetric and Symmetric Electromagnetic Field Tensors

The asymmetric EMF tensor $F_{\mu\nu} = \partial_\mu \mathbf{A}_\nu$, which is a four-dimensional derivative of the electromagnetic potential, is written in the matrix form:

$$F_{\mu\nu} = \partial_\mu \mathbf{A}_\nu = \begin{pmatrix} \frac{1}{c^2}\partial_t\varphi & \frac{1}{c}i\cdot\partial_t A_x & \frac{1}{c}i\cdot\partial_t A_y & \frac{1}{c}i\cdot\partial_t A_z \\ -\frac{1}{c}i\cdot\partial_x\varphi & \partial_x A_x & \partial_x A_y & \partial_x A_z \\ -\frac{1}{c}i\cdot\partial_y\varphi & \partial_y A_x & \partial_y A_y & \partial_y A_z \\ -\frac{1}{c}i\cdot\partial_z\varphi & \partial_z A_x & \partial_z A_y & \partial_z A_z \end{pmatrix} \quad (1)$$

This asymmetric tensor can be decomposed into antisymmetric and symmetric tensors $F_{\mu\nu} = F_{[\mu\nu]}/2 + F_{(\mu\nu)}/2$. The antisymmetric EMF tensor is well known in electrodynamics [27] and it is not given here. Write the symmetric tensor EMF $F_{(\mu\nu)}$ in the matrix form:

members of its diagonal describe the volume deformation of the EMF expansion/contraction, and the remaining terms describe four-dimensional shear deformations.

3. Equations of the Electromagnetic Field Without Field Sources

Let us write down four-dimensional divergences with respect to the indices μ and ν of the asymmetric tensor $F_{\mu\nu} = \partial_\mu \mathbf{A}_\nu$ (differentiation between covariant and contravariant indexes is irrelevant):

$$\partial_\mu F_{\mu\nu} = \partial_\mu (\partial_\mu \mathbf{A}_\nu) \text{ and } \partial_\nu F_{\mu\nu} = \partial_\nu (\partial_\mu \mathbf{A}_\nu) \quad (3)$$

Write these Eqs. in expanded form:

$$\frac{1}{c^2} \partial_{tt} \varphi - \Delta \varphi = 0 \quad (4)$$

$$\frac{1}{c^2} \partial_{tt} \mathbf{A} - \Delta \mathbf{A} = 0 \quad (5)$$

$$\partial_t \left(\frac{1}{c^2} \partial_t \varphi + \nabla \cdot \mathbf{A} \right) = 0 \quad (6)$$

$$-\nabla \left(\frac{1}{c^2} \partial_t \varphi + \nabla \cdot \mathbf{A} \right) = 0 \quad (7)$$

Eqs. (4) and (5) represent Maxwell's canonical equations in the Lorentz gauge $\partial_t \varphi / c^2 + \nabla \cdot \mathbf{A} = 0$. Eqs. (6) and (7) represent, respectively, the derivatives with respect to time and space of the Lorentz gauge condition. By adding Eqs. (4) and (6), and also (5) and (7) the complete divergence of the asymmetric EMF tensor is obtained:

$$2 \frac{1}{c^2} \partial_{tt} \varphi + \partial_t \nabla \cdot \mathbf{A} - \Delta \varphi = 0 \quad (8)$$

$$\frac{1}{c^2} \partial_{tt} \mathbf{A} - \frac{1}{c^2} \partial_t \nabla \varphi - \nabla (\nabla \cdot \mathbf{A}) - \Delta \mathbf{A} = 0 \quad (9)$$

Let us write down four-dimensional divergences with respect to the indices μ and ν of the symmetric tensor $F_{(\mu\nu)} = \partial_\mu \mathbf{A}_\nu + \partial_\nu \mathbf{A}_\mu$. Since the tensor $F_{(\mu\nu)}$ is symmetric, these divergences are equal:

$$\partial_\mu (\partial_\mu \mathbf{A}_\nu + \partial_\nu \mathbf{A}_\mu) = \partial_\nu (\partial_\mu \mathbf{A}_\nu + \partial_\nu \mathbf{A}_\mu) = 0 \quad (10)$$

Writing these equations in expanded form, we obtain two Eqs. (8) and (9), completely describing the motions of the EMF. These two equations replace Maxwell's canonical equations in potentials:

$$\nabla \cdot (-\nabla \varphi - \partial_t \mathbf{A}) = \nabla \cdot \mathbf{E} = 0 \quad (11)$$

$$\frac{1}{c^2} \partial_{tt} \mathbf{A} + \frac{1}{c^2} \partial_t \nabla \varphi + \nabla \times \nabla \times \mathbf{A} = -\frac{1}{c^2} \partial_t \mathbf{E} + \nabla \times \mathbf{B} = 0 \quad (12)$$

For the static case, Eq. (8) describes the Gaussian law for

an electric field without sources $\nabla \cdot \mathbf{E} = -\Delta \varphi = 0$. Eq. (9) can be written in the form:

$$\partial_{tt} \mathbf{A} - 2c^2 \nabla (\nabla \cdot \mathbf{A}) + c^2 \nabla \times \nabla \times \mathbf{A} = -\partial_t \nabla \varphi \quad (13)$$

This equation is an electromagnetic analog of the Lamé equation (or the dynamic Navier-Stokes equation), known in the linear theory of elasticity and describing the wave motion of a continuous elastic medium [28]:

$$\partial_{tt} \mathbf{U} - \nu_1^2 \cdot \nabla (\nabla \cdot \mathbf{U}) + \nu_2^2 \cdot \nabla \times \nabla \times \mathbf{U} = \mathbf{G} \quad (14)$$

where \mathbf{U} is the displacement vector of the medium, ν_1 is the velocity of longitudinal waves, ν_2 is the velocity of transverse waves, and \mathbf{G} is the external force. Comparison of this equation with Eq. (13) shows that the velocity of the longitudinal EMF waves is $\sqrt{2}$ greater than the transverse wave velocity, i.e. speed of light. The right-hand side of Eq. (13) describes the EMF wave source in the form of an alternating potential electric field. This source of electromagnetic waves is an electric dipole in the form of an electric capacitor deployed in space.

Eq. (8) can be written in the form:

$$\frac{2}{c^2} \partial_{tt} \varphi - \Delta \varphi = \nabla \cdot (-\partial_t \mathbf{A}) \quad (15)$$

This equation can be interpreted as a wave equation for a scalar electromagnetic potential. It follows from this that the waves of the scalar electromagnetic potential propagate at a speed that is $\sqrt{2}$ times slower than the speed of light. The source of the waves of the scalar potential is the divergence of the time-varying vector potential or the divergence of the vortex electric field.

4. Equations of the Electromagnetic Field with Field Sources

Since the total four-dimensional divergence of a symmetric tensor can be nonzero, we equate it with the source of the EMF, i.e. to a four-dimensional current density of $\partial_\mu F_{(\mu\nu)} = \mathbf{J}_\nu$ or $\partial_\mu (\partial_\mu \mathbf{A}_\nu + \partial_\nu \mathbf{A}_\mu) = \mathbf{J}_\nu$. Let us write down this complete four-dimensional divergence of a symmetric tensor $F_{(\mu\nu)}$ with sources in expanded form:

$$2 \frac{1}{c^2} \partial_{tt} \varphi + \partial_t \nabla \cdot \mathbf{A} - \Delta \varphi = \rho / \epsilon_0 \quad (16)$$

$$\frac{1}{c^2} \partial_{tt} \mathbf{A} - \frac{1}{c^2} \partial_t \nabla \varphi - \nabla (\nabla \cdot \mathbf{A}) - \Delta \mathbf{A} = \mu_0 \cdot \mathbf{J} \quad (17)$$

Eq. (16) in statics describes the Gaussian law for an electric field with sources $\nabla \cdot \mathbf{E} = -\Delta \varphi = \rho / \epsilon$, and Eq. (17) replaces the Ampere-Maxwell total current equation. Eq. (17) can be written in the form:

$$\frac{1}{c^2} \partial_{tt} \mathbf{A} - \frac{1}{c^2} \partial_t \nabla \varphi - 2 \cdot \nabla (\nabla \cdot \mathbf{A}) + \nabla \times \nabla \times \mathbf{A} = \mu_0 \cdot \mathbf{J} \quad (18)$$

In this equation, the fourth term represents the magnetic field rotor, and the third term describes the gradient of the scalar magnetic field, hypothetically introduced by Nikolaev [23]. This term ensures the fulfillment in electrodynamics of Newton's third law in the interaction of arbitrarily moving

electric charges and non-parallel currents. For the stationary case, Eq. (18) can be written in the form of an equation describing the Ampere law, but in which there is a Nikolaev scalar magnetic field:

$$-2 \cdot \nabla (\nabla \cdot \mathbf{A}) + \nabla \times \nabla \times \mathbf{A} = \mu_0 \cdot \mathbf{J} \quad (19)$$

We take the rotor from both sides of Eq. (17) and obtain the well-known wave equation for the magnetic field:

$$\frac{1}{c^2} \partial_{tt} (\nabla \times \mathbf{A}) - \Delta (\nabla \times \mathbf{A}) = \mu_0 \cdot (\nabla \times \mathbf{J}) \text{ or } \frac{1}{c^2} \partial_{tt} \mathbf{B} - \Delta \mathbf{B} = \mu_0 \cdot (\nabla \times \mathbf{J}) \quad (20)$$

We take the divergence from both sides of Eq. (18) and obtain the equation of longitudinal waves of the divergence of the vector potential:

$$\frac{1}{c^2} \partial_{tt} \nabla \cdot \mathbf{A} - \frac{1}{c^2} \partial_t \Delta \varphi - 2 \cdot \Delta (\nabla \cdot \mathbf{A}) = \mu_0 \nabla \cdot \mathbf{J} \text{ or } \frac{1}{2c^2} \partial_{tt} \nabla \cdot \mathbf{A} - \Delta (\nabla \cdot \mathbf{A}) = \mu_0 \nabla \cdot \mathbf{J} + \frac{1}{c^2} \partial_t \Delta \varphi \quad (21)$$

From this equation follows the previously made conclusion that the velocity of longitudinal EMF waves is $\sqrt{2}$ times greater than the speed of light. It also follows from this equation that in the longitudinal EMF waves there is no magnetic component and they can be called electroscalar waves.

5 Conclusion

The existing EMF theory has a number of controversial and unresolved issues. These include the shape of the energy-momentum tensor, the transfer of the angular momentum of the electromagnetic wave, the fulfillment of Newton's third law, the explanation of plasma phenomena, such as ball lightning, Z-pinches, and others. This leads to a revision of the existing theory. In the existing theory of EMF, the field equations follow from the canonical antisymmetric tensor in the form of its divergence. This divergence equates to the sources of the field. However, the canonical antisymmetric tensor of EMF has four-dimensional divergences with opposite signs for each of the indices, so the introduction of the field source into its divergence by only one of the indices is incorrect. The total divergence of the antisymmetric tensor, like a four-dimensional rotor, equals zero and cannot have a divergence as a source of EMF. Consequently, it cannot be attributed to the field sources in the form of charges and currents.

It is shown that the complete EMF description is an asymmetric second-rank tensor $F_{\mu\nu} = \partial_\mu \mathbf{A}_\nu$, which is a four-dimensional derivative of the electromagnetic potential. This tensor can be decomposed into the canonical antisymmetric and the new symmetric EMF tensors. For a symmetric EMF tensor, the divergences may not be equal to zero, hence, they can be attributed to EMF sources. From this tensor in the form of its complete divergence the EMF equations follow in which one can introduce charges and currents.

These equations describe transverse and longitudinal EMF waves. One of these equations is an electromagnetic analog of the Lamé equation for an elastic medium. Longitudinal waves do not have a magnetic component and propagate at a

speed $\sqrt{2}$ times greater than the speed of light. The field equation replacing the Ampere-Maxwell equation includes a hypothetical scalar magnetic field that ensures the fulfillment of Newton's third law in electrodynamics. Separate equations for the transverse and longitudinal EMF waves are obtained.

References

- [1] I. Brevik, Minkowski momentum resulting from a vacuum-medium mapping procedure, and a brief review of Minkowski momentum experiments, *Annals of Physics* 377 (2017) 10–21.
- [2] R. Medina and J. Stephany, The energy-momentum tensor of electromagnetic fields in matter, arXiv: 1703.02109.
- [3] M. G. Silveirinha, Revisiting the Abraham-Minkowski Dilemma, arXiv: 1702.05919.
- [4] J. J. Bisognano, Electromagnetic Momentum in a Dielectric: a Back to Basics Analysis of the Minkowski-Abraham Debate, arXiv: 1701.08683.
- [5] Yu. A. Spirichev, Electromagnetic energy, momentum and forces in a dielectric medium with losses, arXiv: 1705.08447.
- [6] M. E. Crenshaw, The Role of Conservation Principles in the Abraham-Minkowski Controversy, arXiv: 1604.01801.
- [7] C. Wang, Is the Abraham electromagnetic force physical?, *Optik*, (2016) 127, 2887–2889.
- [8] P. L. Saldanha, J. S. Oliveira Filho, Hidden momentum and the Abraham-Minkowski debate, arXiv: 1610.05785.
- [9] M. Testa, A Comparison between Abraham and Minkowski Momenta, *Journal of Modern Physics*, 2016, 7, 320-328.
- [10] C. J. Sheppard, B. A. Kemp, *Phys. Rev. A* 93 (2016) 053832.
- [11] N. Toptygin, K. Levina, *Phys. Usp.* 59 141 (2016).
- [12] V. V. Nesterenko, A. V. Nesterenko, Ponderomotive forces in electrodynamics of moving medium: The Minkowski and Abraham approaches, arXiv: 1604.01708.
- [13] C. Wang, J. Ng, M. Xiao, C. T. Chan, *Sci. Adv.* 2 (2016) e1501485.

- [14] M. Abragam, *Be. Circ. mat. Palermo* 28, 1 (1909), 31, 527 (1910).
- [15] F. J. Belinfante, *Physica* 6, 887 (1939).
- [16] Yu. A. Spirichev, A new form of the energy-momentum tensor of the interaction of an electromagnetic field with a non-conducting medium. The wave equations. The electromagnetic forces, arXiv: 1704.03815.
- [17] McDonald K T, Bibliography on the Abraham-Minkowski debate (February 17, 2015, updated September 29, 017), <http://physics.princeton.edu/~mcdonald/examples/ambib.pdf>
- [18] E. Leader and C. Lorce, The angular momentum controversy: What's it all about and does it matter? *Phys. Rep.* 541, 163 (2014).
- [19] K. Y. Bliokh, A. Y. Bekshaev and F. Nori, Optical momentum and angular momentum in dispersive media: From the Abraham-Minkowski debate to unusual properties of surface plasmon-polaritons, arXiv: 1706.05493.
- [20] K. Y. Bliokh, A. Y. Bekshaev and F. Nori, Optical Momentum, Spin, and Angular Momentum in Dispersive Media, arXiv: 1706.06406.
- [21] E. Leader, The photon angular momentum controversy: Resolution of a conflict between laser optics and particle physics, *Phys. Lett. B* 756, 303 (2016).
- [22] Aiello, P. Banzer, M. Neugebauer, and G. Leuchs, From transverse angular momentum to photonic wheels, *Nature Photon.* 9, 789 (2015).
- [23] G. V. Nikolaev, *Non-contradictory electrodynamics. Theories, experiments, paradoxes*, (Tomsk, NTL Publishing, 1997).
- [24] A. K. Tomilin, The Fundamentals of Generalized Electrodynamics, arXiv: 0807.2172.
- [25] V. Simulik, Longitudinal electromagnetic waves in the framework of standard classical electrodynamics, arXiv: 1606.01738.
- [26] S. Nedic, Longitudinal Waves in Electromagnetism, *INFOTEH-JAHORINA*, Vol. 16, March 2017.
- [27] L. D. Landau, E. M. Lifshits, *The Classical Theory of Fields*, (Oxford: Pergamon Press, 1983).
- [28] L. D. Landau, E. M. Lifshits, *The Theory of elasticity*, (Oxford: Pergamon Press, 1983).