

A thought about the Riemann hypothesis.

Abstract. In these papers we will try to face the Riemann hypothesis, basing on the study of the functional equation of the Riemann zeta function.

0 Introduction and known properties. [1]

The Riemann zeta function is defined by the Dirichlet series:

$$\mathbf{0.1} \quad \zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

for every complex number s with real part $\text{Re}(s)$ greater than 1.

However, this function can be analytically continued to a holomorphic function on the whole complex s -plane, except for a simple pole at $s=1$, which satisfies the following functional equation:

$$\mathbf{1} \quad \zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s),$$
 where s is a complex number and Γ is the gamma function.

The zeta function has some zeros, called the trivial zeros, and others, known as the non-trivial zeros, which lie in the critical strip $0 < \text{Re}(s) < 1$, symmetrically about the real line.

Thus for every non-trivial zero $z = \sigma + it$:

$$\mathbf{0.2} \quad \exists! z_n \mid z_n = \sigma - it$$

The Riemann hypothesis is about the non-trivial zeros and it asserts that:

“Every non-trivial zero s has $\text{Re}(s)=1/2$ ”.

In addition, for the zeta function **(0.1)** there is the following identity, which is the Euler product formula:

$$\mathbf{0.3} \quad \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \text{ prime}} \frac{1}{1-p^{-s}},$$

where s is a complex number with real part $\text{Re}(s)$ greater than 1 and p is a prime number.

Proof p(0) [Euler]:

We consider the zeta function **(0.1)**

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = 1 + \frac{1}{2^s} + \frac{1}{3^s} + \frac{1}{4^s} + \dots$$

We multiply it by the factor $\frac{1}{2^s}$

$$\frac{1}{2^s} \zeta(s) = \frac{1}{2^s} + \frac{1}{4^s} + \frac{1}{6^s} + \frac{1}{8^s} + \dots$$

Subtracting the second equation from the first we get:

$$\left(1 - \frac{1}{2^s}\right) \zeta(s) = 1 + \frac{1}{3^s} + \frac{1}{5^s} + \frac{1}{7^s} + \dots$$

Doing this, we have removed all elements that have a factor of 2.

Repeating the same procedure for the next terms:

$$\left(1 - \frac{1}{7^s}\right) \left(1 - \frac{1}{5^s}\right) \left(1 - \frac{1}{3^s}\right) \left(1 - \frac{1}{2^s}\right) \zeta(s) = 1$$

In this way, we are progressively removing all the multiples of every term after 1, which is a prime number since it is not a multiple of any lesser number. Therefore, the numbers of the product are all primes.

Repeating infinitely for $\frac{1}{p^s}$, where p is a prime, we obtain:

$$\dots \left(1 - \frac{1}{11^s}\right) \left(1 - \frac{1}{7^s}\right) \left(1 - \frac{1}{5^s}\right) \left(1 - \frac{1}{3^s}\right) \left(1 - \frac{1}{2^s}\right) \zeta(s) = 1$$

Dividing both sides by everything but $\zeta(s)$, we get:

$$\zeta(s) = \frac{1}{\left(1 - \frac{1}{2^s}\right)} \frac{1}{\left(1 - \frac{1}{3^s}\right)} \frac{1}{\left(1 - \frac{1}{5^s}\right)} \frac{1}{\left(1 - \frac{1}{7^s}\right)} \frac{1}{\left(1 - \frac{1}{11^s}\right)} \dots,$$

which is equivalent to:

$$\zeta(s) = \prod_{p \text{ prime}} \frac{1}{1 - p^{-s}} \quad \text{(0.3)}$$

From the Euler product formula (0.3) we deduce an important consequence:

(0.4) "the zeta function has no zeros in the half-plane $\text{Re}(s) > 1$ ".

1 Study of the functional equation

In order to study the functional equation, we factor it.

1.1 factor: 2^s

The factor 2^s is an exponential function of the form $y = a^s$, with $a=2$ and $s \in \mathbb{C}$. Since the base of the exponent is different from 0, there is no value of s such that cancels the factor 2^s and therefore the functional equation (1). Thus:

$$\text{1.11 } \nexists s \in \mathbb{C} \mid 2^s = 0 \rightarrow (1) = 0$$

1.2 factor: π^{s-1}

Doing a similar reasoning to that used for the factor (1.1), we can assert that there is no value of s such that cancels the factor π^{s-1} and therefore the functional equation (1). Thus:

$$\text{1.21 } \nexists s \in \mathbb{C} \mid \pi^{s-1} = 0 \rightarrow (1) = 0$$

1.3 factor: $\sin\left(\frac{\pi s}{2}\right)$

Since the factor $\sin\left(\frac{\pi s}{2}\right)$ is a goniometric function of the form $y = \sin(x)$, it cancels the functional

equation (1) when the argument of the sin equals to $k\pi$, with $k \in \mathbb{Z}$.

In this case we have that:

$$\frac{\pi s}{2} = k\pi$$

$$\pi s = 2k\pi$$

$$\text{1.31 } s = 2k, \text{ with } k \in \mathbb{Z}$$

Because the zeta function (0.1) has no zeros for (0.4), the formula (1.31) becomes:

$$\text{1.32 } s = -2n, \text{ with } n \in \mathbb{N} - \{0\}$$

Thus, when $s = -2n$, with $n \in \mathbb{N} - \{0\}$, the functional equation (1) equals 0. Therefore:

$$\text{1.33 } \zeta(-2n) = 0, \text{ with } n \in \mathbb{N} - \{0\}$$

The trivial zeros of the zeta function derive from the formula **(1.33)**.

1.4 factor: $\Gamma(1 - s)$ [2]

Since the gamma function has the property to be non-zero everywhere, there is no value of s such that cancels the factor $\Gamma(1 - s)$ and therefore the functional equation **(1)**. Thus:

$$1.41 \quad \nexists s \in \mathbb{C} \mid \Gamma(1 - s) = 0 \rightarrow (1) = 0$$

1.5 factor: $\zeta(1 - s)$

[In the next paragraph we will concentrate upon the factor **(1.5)**].

2 Study of the factor $\zeta(1 - s)$

In order to study this factor, we will think about the following question:

*"What happen when $\zeta(1 - s)=0$, canceling the functional equation **(1)**?"*

We formulate two hypothesis:

Hypothesis A: *"If $\zeta(1 - s)=0$, then $\zeta(1 - s) \neq \zeta(s)$ ".*

Hypothesis B: *"If $\zeta(1 - s)=0$, then $\zeta(1 - s) = \zeta(s)$ ".*

From the functional equation we demonstrate that the hypothesis **A** is false, while the hypothesis **B** is true.

Proof p(1):

Considered the functional equation **(1)**, if $\zeta(1 - s)=0$, for the zero-product property, the functional equation equals 0. From this, we work out the following relation:

$$2.1 \quad \zeta(1 - s)=0 \rightarrow \zeta(s) = 0$$

In fact:

$$\text{If } \zeta(1 - s)=0$$

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1 - s) \zeta(1 - s)$$

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1 - s) 0$$

$$\zeta(s) = 0$$

Therefore, $\zeta(1 - s)=0 \rightarrow \zeta(s) = 0$ **(2.1)**

From the relation **(2.1)**, the hypothesis **(A)** is false, whilst the hypothesis **(B)** is true:

(A) $\zeta(1 - s)=0 \rightarrow \zeta(s) = 0 \rightarrow \zeta(1 - s) \neq \zeta(s) \rightarrow$ (false for proof **p(1)**).

(B) $\zeta(1 - s)=0 \rightarrow \zeta(s) = 0 \rightarrow \zeta(1 - s) = \zeta(s) \rightarrow$ (true for proof **p(1)**).

From the veracity of the hypothesis **(B)** we deduce a key condition satisfied when $\zeta(1 - s)=0$ that is:

$$2.2 \quad \zeta(s) = \zeta(1 - s)$$

From the condition **(2.2)** we deduce an important consequence:

"The trivial zeros of the zeta function don't derive from the factor $\zeta(1 - s)$ ".

Proof p(2):

Let $s = -2n$, with $n \in \mathbb{N} - \{0\}$, we substitute it in the functional equation **(1)**:

$$\zeta(-2n) = 2^{-2n} \pi^{-2n-1} \sin\left(\frac{\pi(-2n)}{2}\right) \Gamma(1 - (-2n)) \zeta(1 - (-2n))$$

$$\zeta(-2n) = 2^{-2n} \pi^{-2n-1} \sin(-n\pi) \Gamma(1+2n) \zeta(1+2n)$$

Done the substitution and the related calculations, $\zeta(s)$ and $\zeta(1-s)$ have become respectively $\zeta(-2n)$ and $\zeta(1+2n)$.

For the formula **(1.33)** $\zeta(-2n) = 0$, while $\zeta(1+2n) \neq 0$, because the zeta function of the form **(0.1)** has no zeros.

Therefore $\zeta(-2n) \neq \zeta(1+2n)$ and thus $\zeta(s) \neq \zeta(1-s)$.

Consequently the condition **(2.2)** is not satisfied and so the trivial zeros don't derive from the factor **(1.5)**, but only from **(1.3)**.

3 Study of the condition $\zeta(s) = \zeta(1-s)$ in the critical strip

We consider the functional equation **(1)**. From what we have said in the previous paragraphs, we deduce that the non-trivial zeros of the zeta function don't derive from the factors **(1.1)**, **(1.2)**, **(1.4)**, because they don't cancel the functional equation, and even from the factor **(1.3)**, from which the sole trivial zeros derive. Thus they only derive from the factor **(1.5)** and satisfy the condition **(2.2)**.

Now, we suppose that z_0 is a non-trivial zero with real part $\sigma \neq \frac{1}{2}$ and imaginary part it :

$$z_0 = \sigma + it$$

We substitute z_0 in the condition **(2.2)**:


$$\zeta(z_0) = \zeta(1-z_0)$$

From this, we obtain a second non-trivial zero z_1 , which is equal to $1-z_0$:

$$z_1 = 1 - z_0 = (1 - \sigma) - it$$

We represent z_0 and z_1 in the complex plane:

Picture 1




From the property **(0.2)** of the non-trivial zero we obtain z_2 and z_3 :

$$z_2 = \sigma - it$$

$$z_3 = (1 - \sigma) + it$$

We represent all the non-trivial zeros in the complex plane:

Picture 2



Since z_0, z_1, z_2 and z_3 are non-trivial zeros, we deduce that $\zeta(z_0) = \zeta(z_1) = \zeta(z_2) = \zeta(z_3)$. Therefore every couple of non-trivial zeros satisfies the following relation:

$$\mathbf{3.1} \quad \zeta(z_j) = \zeta(z_k)$$

So, for the non-trivial zeros both relations **(2.2)** and **(3.1)** are satisfied and combining them together we obtain this final relation, which must be satisfied by all the couples of non-trivial zeros, which have imaginary part $\pm it$:

$$\mathbf{3.2} \quad \begin{cases} \zeta(z_k) = \zeta(s) \\ \zeta(z_j) = \zeta(1 - s) \end{cases}$$

We have shown that if z_0 is a non-trivial zero with real part $\sigma \neq \frac{1}{2}$, there will be another three non-trivial zeros (z_1, z_2, z_3) such that we will have the situation represented in the picture **2**.

Now, we consider all the possible couples of non-trivial zeros:

$$\{(z_0; z_1), (z_0; z_2), (z_0; z_3), (z_1; z_2), (z_1; z_3), (z_2; z_3)\}$$

From the relation **(3.2)** we deduce that only the couples $(z_0; z_1)$ and $(z_2; z_3)$ satisfy the relation. In fact, since in the relation **(3.2)** one zero is equal to $\zeta(s)$ and the other is equal to $\zeta(1 - s)$, these zeros must be symmetric about the point $P(\frac{1}{2}; 0)$.

Therefore the disposition of the non-trivial zeros, shown in the picture 2, is an absurd because the couples $(z_0; z_2)$, $(z_0; z_3)$, $(z_1; z_2)$ and $(z_1; z_3)$ don't satisfy the relation (3.2); in fact they aren't symmetric about the point $P(\frac{1}{2}; 0)$, but about the real line, for $(z_0; z_2)$ and $(z_1; z_3)$, and about the critical line $x=\frac{1}{2}$ for $(z_0; z_3)$ and $(z_1; z_2)$.

The only case, for which the relation (3.2) and the property (0.2) of the non-trivial zeros are both true, is when z_0 and z_1 coincide respectively with z_3 and z_2 .

Thus we have:

$$z_0 = z_3$$

$$\sigma + it = (1 - \sigma) + it$$


$$\sigma = 1 - \sigma$$

$$2\sigma = 1$$

$$\sigma = \frac{1}{2}$$

We represent the situation in the complex plane:

Picture 3



From the result, represented in the picture 3, we substitute a complex number of the form $\frac{1}{2} + it$ in the condition (2.2) and we get:

$$\mathbf{3.3} \quad \zeta\left(\frac{1}{2} + it\right) = \zeta\left(\frac{1}{2} - it\right)$$

This relation is true only if the complex number $\frac{1}{2} + it$ is a zero that is when $\zeta\left(\frac{1}{2} + it\right) = 0$.

Therefore the formula (3.3) is true if:

$$\mathbf{3.4} \quad \zeta\left(\frac{1}{2} \pm it\right) = 0$$

The formula **(3.4)** confirms the relation **(3.2)** and the property **(0.2)** of the non-trivial zeros, stated in the introduction.

All the zeros, known as the non-trivial zeros of the zeta function, derive from this formula.

In conclusion, in the paragraph **3** we have shown that all the non-trivial zeros of the zeta function must be of the form $\frac{1}{2} + it$ so that the relation **(3.2)** and the property **(0.2)**, both satisfied by all the non-trivial zeros, are true and this confirms the veracity of the Riemann hypothesis.

4 Clarifications and Conclusions

This proposal of demonstration of the Riemann hypothesis is based on the fact that since the non-trivial zeros derive only from the factor **(1.5)** and thus they satisfy the condition **(2.2)**, they must be symmetric about the point $P(\frac{1}{2};0)$ in the complex plane. The idea is to consider the non-trivial zeros no more as single points but as couples, which we can deduce from the functional equation **(1)**. The couples $(z_0; z_2)$, $(z_0; z_3)$, $(z_1; z_2)$ and $(z_1; z_3)$ are not valid, because, chosen an initial non-trivial zero, we can't calculate the second through the factor **(1.5)** and, since the non-trivial zeros derive only from this factor and because this couples don't derive from it, $(z_0; z_2)$, $(z_0; z_3)$, $(z_1; z_2)$ and $(z_1; z_3)$ don't derive from functional equation. The annullment of the functional equation, due to the non-trivial zeros through the factor **(1.5)**, includes a domain of definition, regarding the position of these zeros, for which the existence of couples of non-trivial zeros, with imaginary part $\pm it$ and asymmetric about the point $P(\frac{1}{2};0)$, is an absurd, despite the fact that these zeros satisfy the functional equation from a numerical point of view.

The picture **1** could show a valid couple of non-trivial zeros for what we have just said; however, it doesn't satisfy a known property of the non-trivial zeros **(0.2)** for which they must be symmetric about the real line.

The Riemann hypothesis is true, because it is the only case where the non-trivial zeros are symmetric both about the point $P(\frac{1}{2};0)$ and about the real line.

References

- [1] *Wikipedia, Riemann zeta function: https://en.wikipedia.org/wiki/Riemann_zeta_function*
- [2] *Wikipedia, Gamma function: https://en.wikipedia.org/wiki/Gamma_function*