

# The Seiberg-Witten equations for vector fields

A.Balan

October 14, 2018

## Abstract

By analogy with the Seiberg-Witten equations, we define equations for a spinor and a vector field.

## 1 Recalls of differential geometry

The  $Spin - C$ -structures are reductions of a  $SO(n).S^1$ - fiber bundle to the group  $Spin^C(n) = Spin(n) \times_{\{1,-1\}} S^1$ . For a four-manifold it exists always a  $Spin - C$ -structure for the tangent fiber bundle [F].

The Dirac operator is defined over the  $Spin - C$ -structure with help of a connection  $A$  for the associated line bundle.

$$\mathcal{D}_A = \sum_i e_i \cdot \nabla_{e_i}^A$$

with  $\nabla^A$  the connection defined by the Levi-Civita connection and the connection  $A$  of the determinant fiber bundle of the  $Spin - C$ -structure.

The self-dual part of the curvature (which is a 2-form) of the connection  $A$  is considered:

$$\Omega_A^+$$

A self-dual 2-form with imaginary values, bound to a spinor  $\psi \in S^+$  is also defined by [F]:

$$\omega(\psi)(X, Y) = \langle X.Y.\psi, \psi \rangle + \langle X, Y \rangle |\psi|^2$$

## 2 Recalls of the Seiberg-Witten equations

The Seiberg-Witten equations are the following ones [F] [M]:

1)

$$\mathcal{D}_A(\psi) = 0$$

2)

$$\Omega_A^+ = -(1/4)\omega(\psi)$$

### 3 The SWX equations

#### 3.1 Definition

By analogy with the usual Seiberg-Witten equations, we are tempted to define equations for a spinor  $\psi$  and a vector field  $X$ :

1)

$$\mathcal{D}_X(\psi) = (\mathcal{D} + iX)(\psi) = 0$$

2)

$$id(X^*)^+ = -(1/4) \frac{\omega(\psi)}{|\psi|^2}$$

with  $X^*$  the dual form of  $X$ ,  $d$  is the differential of the forms. The first equation makes use of the Clifford multiplication. We call these two equations, the SWX equations.

#### 3.2 The gauge group

The gauge group acts:

$$f.(X, \psi) = (X - (df)^*, if\psi)$$

#### 3.3 The moduli spaces

We verify that we can define the quotient of the solutions of the SWX equations by the gauge group, it is the moduli space.

### References

- [B] N.Berline, E.Getzler, M.Vergne, "Heat kernels and Dirac operators", Springer-Verlag, 1992.
- [F] T.Friedrich, "Dirac Operators in Riemannian Geometry", Graduate Studies in Mathematics vol 25, AMS, 2000.
- [K] M.Karoubi, "Algèbres de Clifford et K-théorie", Ann.Scient.Ec.Norm.Sup. 4 ser. 1 (1968), 161-270.
- [M] J.Morgan, "The Seiberg-Witten equations and applications to the topology of smooth four-manifolds", Mathematical Notes, Princeton University Press, 1996.
- [W] E.Witten, "Monopoles and four-manifolds", Math.Res.Lett. 1(1994), 769-796.