

# The Origin of the Experimental Weinberg Angle

Sylwester Kornowski

**Abstract:** Here, within the non-perturbative Scale-Symmetric Theory (SST), we derived the Paschos-Wolfenstein (PW) relationship. The derivation shows the true origin of the experimental Weinberg angle. We also showed that the PW ratio leads to the CP-violation sector in the PMNS matrix.

## 1. Introduction

The Paschos-Wolfenstein (PW) relationship/ratio  $R^-$  for an isoscalar target (i.e. the isospin is  $T = 0$ , i.e. there is equal number of neutrons and protons) is [1]

$$\begin{aligned} R^- &= (\sigma^{vN}_{NC} - \sigma^{v(\text{anti})N}_{NC}) / (\sigma^{vN}_{CC} - \sigma^{v(\text{anti})N}_{CC}) = \\ &= 1/2 - \sin^2\Theta_w = 1/2 - s_w^2, \end{aligned} \quad (1)$$

where  $\sigma^{vN}_{NC}$  and  $\sigma^{vN}_{CC}$  are the deep inelastic neutrino-nucleon cross sections for neutral-current (NC) and charged-current (CC) interactions while  $\sigma^{v(\text{anti})N}_{NC}$  and  $\sigma^{v(\text{anti})N}_{CC}$ , due to the opposite internal helicity of the muon-antineutrinos and nucleons [2], are the elastic cross sections as it is showed in this paper. The  $\Theta_w$  is the experimental Weinberg angle.

The orthodox electroweak (EW) theory leads to following formula for the weak mixing angle

$$s_w^2 = 1 - (W^+ / Z)^2 = 0.22301(27), \quad (2)$$

where  $W^\pm = 80.379(12)$  GeV and  $Z = 91.1876(21)$  GeV [3] are the orthodox bosons carrying the weak interactions. Emphasize that at this time, there is no generally accepted theory that explains why the experimental value is so and not different.

Here the very simple derivation of the PW ratio on base of the Scale-Symmetric Theory (SST) [2] shows the true origin of the  $s_w^2$ .

SST leads to the atom-like structure of baryons [2]. There are three main parts in nucleons: the spin-1/2 torus/electric-charge  $X^+$  which has left-handed internal helicity ( $\bar{L}$ ) and positive

electric charge ( $E_+$ ) so the signature is  $LE_+$ , the spin-0 central condensate  $Y$  responsible for the weak interactions which is composed of the Einstein-spacetime (ES) components i.e. of the spin-1 neutrino-antineutrino pairs, and the relativistic pion  $W$  (it is not the  $W^\pm$  boson).

On the other hand, the muon-neutrinos have also the left-handed internal helicity ( $L$ ) and negative weak charge ( $W_-$ ) so the signature is  $LW_-$  (see Table 6 in [2]). Two whirls with antiparallel spins on the same straight line and with the same internal helicity attract each other. Moreover, the opposite electric charge and weak charge cause that there is additional attraction. It leads to conclusion that the muon-neutrinos are scattered on the  $X^+$  and such scattering is the deep inelastic scattering because pions are produced inside the torus. According to SST, the muon-antineutrinos have the signature  $RW_+$  so they are repulsed by  $X^+$ . It leads to conclusion that there is the elastic scattering on  $X^+$ . To conserve the spin and electric charge of the torus  $X^+$ , the muon-antineutrinos produce pairs of the spin-1 pairs – it can be the two spin-1 bare electron-positron pairs  $(e^+e^-)_{\text{bare}}$  or two spin-1 neutrino-antineutrino pairs, for example,  $\nu_\mu\nu_{\mu,\text{anti}}$  plus  $\nu_e\nu_{e,\text{anti}}$  – such an object of four neutrinos is perfectly neutral (see Table 6 in [2]).

First the scattered neutrino produces the charged pions  $\pi^{+,-} \rightarrow \pi^0 + \Delta\pi^\pm$  or the bare pairs  $(e^+e^-)_{\text{bare}}$  and next such neutrino is scattered on electron/positron in  $\Delta\pi^\pm = 4.5936(5)$  [3] or on electron/positron in  $(e^+e^-)_{\text{bare}} = 1.020814$  MeV [2].

More and more massive virtual fermions/tori produced in ES, have smaller and smaller sizes so lower and lower cross sections. Mass of torus is inversely proportional to its squared radius whereas cross section is directly proportional to squared radius so cross section is inversely proportional to mass of torus

$$\sigma^{vN} \sim 1 / \Delta\pi^\pm \sim 1 / (e^+e^-)_{\text{bare}} . \quad (3)$$

We showed here also that  $1/R^-$  leads to the SST invariant for the CP-violation sector in the SST PMNS matrix: 3.6018 [4].

## 2. Calculations

In the simplest deep inelastic NC scattering of the muon-neutrinos on the torus  $X^+$  is

$$\nu_\mu + p \rightarrow \nu_\mu + p + \pi^+ + \pi^- . \quad (4)$$

There are two charged pions so we have

$$\sigma^{vN}_{\text{NC}} \rightarrow 1 / (2\Delta\pi^\pm) . \quad (5)$$

In the simplest elastic NC scattering of the muon-antineutrinos on the torus  $X^+$  is

$$\nu_{\mu,\text{anti}} + p \rightarrow \nu_{\mu,\text{anti}} + p , \quad (6)$$

so  $\sigma^{v(\text{anti})N}_{\text{NC}}$  we can neglect.

From (4) – (6) we have

$$\sigma^{vN}_{\text{NC}} - \sigma^{v(\text{anti})N}_{\text{NC}} \rightarrow 1 / (2\Delta\pi^\pm) . \quad (7)$$

The muon is produced from selected particles from the two quadrupoles (the electron quadrupole and neutrino quadrupole). In the simplest deep inelastic CC scattering of the muon-neutrinos on the torus  $X^+$  is

$$\nu_\mu + n + 2(e^+e^-)_{\text{bare}} + (\nu_\mu\nu_{\mu,\text{anti}} + \nu_e\nu_{e,\text{anti}}) \rightarrow \mu^- + (\Sigma^+ \rightarrow n + \pi^+) + \gamma + \nu. \quad (8)$$

There is one charged pion and  $2(e^+e^-)_{\text{bare}}$  on the left side so we have

$$\sigma^{\nu N}_{\text{CC}} \rightarrow 1 / [\Delta\pi^\pm - 2(e^+e^-)_{\text{bare}}]. \quad (9)$$

In the simplest elastic CC scattering of the muon-antineutrinos on the condensate  $X^+$  is

$$\nu_{\mu,\text{anti}} + n \rightarrow \nu_{\mu,\text{anti}} + n, \quad (10)$$

so  $\sigma^{\nu(\text{anti})N}_{\text{CC}}$  we can neglect.

For the CC scattering from (8) – (10) we have

$$\sigma^{\nu N}_{\text{CC}} - \sigma^{\nu(\text{anti})N}_{\text{CC}} \rightarrow 1 / [\Delta\pi^\pm - 2(e^+e^-)_{\text{bare}}]. \quad (11)$$

From (1), (7) and (11) we have

$$\begin{aligned} R^- &= [\Delta\pi^\pm - 2(e^+e^-)_{\text{bare}}] / (2\Delta\pi^\pm) = 1/2 - (e^+e^-)_{\text{bare}} / \Delta\pi^\pm = 1 / 3.6000 = \\ &= 1/2 - s_w^2. \end{aligned} \quad (12)$$

From (12) follows that  $R^- = 1 / 3.6000$  and that  $s_w^2 = (e^+e^-)_{\text{bare}} / \Delta\pi^\pm = 0.22223(3)$ .

### 3. Summary

Presented here derivation leads to the true origin of the experimental  $s_w^2$  – it is the ratio of mass of the bare electron-positron pair to mass distance between the charged and neutral pion. We obtained  $s_w^2 = 0.22223(3)$ . Most important are the internal helicities of neutrinos and nucleons and their respectively weak charges and electric charges described within SST [2].

We can see that the  $1/R^- = 3.6000$  is very close to value of the SST invariant for the CP-violation sector in the SST PMNS matrix 3.6018 (see Fig.7 in [4]). The SST invariant leads to  $s_w^2 = 0.22236$ .

Both SST results are consistent with the SLD result  $s_w^2 = 0.22228(54)$  [3].

It is just a coincidence that the orthodox value (see formula (2)) is close to the two values received in this paper. There is not a dependence of  $s_w^2$  on squared masses of the  $W^\pm$  and  $Z$  bosons and such bosons are the composite particles [5]. Notice that both SST values are lower than the lower limit resulting from the orthodox formula.

### References

- [1] E. A. Paschos and L. Wolfenstein, Phys. Rev. D **7**, 91 (1973)
- [2] Sylwester Kornowski (23 February 2018). ‘‘Foundations of the Scale-Symmetric Physics (Main Article No 1: Particle Physics)’’  
<http://vixra.org/abs/1511.0188>

- [3] M. Tanabashi *et al.* (Particle Data Group).  
Phys. Rev. D **98**, 030001 (2018)  
“10. Electroweak Model and Constraints on New Physics” – Revised March 2018 by J. Erler and A. Freitas.
- [4] Sylwester Kornowski (21 June 2019). “The Relationship Between CKM and PMNS Matrices in the CP-Violation Sectors”  
<http://vixra.org/abs/1901.0226>
- [5] Sylwester Kornowski (7 June 2017). “The Origin of the Z and W Bosons”  
<http://vixra.org/abs/1705.0202>