

Proof that there are no odd perfect numbers

Kouji Takaki

May 08th, 2019

1. Abstract

For y to be a perfect number, if one of the prime factors is p , the exponent of p is an integer $n(n \geq 1)$, the prime factors other than p are $p_1, p_2, p_3, \dots, p_r$ and the even exponent of p_k is q_k ,

$$y/p^n = (1 + p + p^2 + \dots + p^n) \prod_{k=1}^r (1 + p_k + p_k^2 + \dots + p_k^{q_k}) / (2p^n) = \prod_{k=1}^r p_k^{q_k}$$

must be satisfied. Let m and q be non-negative integers,

$$n = 4m + 1$$

$$p = 4q + 1$$

Letting b and c be odd integers, satisfying following expressions,

$$b = \prod_{k=1}^r p_k^{q_k}$$

$$c = \prod_{k=1}^r (1 + p_k + p_k^2 + \dots + p_k^{q_k}) / p^n$$

$$2b = c(p^n + \dots + 1)$$

is established. This is a known content. By the consideration of this research paper, since it turns out that there are no solutions that satisfy the condition of this problem in n order equation of p , we have obtained the conclusion that there are no odd perfect numbers.

2. Introduction

The perfect number is one in which the sum of the divisors other than itself is the same value as itself, and the smallest perfect number is

$$1 + 2 + 3 = 6$$

It is 6. Whether an odd perfect number exists or not is currently an unsolved problem.

3. Proof

An odd perfect number is y , one of them is an odd prime number p , an exponent of p is an integer n ($n \geq 1$). Let $p_1, p_2, p_3, \dots, p_r$ be the odd prime numbers of factors other than p , q_k the index of p_k , and variable a be the sum of product combinations other than prime p .

$$a = \prod_{k=1}^r (1 + p_k + p_k^2 + \dots + p_k^{q_k}) \dots \textcircled{1}$$

The number of terms N of variable a is

$$N = \prod_{k=1}^r (q_k + 1) \dots \textcircled{2}$$

When y is a perfect number,

$$y = a(1 + p + p^2 + \dots + p^n) - y \quad (n > 0)$$

is established.

$$a \sum_{k=0}^n p^k / 2 = y$$

$$a \sum_{k=0}^n p^k / (2p^n) = y/p^n \dots \textcircled{3}$$

3.1. If q_k has at least one odd integer

Letting the number of terms where q_k is an odd integer be a positive integer u , because $y/p^n = \prod_{k=1}^r p_k^{q_k}$ is an odd integer, the denominator on the left side of expression $\textcircled{3}$ has a prime factor 2, from expression $\textcircled{2}$ variable a has more than u prime factor 2 and variable a is an even integer. Therefore $\sum_{k=0}^n p^k$ must be an odd integer, n is an even integer and u is 1.

3.2. When all q_k are even integers

y/p^n is an odd integer, the denominator on the left side of expression $\textcircled{3}$ is an even integer, and since N is an odd integer when q_k are all even integers, variable a is an odd integer. Therefore $\sum_{k=0}^n p^k$ is necessary to include one prime factor 2, $\sum_{k=0}^n p^k \equiv 0 \pmod{2}$ is established, and n must be an odd integer.

From 3.1, 3.2, in order to have an odd perfect number, only one exponent of the prime factor of y must be an odd integer and variable a must be an odd integer. We consider the case of 3.2 below.

In order for y to be a perfect number, the following expression must be established.

$$y/p^n = (1 + p + p^2 + \dots + p^n) \prod_{k=1}^r (1 + p_k + p_k^2 + \dots + p_k^{q_k}) / (2p^n) = \prod_{k=1}^r p_k^{q_k}$$

However, q_1, q_2, \dots, q_r are all even integers.

Here, let b be an integer

$$b = \prod_{k=1}^r p_k^{q_k} \dots \textcircled{4}$$

A following expression is established.

$$y/p^n = a(1 + p + p^2 + \dots + p^n) / (2p^n) = b$$

$$a(p^{n+1} - 1) / (2(p - 1)p^n) = b$$

$$(a - 2b)p^{n+1} + 2bp^n - a = 0 \dots \textcircled{5}$$

Because it is an $n + 1$ order equation of p , the solution of the odd prime p is $n + 1$ at most.

$$(ap - 2bp + 2b)p^n = a$$

Since $ap - 2bp + 2b$ is an odd integer, a/p^n is an odd integer, which is c .

$$ap - 2bp + 2b = c \ (c > 0) \dots \textcircled{6}$$

$$(2b - a)p = 2b - c$$

Since variable a is an odd integer, $2b - a$ is an odd integer and $2b - a \neq 0$

$$p = (2b - c) / (2b - a)$$

Since $n \geq 1$

$$a - c = cp^n - c \geq cp - c > 0$$

$$a > c$$

is.

From equation ⑥

$$2b(p - 1) - (ap - c) = 0$$

$$2b - c(p^{n+1} - 1)/(p - 1) = 0$$

$(p^n + \dots + 1)/2$ is an odd integer, $n = 4m + 1$ is required with m as an integer.

$$2b(p - 1) = c(p^{n+1} - 1)$$

$$2b = c(p^n + \dots + 1)$$

$$2b = c(p + 1)(p^{n-1} + p^{n-3} + \dots + 1) \dots \textcircled{7}$$

b is an odd integer when $p + 1$ is not a multiple of 4. It is necessary that $p - 1$ be a multiple of 4. A positive integer is taken as q .

$$p = 4q + 1$$

is established.

When $p > 1$

$$p^n - 1 < p^n$$

$$(p^n - 1)/(p - 1) < p^n/(p - 1)$$

$$p^{n-1} + \dots + 1 < p^n/(p - 1) \dots \textcircled{8}$$

Since p is an odd prime number satisfying $p = 4q + 1$ and $p \geq 5$

$$p^{n-1} + \dots + 1 < p^n/4$$

$$2b - a = c(p^n + \dots + 1) - cp^n = c(p^{n-1} + \dots + 1)$$

$$2b - a < cp^n/4 = a/4$$

$$2b < 5a/4$$

$$a > 8b/5 \dots \textcircled{9}$$

Let a_k and b_k be integers and if

$$a_k = 1 + p_k + p_k^2 + \dots + p_k^{q_k}, \quad b_k = p_k^{q_k},$$

$$a_k - b_k < b_k/(p_k - 1)$$

$$a_k < b_k p_k/(p_k - 1)$$

$$a = \prod_{k=1}^r a_k < \prod_{k=1}^r b_k p_k/(p_k - 1) = b \prod_{k=1}^r p_k/(p_k - 1)$$

$$a/b < \prod_{k=1}^r p_k/(p_k - 1)$$

When $r = 1$, since $a/b < 3/2$ is established, it becomes inappropriate contrary to inequality ⑨.

From expression ⑦,

$$b = c(p + 1)/2 \times (p^{n-1} + p^{n-3} + \dots + 1)$$

holds. Since $(p + 1)/2$ is the product of only prime numbers of b , let d_k be the index,

$$(p + 1)/2 = \prod_{k=1}^r p_k^{d_k}$$

$$p = 2 \prod_{k=1}^r p_k^{d_k} - 1$$

From $a = cp^n$ and expression ⑦,

$$2bp^n = a(p^n + \dots + 1)$$

$$a(p^n + \dots + 1)/(2bp^n) = 1 \dots (A)$$

When $r = 1$,

$$a = (p_1^{q_1+1} - 1)/(p_1 - 1)$$

$$b = p_1^{q_1}$$

Equation (A) does not hold since there is no odd perfect number when $r = 1$.

Let R be a rational number,

$$R = a(p^n + \dots + 1)/(2bp^n)$$

Let b' be a rational number and let A and B to be an integer,

$$b' = (p_k^{q_k+1} - 1)/(p_k^{q_k}(p_k - 1)) > 1$$

$$A = (p_k^{q_k+1} - 1)/(p_k - 1)$$

$$B = p_k^{q_k}$$

Multiplying R by b', there are both cases that p_k increases p or does not change.

When multiplied by b', the rate of change of R is $Ap^n(p'^n + \dots + 1)/(Bp'^n(p^n + \dots + 1))$, if p after variation is p'. If the rate of change of R is 1,

$$Ap^n(p'^n + \dots + 1)/(Bp'^n(p^n + \dots + 1)) = 1$$

$$Ap^n(p'^n + \dots + 1) = Bp'^n(p^n + \dots + 1)$$

This expression does not hold, since the right side is not a multiple of p when p' > p, and A > B holds when p' = p. Due to this operation, R may be larger or smaller than the original value, since the rate of change of R does not become 1.

From $R \neq 1$ and $a = cp^n$ for some r, also multiplying fractions $b' = A_1/B_1$, $b'' = A_2/B_2$, $\dots b'''\dots' = A_x/B_x$, if $R = 1$ holds finally,

$$, \text{ if } a(p^n + \dots + 1)/(2bp^n) \times A_1p^n(p_1^n + \dots + 1)/(B_1p_1^n(p^n + \dots + 1)) \times A_2p_1^n(p_2^n + \dots + 1)/(B_2p_2^n(p_1^n + \dots + 1)) \dots A_xp_{x-1}^n(p_x^n + \dots + 1)/(B_xp_x^n(p_{x-1}^n + \dots + 1)) = 1$$

$$a/(2b) \times A_1/B_1 \times A_2/B_2 \dots A_x(p_x^n + \dots + 1)/(B_xp_x^n) = 1$$

$$a(p_x^n + \dots + 1)A_1A_2 \dots A_x = 2bp_x^nB_1B_2 \dots B_x$$

$$cp^n(p_x^n + \dots + 1)A_1A_2 \dots A_x = 2bp_x^nB_1B_2 \dots B_x$$

When $p_x > p$, it becomes inconsistent since the right side of this expression does not include p as a factor.

When $p_x = p$,

$$cp^n(p^n + \dots + 1)A_1A_2 \dots A_x = c(p^n + \dots + 1)p^n$$

$$A_1A_2 \dots A_x = 1$$

It becomes contradiction, since this expression is not established. Therefore, $a = cp^n$ holds at one point where $R = 1$.

Assuming that $R = 1$ in some r by multiplying fractions $b' = A_1/B_1$, $b'' = A_2/B_2$, \dots
 $b'''\dots' = A_x/B_x$, if $R = 1$ holds,

$$1 \times A_1 p^n (p_1^n + \dots + 1) / (B_1 p_1^n (p^n + \dots + 1)) \times A_2 p_1^n (p_2^n + \dots + 1) / (B_2 p_2^n (p_1^n + \dots + 1)) \dots A_x p_{x-1}^n (p_x^n + \dots + 1) / (B_x p_x^n (p_{x-1}^n + \dots + 1)) = 1$$

$$A_1 A_2 \dots A_x p^n (p_x^n + \dots + 1) = B_1 B_2 \dots B_x p_x^n (p^n + \dots + 1)$$

When $p_x > p$, it becomes inconsistent since the right side of this expression does not include p as a factor.

When $p_x = p$,

$$A_1 A_2 \dots A_x = B_1 B_2 \dots B_x$$

is established. It becomes contradiction. Therefore, when n is fixed, the number of values of r for which $R = 1$ is one or less.

Assuming that $R = 1$ in some r by multiplying fractions $b' = A_1/B_1$, $b'' = A_2/B_2$, \dots
 $b'''\dots' = A_x/B_x$ and reciprocal of fraction previously multiplied, if $R = 1$ holds. At this time, assuming that n also changes, the change rate when multiplying by A_1/B_1 is

$$A_1 p^n (p_{r+1}^{n_{r+1}} + \dots + 1) / (B_1 p_{r+1}^{n_{r+1}} (p^n + \dots + 1))$$

The rate of change when multiplying $p_{r+2}^{q_{r+2}} / (p_{r+2}^{q_{r+2}} + \dots + 1)$ after this is

$$p_1^{q_1} p_{r+1}^{n_{r+1}} (p_{r+2}^{n_{r+2}} + \dots + 1) / ((p_1^{q_1} + \dots + 1) p_{r+2}^{n_{r+2}} (p_{r+1}^{n_{r+1}} + \dots + 1))$$

$$1 \times A_1 p^n (p_{r+1}^{n_{r+1}} + \dots + 1) / (B_1 p_{r+1}^{n_{r+1}} (p^n + \dots + 1)) \times p_1^{q_1} p_{r+1}^{n_{r+1}} (p_{r+2}^{n_{r+2}} + \dots + 1) / ((p_1^{q_1} + \dots + 1) p_{r+2}^{n_{r+2}} (p_{r+1}^{n_{r+1}} + \dots + 1)) \dots = 1$$

When A_z and B_z are reduced when multiplied by the reciprocal, if the products excluding the reduced variable are expressed as $A_1 A_2 \dots A_x$ and $B_1 B_2 \dots B_x$,

$$A_1 A_2 \dots A_x p^n (p_x^{n_x} + \dots + 1) = B_1 B_2 \dots B_x p_x^{n_x} (p^n + \dots + 1)$$

Since it becomes contradiction like above proof, when n is arbitrary, the number of combinations (a, b, p, n) of solutions for $R = 1$ is one or less.

When $n = 0$,

$$R = a(p^n + \dots + 1) / (2bp^n) = a/2b = c/c = 1$$

holds. There are no solution that satisfies equation (A) in the range of $n > 0$.

Therefore, there are no odd perfect numbers.

4. Complement

From equation ⑤,

$$2bp^n(p-1) = a(p^{n+1}-1)$$

$$2 = a(p^{n+1}-1)/(bp^n(p-1))$$

$$\begin{aligned} 2 &= (p_1^{q_1+1}-1)(p_2^{q_2+1}-1) \dots (p_r^{q_r+1}-1)(p^{n+1}-1) \\ &\quad / (p_1^{q_1}p_2^{q_2} \dots p_r^{q_r}p^n(p_1-1)(p_2-1) \dots (p_r-1)(p-1)) \\ 2(p_1^{q_1+1}-p_1^{q_1})(p_2^{q_2+1}-p_2^{q_2}) \dots (p_r^{q_r+1}-p_r^{q_r})(p^{n+1}-p^n) \\ &= (p_1^{q_1+1}-1)(p_2^{q_2+1}-1) \dots (p_r^{q_r+1}-1)(p^{n+1}-1) \end{aligned}$$

We consider when $r = 2$.

$$(p_1^{q_1+1}-1)(p_2^{q_2+1}-1)(p^{n+1}-1) = 2(p_1^{q_1+1}-p_1^{q_1})(p_2^{q_2+1}-p_2^{q_2})(p^{n+1}-p^n)$$

Let s, t, u be integers,

$$s = p_1^{q_1+1} - 1$$

$$t = p_2^{q_2+1} - 1$$

$$u = p^{n+1} - 1$$

are.

$$stu = 2(p_1^{q_1+1} - 1 - (p_1^{q_1} - 1))(p_2^{q_2+1} - 1 - (p_2^{q_2} - 1))(p^{n+1} - 1 - (p^n - 1))$$

$$stu = 2(s - (s+1)/p_1 + 1)(t - (t+1)/p_2 + 1)(u - (u+1)/p + 1)$$

$$pp_1p_2stu = 2((s+1)p_1 - (s+1))((t+1)p_2 + (t+1))((u+1)p + (u+1))$$

$$pp_1p_2stu = 2(s+1)(p_1-1)(t+1)(p_2-1)(u+1)(p-1)$$

$$stu/((s+1)(t+1)(u+1)) = 2(p_1-1)(p_2-1)(p-1)/(p_1p_2p)$$

Since $stu/((s+1)(t+1)(u+1))$ is a monotonically increasing function for variables

s, t and u , if

$$s \geq 3^{2+1} - 1 = 26, \quad p_1 = 3, \quad q_1 = 2$$

$$t \geq 7^{2+1} - 1 = 342, \quad p_2 = 7, \quad q_2 = 2$$

$$u \geq 5^2 - 1 = 24, \quad p = 5, \quad n = 1$$

holds,

$$stu/((s+1)(t+1)(u+1)) \geq 26 \times 342 \times 24 / (27 \times 343 \times 25) = 7904/8575$$

$$2(p_1-1)(p_2-1)(p-1)/(p_1p_2p) = 2 \times 2 \times 6 \times 4 / (3 \times 7 \times 5) = 32/35$$

Since $stu/((s+1)(t+1)(u+1))$ is limited to 1 when s, t and u are infinite,
 $stu/((s+1)(t+1)(u+1)) < 1$

If $f(p_1, p_2, p) = 2(p_1 - 1)(p_2 - 1)(p - 1)/(p_1 p_2 p)$ holds, it is sufficient to consider a combination where $f(p_1, p_2, p) < 1$.

$$f(3,7,5) = 2 \times 2 \times 6 \times 4 / (3 \times 7 \times 5) = 32/35$$

$$f(3,11,5) = 2 \times 2 \times 10 \times 4 / (3 \times 11 \times 5) = 32/33$$

$$f(3,13,5) = 2 \times 2 \times 12 \times 4 / (3 \times 13 \times 5) = 64/65$$

$$f(3,17,5) = 2 \times 2 \times 16 \times 4 / (3 \times 17 \times 5) = 256/255$$

$$f(3,7,13) = 2 \times 2 \times 6 \times 12 / (3 \times 7 \times 13) = 96/91$$

$$f(3,5,17) = 2 \times 2 \times 4 \times 16 / (3 \times 5 \times 17) = 256/255$$

From the above, when $r = 2$, a combination $(p_1, p_2, p) = (3,7,5), (3,11,5), (3,13,5)$ can be considered.

Let q_k be 2 and $n = 1$, if $g(p_1, p_2, p) = (p_1^3 - 1)(p_2^3 - 1)(p^2 - 1)/(p_1^3 p_2^3 p^2)$,

$$g(3,7,5) = 26 \times 342 \times 24 / (3^3 7^3 5^2) = 7904/8575 > 32/35$$

$$g(3,11,5) = 26 \times 1330 \times 24 / (3^3 11^3 5^2) = 55328/59895$$

$$g(3,13,5) = 26 \times 2196 \times 24 / (3^3 13^3 5^2) = 3904/4225$$

Since the function g is the minimum in the case of $q_k = 2$ and $n = 1$, there is no solution q_k and n when $g > f$, so the case of $(p_1, p_2, p) = (3,7,5)$ becomes unsuitable.

$$\begin{aligned} stu/((s+1)(t+1)(u+1)) &= 2(p_1 - 1)(p_2 - 1)(p - 1)/(p_1 p_2 p) \\ (p_1^{q_1+1} - 1)(p_2^{q_2+1} - 1)(p^{n+1} - 1) &/ (p_1^{q_1+1} p_2^{q_2+1} p^{n+1}) \\ &= 2(p_1 - 1)(p_2 - 1)(p - 1)/(p_1 p_2 p) \end{aligned}$$

If $F(p_1, p_2, p) = (p_1 - 1)(p_2 - 1)(p - 1)/(p_1 p_2 p)$,

$$F(p_1^{q_1+1}, p_2^{q_2+1}, p^{n+1}) = 2F(p_1, p_2, p)$$

5. Acknowledgement

In writing this research document, we asked anonymous reviewers to point out several tens of mistakes. We would like to thank you for giving appropriate guidance and counter-arguments.

6. References

Hiroyuki Kojima "The world is made of prime numbers" Kadokawa Shoten, 2017

Fumio Sairaiji · Kenichi Shimizu "A story that prime is playing" Kodansha, 2015