

Elementary Aspects of the Fundamental Theorem of Algebra

Timothy W. Jones

October 29, 2019

Abstract

We give a sequence of inferences from high school algebra that can be related to the fundamental theorem of algebra (FTA). The sequence builds to an easy proof of this theorem. In passing we mention two proofs given in complex analysis courses; these theorems use Rouché and Liouville. These proofs, although short, require differential and integral calculus for complex variables. They do, however, have simple concepts that can help with a simpler, albeit longer, proof. The proof given here is easy enough for good high school students. It may also pique their curiosity about more advanced complex and real analysis courses.

Introduction

How close can a typical high school algebra student come to understanding the fundamental theorem of algebra? Currently some of the ingredients for a good understanding are present after a typical algebra 1, algebra 2, and pre-calc (or trigonometry) sequence, but the dots aren't connected. Thus students are familiar with quadratics and cubics and general polynomials, as well as Euler's and DeMoivre's formula and theorem; they also are told the fundamental theorem of algebra; but, in no course are they encouraged to see how polynomials might always have roots in the complex numbers, how or why the fundamental theorem of algebra is plausible and might be proved. There are inferences that can be made that suggest that this is a plausible conclusion. Indeed, it is possible to give a proof of this result using high school math, albeit with a couple of unproven but intuitively easy assumptions. That's the main trajectory of this article.

We use proofs and results from *Rudin's Principles of Mathematical Analysis* [4] and Spiegel's *Complex Variables* [5]. For high school mathematics, we reference Blitzer's *Algebra and Trigonometry* [1]. When we wish to stress complex variables we use z and for real variables x .

Review

We know using the quadratic formula [1] that all quadratics can be solved in the complex numbers. So $z^2 + 1 = 0$ is solved by $\pm i$. We also know that some quadratics have graphs that don't show any x-intercepts. z^2 touches the origin and has a double root at $(0, 0)$. It is concave up, meaning it opens up or holds water. When transformed vertically up by 1, its vertex moves away from the origin; it has no real roots. The fundamental theorem of algebra states that all polynomials of degree n with rational coefficients have roots – all n of them in the complex plane. These statements should all be easy enough recollections from high school algebra.

What's missing

What might have escaped a student's attention is that a quadratic like formula for the general degree n polynomial's roots is not given. In addition, reglosses of some results show the plausibility of the FTA. In this section we mention these.

Easy example one

There is hope via a simple observation from a chapter on trigonometry, Blitzer's Chapter 7, Section 5: you can solve any n th degree polynomial of the form

$$z^n + a = 0. \tag{1}$$

The n solutions are the n th roots of

$$z = \sqrt[n]{a} = \sqrt[n]{|a|} \left(\cos \frac{\theta + 2\pi k}{n} + i \sin \frac{\theta + 2\pi k}{n} \right),$$

where $0 \leq k \leq n - 1$ and $a = \cos \theta + i \sin \theta$.

Of course polynomials like (1) are not the rule, but we can discern why polynomials might always be solved by complex numbers. There's at least one example of every degree that can so be solved.

Easy example two

Notice that a quadratic like $ax^2 + bx + c = 0$ has the standard form $a(x - h)^2 + k$; Blitzer Chapter 3, Section 1. This is really another $z^2 + k$, only its transformed a little – moved to the right or left and the shape of the parabola legs are squeezed together or spread apart via the a coefficient; neither deformation changes that it has x-intercepts, roots (or doesn't); see Blitzer's Chapter 2, Section 5 on transformations. When switched to complex variables one knows, via the quadratic formula, roots will exist in the complex plane.

Easy example three

Note that higher powers of the form $(z - a)^3 + b = 0$ will just be up and over transformation of z^3 . Using the binomial theorem (Blitzer's Chapter 11, Section 5), we know the $(z - a)^3$ part will generate all terms in a cubic:

$$(z - a)^3 = z^3 - 3az^2 + 3a^2z - a^3.$$

Are all polynomials just transformations of the base type z^n ? and so just as $z^n = 0$ has $z = 0$ as a root of multiplicity of n so do all polynomials have n roots somewhere. Rouché's theorem (see Spiegel, page 128, problem 19 [5]) will get something close to this result: $z^n = 0$ helps to prove the general n th degree polynomial has n roots.

These examples make the FTA plausible and are already in the high school curriculum. Next, we analyze the situation and evolve a proof of the FTA.

The big picture

Let's dig deeper into complex numbers and functions. A mind experiment is to imagine the $x + iy$ of one complex plane mapping to $u + iv$ of another. Imagine two computer windows and as you drag your mouse around on the left plane there is a corresponding cross-hair on the right window. You would hope you could adjust the mouse pointer so as to find the origin on the right: a kind of sophisticated cat and mouse game. The connection between the two windows is a function from the complex numbers to the complex numbers. For $z^2 + 1 = 0$, left i and $-i$ map to the right window's origin. For $z^n + a = 0$ tracing a circle on the left generates periodic bulls-eyes at the origin on the right. A nice academic question is what properties of the

function make its range the entire complex plane. Hint: in general books on complex analysis treat objects called *entire* functions.

What is an example of a function that doesn't have a root for sure. The constant function, $c(z) = c$, $c \neq 0$ is such an example. What is an example of function that has an absolute value that is bounded? The constant function is also such that $|c(z)| < 2|c|$. The constant function is also continuous. If a polynomial doesn't have a root, then $1/p(z)$ is defined for all z and its range excludes the origin; it is a good function – there is no division by zero. There is a theorem that says if $f(z)$ is continuous so is $1/f(z)$, provided $f(z) \neq 0$ for all z in a given domain. If we could show that $|f(z)|$ is unbounded, that would mean $1/|f(z)|$ is bounded on a certain domain. Putting this all together it is possible that our polynomial with these properties must be a constant, which it isn't. It must have a zero; i.e. the FTA is proven. That's Liouville's theorem (Spiegel, page 125, problem 10) applied to show the FTA. Liouville's theorem requires complex differentiation and integration. We seek an easier approach that is almost within the range of high school algebra – no calculus.

Proofs

Here are a results in problem form which will help you become familiar with the ideas of a proof of the FTA. Do the following for missing cases: linear, quadratic, and cubic polynomials $p(z)$. Assume coefficients can be complex numbers.

1. Show that $|p(z)|$ values go to infinity as $R > 0$, in $z = Rei\theta$, goes to infinity.
2. Show that $|p(z)|$ values can be made less than the absolute value of $p(z)$'s constant term, $|p(0)|$.

Lemma 1.

$$|A + B| \geq |A| - |B| \quad (2)$$

Proof. By the triangle inequality,

$$|A + B + (-B)| \leq |A + B| + |B|$$

and this gives (2). □

Lemma 2.

$$|A + B + C| \geq |A| - |B| - |C| \quad (3)$$

Proof. By Lemma 1,

$$|A + (B + C)| \geq |A| - |B + C|$$

and

$$\geq |A| - (|B| + |C|) = |A| - |B| + |C|$$

and this is

$$\geq |A| - |B| - |C|$$

and this gives (3). \square

Clearly, an induction proof (Blitzer's Chapter 11, Section 4) will yield the general result. One can, of course, simply say that if you start with A and subtract rather than add potentially positive numbers you will decrease its value. I.e. it's kind of obvious.

Theorem 1. $|p(z)|$ can be made as large as we please.

Proof. *Quadratic case:* Let $|z| = R$ and suppose

$$p(z) = a_2 z^2 + a_1 z + a_0.$$

Then

$$|p(R)| = |a_2 R^2 + a_1 R + a_0| \geq \tag{4}$$

$$|a_2| R^2 - |a_1| R - |a_0| = R^2[|a_2| - |a_1| R^{-1} - |a_0| R^{-2}].$$

We've used our lemma. The factor in brackets shrinks to $|a_2|$ with growing R and this implies $|p(z)|$ can be made as large as we please. \square

Of course one could use the end-behavior of real polynomials to make the same conclusion $p(x)$: the left and right tails of all absolute values of real polynomials will go to infinity; see Blitzer, Chapter 3, Section 2. Positive headed to the x-axis bounce off of it and head north, for example. But in this theorem we allow for complex coefficients and arguments (z), so this image can't necessarily be relied upon. The complication of allowing complex coefficients forces the constant reliance on conversion to statements with absolute values. We see this in our lemmas and theorems.

First, we need that the function $re^{i\theta}$ has as its range all of \mathbb{C} . This is not hard to imagine. Using

$$re^{i\theta} = r(\cos \theta + i \sin \theta), \tag{5}$$

we see that any point in the complex plane can be expressed in polar coordinates – that’s it. Blitzter states, page 717, but doesn’t prove (5), but see [2].

As a side note, notice that $i\theta$ in the right hand side of (5) is like a linear function going up and down the imaginary i -axis; this generates a circle in the u - v plane. That is a line in the x - y plane goes to a circle in the u - v . This type of transformation broadens modeling opportunities. It might just prove Fermat’s last theorem [3]. Back to our main thread.

Lemma 3. *For every real non-zero number a there exists a real θ such that $ae^{i\theta} = -|a|$. This is also true for complex a .*

Proof. If $a < 0$ then let $\theta = 0$ and $ae^{0i} = -|a|$. If $a > 0$ then let $\theta = \pi$ and $ae^{\pi i} = -a = -|a|$.

For complex a , we just note $ae^{i\theta} = -|a|$ implies

$$e^{i\theta} = \frac{-|a|}{a}.$$

Not that it is necessary to note, we note $|-|a|/a| = 1$, so r in the polar coordinate representation of this number is 1. \square

Theorem 2. *$|p(z)|$ can be made less than the absolute value of its constant.*

Proof. Cubic case: Let $p(z) = a_0 + a_1z + a_2z^2 + a_3z^3$. We always will have a non-zero constant and here we assume a_1 is the first non-zero coefficient. It could be a_2 or a_3 . The argument won’t change. Then

$$p(re^{i\theta}) = a_0 + a_1re^{i\theta} + a_2r^2e^{2i\theta} + a_3r^3e^{3i\theta}.$$

Using Lemma 3, there exists θ such that $a_1e^{i\theta} = -|a_1|$. So now we have

$$|p(re^{i\theta})| = |a_0 - |a_1|r + a_2r^2e^{2i\theta} + a_3r^3e^{3i\theta}|$$

and taking the absolute value of the first constant term and the terms after $-|a_1|r$ increases the value of the right hand side. So

$$|p(re^{i\theta})| \leq |a_0| - |a_1|r + |a_2|r^2 + |a_3|r^3. \quad (6)$$

We’ve used $r > 0$ and $|e^{ik\theta}| = 1$. Rearranging (6),

$$|p(re^{i\theta})| \leq |a_0| - r\{|a_1| - |a_2|r - |a_3|r^2\}.$$

Now for small enough r the value in the braces is positive, so the right hand side drops below $|a_0|$, as needed. \square

We are now in a position to prove, with a couple of assumptions from advanced mathematics, the FTA. The assumptions are intuitively easy to understand because the results parallel real cases that are easily graphed and comprehended. First, real polynomials (with argument x) are continuous on any closed real interval, $[a, b]$. They will have an absolute minimum and maximum on any such interval. Of course, the interval might have to be widened sufficiently to guarantee you have found such a value for all of \mathbb{R} . The quadratic case makes this clear. Its vertex is (h, k) when it is defined using the standard form, $a(x - h)^2 + k$. If a closed interval is such that $h \in [a, b]$ then its absolute maximum, if $a < 0$ is in this interval; likewise, if $a > 0$, its absolute minimum. We also know that polynomials of higher degree have turning points – points in their domains where they turn from an increasing to a decreasing function or vice-versa. A large enough closed interval will capture any local and hence any absolute finite maximum or minimum. The end behavior allows for maximum and/or minimum's to be $\pm\infty$.

All of these properties hold for $|f(z)|$, z complex, when we consider closed discs of the form $\{z \in \mathbb{C} \mid |z| \leq R\}$. These are discs because $|z| = \sqrt{x^2 + y^2}$ gives a circle of complex points $z = (x, y)$. In more advanced math it is proven that complex polynomials are continuous and will have points corresponding to their maximums and minimums in the equivalent of closed intervals, closed discs. The center of discs we consider is the origin, but the discs could have different centers. Like the real case, we may need to extend our domain, increase the radius of our disc in order to ensure that we capture the z value corresponding to $|f(z)|$ minimum and also that we have a value that gives its minimum. So, for the real case say we have $p(x) = 3(x - 7) + 10$ and we are considering the interval $[-3, 0]$, we know this polynomial has a minimum, so we can say for some $m > 0$, $x_0 \in [-m, m]$ and $p(x_0)$ is the minimum value possible for $p(x)$. We actually know that value is 10. In the same way, if we switch to $p(z) = 3(z - 7) + 10$ and consider $D_0(2)$, the disc with center at the origin and radius 2, can we say $z_0 \in D_0(2)$ and $|p(z_0)|$ is the minimum value of $|p(z)|$ in $D_0(2)$. Well we don't know. What we do know is that if the radius is large enough then this will be true; the continuity of complex polynomials on closed discs tells us this result. Rudin's book on real and some complex analysis gives proofs of this [4].

We can also give the idea by way of contrast. Polynomials are not rational functions. Rational functions, like $r(z) = 1/x$, does not reach its maximum on $[0, \infty)$, for example. It isn't defined at $x = 0$; $r(z) = -1/z$ doesn't reach its minimum. These functions asymptotically approach values,

see Blitzer Chapter 3, Section 5; but there is no z_0 such that $r(z_0) = 0$. In contrast, $|p(z)|$ will have a value z_0 such that $|p(z_0)|$ is its minimum value; its a continuous function on any disc. Lemma 1 says that any polynomial will exceed any value on a disk, so we know $|p(z_0)|$ will hit its minimum. That's the hardest thing to prove, the biggest assumption we make.

One more time. The quadratic $a(x-h)^2 + k$ with $k > 0$ and $a > 0$ has a minimum at k , see Blitzer on this the standard form of a quadratic, Chapter 3, Section 1. Apart from the horizontal shift, this function is $x^2 + k$. We have shown that using complex numbers the absolute value of this polynomial can fall below k , its constant. How far below k can it go? It reaches zero when evaluated at $\pm i\sqrt{k}$. We need a way to build on the assumption that it doesn't reach zero. For a proof by contradiction assume there is a polynomial such that the minimum of its absolute value is not 0. The following proof of the FTA is based on that given in Rudin [4]. We are ready.

Theorem 3. *If $p(z)$ is a polynomial, then there exists $z_0 \in \mathbb{C}$ such that $p(z_0) = 0$.*

Proof. We assume $|p(z)|$ is a continuous function and that its minimum occurs at $z = z_0$. To derive a contradiction, we will assume $|p(z_0)| = \mu \neq 0$. Consider the polynomial defined by

$$Q(z) = \frac{p(z + z_0)}{p(z_0)}.$$

Then the constant of $Q(z)$ is, as it is with all polynomials, given by an evaluation at 0, given by $Q(0)$; $Q(0) = 1$. As $|p(z_0)|$ is the minimum value, all other z values make $|Q(z)| > 1$. But this contradicts Theorem 2. We can't get below this polynomial's constant. \square

Conclusion

The fastest avenue to believing and proving the FTA is to notice that $p(z) = z^n + 1 = 0$ can be solved in \mathbb{C} and this means that \mathbb{C} values drop this function's absolute value below its constant. Show that for all absolute values of polynomials, $p(z)$, there are values, z , in \mathbb{C} such that $|p(z)|$ is less than the absolute value of the polynomial's constant. Note: if a polynomial has no constant, then its terms have a common z factor; $z = 0$ is a root; done. Next, show that with the assumption that the minimum value of $|p(z)| > 0$, there is a polynomial that never goes below its constant, a contradiction.

Does the FTA satisfy all curiosity about polynomial roots? Well not really. We know that roots are in the complex field, but we still don't have a general formula for the solutions to a polynomial of degree n . To really be satisfied, we'd like a formula like the quadratic for such polynomials. In order to achieve this we'd have to limit the complex field. The quadratic formula suggests an avenue. Consider $\mathbb{Q}[\sqrt{b^2 - 4ac}]$ is this a field? Could extending the rationals to include certain roots give us a smaller field than \mathbb{C} to look in? These are some of the puzzles abstract algebra addresses.

References

- [1] R. Blitzer, *Algebra and Trigonometry*, 4th ed., Pearson, Upper Saddle, NJ, 2010.
- [2] T.W. Jones, Euler's Formula Motivated, Trigonometric Derivatives Eased Too, (2019), available at <http://vixra.org/abs/1906.0466>.
- [3] T.W. Jones, Using Scientific Notation To Express Fermat's Last Theorem, (2019), available at <http://vixra.org/abs/1805.0276>.
- [4] W. Rudin, *Principles of Mathematical Analysis*, 3rd ed., McGraw-Hill, New York, 1976.
- [5] M. R. Spiegel and J. Liu, *Complex Variables*, 2nd ed., McGraw-Hill, New York, 1999.