

# Proof of the Riemann hypothesis

Toshiro Takami\*  
mmm82889@yahoo.co.jp

## Abstract

Up to now, I have tried to expand this equation and prove Riemann hypothesis with the equation of cos, sin, but the proof was impossible.

However, I realized that a simple formula before expansion can prove it.

The real value is zero only when the real part of s is 1/2. Non-trivial zeros must always have a real value of zero.

The real part of s being 1/2 is the minimum requirement for s to be a non-trivial zeros.

This is true whether the imaginary value increases or decreases to the limit.

## key words

Riemann hypothesis, non-trivial zeros, 1/2, minimum requirement, to the limit

## 1 introduction

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s) \zeta(1-s) \quad (1)$$

$$\begin{aligned} \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i14.1347\} &= -0.950558 - 0.310547i \\ \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i21.022\} &= -0.904282 + 0.426936i \\ \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i25.0109\} &= -0.784761 - 0.619798i \\ \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i30.4249\} &= -0.475849 + 0.879527i \\ \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i32.9351\} &= -0.410261 - 0.911968i \\ \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i37.58618\} &= -0.832147 + 0.554555i \\ \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i40.91872\} &= -0.917431 + 0.397894i \\ \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i43.32707\} &= -0.275249 - 0.961373i \\ \{2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)\}, \{s = 1/2 + i48.00515\} &= 0.130432 + 0.991457i \end{aligned}$$

---

\*Isahaya-shi, Nagasaki-prefecture, 854-0067 Japan

$$\begin{aligned}
& \left\{ 2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s) \right\}, \{s = 1/2 + i49.77383\} = -0.579292 - 0.81512i \\
& \left\{ 2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s) \right\}, \{s = 1/2 + i52.97032\} = -0.867736 - 0.497025i \\
& \left\{ 2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s) \right\}, \{s = 1/2 + i56.44625\} = -0.752855 + 0.658186i \\
& \left\{ 2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s) \right\}, \{s = 1/2 + i315.4756\} = -0.286121 - 0.958193i \\
& \left\{ 2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s) \right\}, \{s = 1/2 + i1393.4334\} = 0.973556 - 0.228449i \\
& \left\{ 2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s) \right\}, \{s = 1/2 + i74920.8275\} = -0.827399 - 0.561615i
\end{aligned}$$

From the above calculation, in Euler's formula Eq.(1),  $\zeta(s)=0$  ( $s$  is non-trivial zeros) is not from  $2^s \pi^{s-1} \sin\left(\frac{s\pi}{2}\right) \Gamma(1-s)$  but  $\zeta(1-s) = 0$ .

$$\zeta(s) = \zeta(1-s) = 0 \quad (2)$$

At the non-trivial zeros,  $\zeta(s) = \zeta(1-s) = 0$  holds. in this case. Eq.(10)=0, Eq(11)=0, and  $\eta(s) = \eta(1-s) = 0$  holds.

$$\eta(1-s) = (1 - \frac{2}{2^{1-s}})\zeta(1-s) = \frac{2^{1-s} - 2}{2^{1-s}}\zeta(1-s) = \frac{2 - 2^{s+1}}{2}\zeta(1-s) = (1 - 2^s)\zeta(1-s) \quad (3)$$

$$\eta(s) + \frac{2}{2^s}\zeta(s) = \eta(1-s) + \frac{2}{2^{1-s}}\zeta(1-s) \quad (4)$$

$$\eta(s) = \frac{2^s - 2}{2^s} \frac{2^{1-s}}{2^{1-s} - 2} \eta(1-s) = \frac{2^s - 2}{2^s} \zeta(1-s) = (1 - \frac{2}{2^s})\zeta(1-s) = 0 \quad (5)$$

$$\begin{aligned}
& \left\{ \frac{2^s - 2}{2^s} \zeta(1-s) \right\}, \{s = 1/2 + i25.0109\} = 0.0000600703 - 0.0000774542i \\
& \left\{ \frac{2^s - 2}{2^s} \zeta(1-s) \right\}, \{s = 1/2 + i30.4249\} = 2.30973 \times 10^{-7} - 0.0000678699i \\
& \left\{ \frac{2^s - 2}{2^s} \zeta(1-s) \right\}, \{s = 1/2 + i32.9351\} = -9.25931 \times 10^{-6} - 0.000117068i \\
& \left\{ \frac{2^s - 2}{2^s} \zeta(1-s) \right\}, \{s = 1/2 + i37.5862\} = -0.0000437932 - 0.0000195875i \\
& \left\{ \frac{2^s - 2}{2^s} \zeta(1-s) \right\}, \{s = 1/2 + i40.9187\} = -0.0000173311 + 0.0000661198i \\
& \left\{ \frac{2^s - 2}{2^s} \zeta(1-s) \right\}, \{s = 1/2 + i74919.0752\} = -0.0000827382 - 0.000177009i \\
& \left\{ \frac{2^s - 2}{2^s} \zeta(1-s) \right\}, \{s = 1/2 + i74920.8275\} = -0.0000166426 + 8.31396 \times 10^{-6}i
\end{aligned}$$

From the above calculation,  $\eta(s)=0$  ( $s$  is non-trivial zeros).

$$\eta(1-s) = \frac{2^{1-s} - 2}{2^{1-s}} \frac{2^s}{2^s - 2} \eta(s) = \frac{2^{1-s} - 2}{2^{1-s}} \zeta(s) = (1 - \frac{2}{2^{1-s}})\zeta(s) = 0 \quad (6)$$

$$\zeta(s) = \frac{2^s}{2^s - 2} \eta(s) = (\frac{2^s - 2 + 2}{2^s - 2})\eta(s) = (1 + \frac{2}{2^s - 2})\eta(s) \quad (7)$$

$$= (1 + \frac{2}{2^s} \frac{2^s}{2^s - 2})\eta(s) = \eta(s) + \frac{2}{2^s} \zeta(s) = \eta(s) + \frac{2}{2^s} [\eta(s) + \frac{2}{2^s} \zeta(s)] \quad (8)$$

$$= \eta(s) + \frac{2}{2^s} [\eta(s) + \frac{2}{2^s} (\eta(s) + \frac{2}{2^s} \zeta(s))] = \eta(s) + \frac{2}{2^s} \eta(s) + (\frac{2}{2^s})^2 \eta(s) + (\frac{2}{2^s})^3 \zeta(s) \quad (9)$$

$$= \eta(s)[1 + \frac{2}{2^s} + (\frac{2}{2^s})^2] + (\frac{2}{2^s})^3 \zeta(s) \neq \eta(s)[\frac{1 - (\frac{2}{2^s})^k}{1 - \frac{2}{2^s}}] + (\frac{2}{2^s})^{k+1} \zeta(s) \quad (10)$$

when k=2 (If the formula is a geometric sequence and is the same up to the k-th term, = holds.)

$$\neq \eta(s)[\frac{1 - (\frac{2}{2^s})^2}{1 - \frac{2}{2^s}}] + (\frac{2}{2^s})^3 \zeta(s) = (1 - \frac{2}{2^s})\zeta(s)[\frac{1 - (\frac{2}{2^s})^2}{1 - \frac{2}{2^s}}] + (\frac{2}{2^s})^3 \zeta(s) \quad (11)$$

$$= \zeta(s)[1 - (\frac{2}{2^s})^2 + (\frac{2}{2^s})^3] \quad (12)$$

$$\zeta(1-s) = \frac{2^{1-s}}{2^{1-s} - 2} \eta(1-s) = (\frac{2^{1-s} - 2 + 2}{2^{1-s} - 2}) \eta(1-s) = (1 + \frac{2}{2^{1-s} - 2}) \eta(1-s) \quad (13)$$

$$= (1 + \frac{2}{2^{1-s}} \frac{2^{1-s}}{2^{1-s} - 2}) \eta(1-s) = \eta(1-s) + \frac{2}{2^{1-s}} \zeta(1-s) \quad (14)$$

$$= \eta(1-s) + \frac{2}{2^{1-s}} [\eta(1-s) + \frac{2}{2^{1-s}} \zeta(1-s)] = \eta(1-s) + \frac{2}{2^{1-s}} \eta(1-s) + (\frac{2}{2^{1-s}})^2 \zeta(1-s) \quad (15)$$

$$= \eta(1-s) + \frac{2}{2^{1-s}} \eta(1-s) + (\frac{2}{2^{1-s}})^2 [\eta(1-s) + \frac{2}{2^{1-s}} \zeta(1-s)] \quad (16)$$

$$= \eta(1-s) + \frac{2}{2^{1-s}} \eta(1-s) + (\frac{2}{2^{1-s}})^2 \eta(1-s) + (\frac{2}{2^{1-s}})^3 \zeta(1-s) \quad (17)$$

when  $\frac{2}{2^{1-s}} = 2^s$

$$= \eta(1-s) + 2^s \eta(1-s) + (2^s)^2 \eta(1-s) + (2^s)^3 \zeta(1-s) \quad (18)$$

$$= \eta(1-s)[1 + 2^s + (2^s)^2] + (2^s)^3 \zeta(1-s) \quad (19)$$

$$\neq \eta(1-s)[\frac{1 - (2^s)^k}{1 - 2^s}] + (2^s)^{k+1} \zeta(1-s) \quad (20)$$

when k=2

$$\neq \eta(1-s)[\frac{1 - 2^{2s}}{1 - 2^s}] + 2^{3s} \zeta(1-s) \quad (21)$$

$$= \zeta(1-s)(1-2^s)[\frac{1-2^{2s}}{1-2^s}] + 2^{3s}\zeta(1-s) \quad (22)$$

$$= \zeta(1-s)[1-2^{2s}] + 2^{3s}\zeta(1-s) \quad (23)$$

$$= \zeta(1-s)[1-2^{2s}+2^{3s}] \quad (24)$$

from Eq.(12) and Eq.(24)

$$\zeta(s)[1 - (\frac{2}{2^s})^2 + (\frac{2}{2^s})^3] = \zeta(1-s)[1 - 2^{2s} + 2^{3s}] \quad (25)$$

$$\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] = \zeta(1-s)[1 - 2^{2s} + 2^{3s}] \quad (26)$$

## 2 Discussion

Define  $0 < \Re(s) < 1$

from Eq.(26)

$$\begin{aligned} &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i14.1347\} = 0.0000916985i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - 14.1347\} = -0.0000916985i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i15.1347\} = 8.54441i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - 15.1347\} = -8.54441i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i16.1347\} = -3.42962i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - 16.1347\} = 3.42962i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i17.1347\} = 0.999351i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - 17.1347\} = -0.999351i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i18.1347\} = -1.36168i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - 18.1347\} = 1.36168i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i21.022\} = -0.000426108i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - i21.022\} = 0.000426108i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i210000.022\} = -0.338902i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - i210000.022\} = 0.338902i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i210000000.022\} = 7.00022i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - i210000000.022\} = -7.00022i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 + i210000000.022\} = 7.00022i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 1/2 - i210000000.022\} = -7.00022i \\ &\{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.501 + i2000000.022\} = 0.13935 + 21.9799i \end{aligned}$$

$$\begin{aligned}
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.501 - i2000000.022\} = 0.13935 - 21.9799i \\
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.499 + i2000000.022\} = -0.13935 + 21.9799i \\
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.499 - i2000000.022\} = -0.13935 - 21.9799i \\
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.50001 + i2000000.022\} = 0.00139345 + 21.9792i \\
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.50001 - i2000000.022\} = 0.00139345 - 21.9792i \\
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.50000001 + i2000000000.022\} = 1.39345 \times 10^{-6} + 21.9792i \\
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.50000001 - i2000000000.022\} = 1.39345 \times 10^{-6} - 21.9792i \\
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.49999999 + i2000000000.022\} = -1.39345 \times 10^{-6} + 21.9792i \\
& \{\zeta(s)[1 - 2^{2(1-s)} + 2^{3(1-s)}] - \zeta(1-s)[1 - 2^{2s} + 2^{3s}]\}, \{s = 0.49999999 - i2000000000.022\} = -1.39345 \times 10^{-6} - 21.9792i
\end{aligned}$$

As in these examples, when the real part of s is 1/2, the real value is 0, but the imaginary value remains.

Even if s is a non-trivial zero, the imaginary value is close to 0 but not 0.

However, even if the real part of s is 1/2, a real value close to 0 but not 0 is frequently generated.

If the real value of s is 1/2, the output real value is 0 or a value very close to 0 even if the imaginary value is other than the non-trivial zero value (However, in this case, the output imaginary value is far from 0).

**That is, the minimum requirement for the non-trivial zeros is that the real part of s is 1/2.**

That is, the lowest condition in which a non-trivial zero exists is a real part value of 1/2.

That is, a non-trivial zero can have a real part only 1/2.

$$\Re(s) = \frac{1}{2} \tag{27}$$

Proof complete.

### 3 Postscript

These calculations were performed with WolframAlpha.

## References

- [1] B.Riemann.: Über die Anzahl der Primzahlen unter einer gegebenen Grosse, Mon. Not. Berlin Akad pp.671-680, 1859
- [2] John Derbyshire.: Prime Obsession: Bernhard Riemann and The Greatest Unsolved Problem in Mathematics, Joseph Henry Press, 2003
- [3] S.Kurokawa.: Riemann hypothesis, Japan Hyoron Press, 2009
- [4] Marcus du Sautoy.: The Music of The Primes, Zahar Press, 2007
- [5] T.Takami.: Consideration of the Riemann hypothesis, viXra:1905.0546

Please raise the prize money to son and daughter who are still young.