

# Elementary proof of the Syracuse conjecture

by Ahmed Idrissi Bouyahyaoui

\*\*\*

## Abstract :

Variables used in the proof are :

$x$  and  $(x + V)$  are positive integer variables.  $(x + V)$  is the successor of  $x$ .

$V$  is an integer variable of adjustment, at first step  $V_0 = 2$  and  $x_0 = y_0 > 2$ .

Representation :  $x = 2^\alpha * (y)$  and  $V = 2^\beta * (z)$ ,  $\alpha * \beta = 0$ ,  $\alpha$  and  $\beta$  are non negative integer variables,  $y$  and  $z$  are odd integer variables,  $y > 0$ .

The Collatz algorithm  $(3 * + 1)$  is applied **simultaneously** to  $x$  and  $(x + V)$ , so we have for rule 2 -  $(x3 + 1)$  - of the algorithm and adjustment :

$$(x + V) := (2^\alpha * (3 * y + 1)) + (2^\beta * (3 * z) - (2^\alpha - 1)) = 3(2^\alpha * (y) + 2^\beta * (z)) + 1$$

That gives :

$$x := 2^\alpha * (3 * y + 1) = 2^{\alpha'} * (y'), V := 2^\beta * (3 * z) - (2^\alpha - 1) = 2^{\beta'} * (z') \quad (1)$$

We deduce the rule :

$$(x := 2^\alpha * (3 * y + 1)) \wedge (V := 2^\beta * (3 * z) - (2^\alpha - 1)) \implies V < x$$

For  $x := 1 * (y_0)$  and  $V := V_0 = 2 * (1)$ , rule 2 -  $(x3 + 1)$  - of the algorithm gives :

$$(x + V) := (1 * (3 * y_0 + 1)) + (2 * (3) - (1 - 1)) > 0 \implies x + V > 0$$

As  $x + V > 0$  and  $V < x$ , we deduce the rule :

$$(V < x) \wedge (x + V > 0) \implies (0 < x + V < 2 * x)$$

This rule shows that sequence  $S(x_0+2)$  is bounded because it is upper bounded by sequence of Syracuse  $S(2 * x_0) = 2 * S(x_0)$  and lower bounded by 0.

By hypothesis  $S(x_0)$  is a sequence of Syracuse.

The **bounded sequence  $S(x_0+2)$**  and the sequence of Syracuse  $S(2 * x_0) = 2 * S(x_0)$  – **upper bound** - converge to the only trivial cycle : [4, 2, 1].

So by recurrence, every positive integer gives a sequence of Syracuse.

\*\*\*

## Syracuse conjecture (Collatz conjecture)

Algorithm of Collatz :

Let  $x$  a positive integer number.

1 - if  $x$  is even then  $x := x/2$

2 - if  $x$  is odd then  $x := x * 3 + 1$

We repeat 1 - 2 until obtain a cycle (is only cycle ?) or  $x$  tends to infinity.

The symbol  $:=$  means : assign value on right to variable on left.

## Representation of variables :

$x$  and  $(x + V)$  are positive integer variables.  $(x + V)$  is the successor of  $x$ .

$V$  is a variable of adjustment, at first step  $V_0 = 2$  and  $x_0 = y_0 > 2$ .

The variables  $x$  and  $V$  are written in the form :

$x := a^*(y)$  with  $a := 2^\alpha$  and  $V := b^*(z)$  with  $b := 2^\beta$ .

$\alpha$  and  $\beta$  are non negative integer variables, such as  $\alpha * \beta = 0$ .

$y$  and  $z$  are odd integer variables,  $y > 0$ .

$(x + V) := a^*(y) + b^*(z) = 2^{\alpha*}(y) + 2^{\beta*}(z)$  and  $\alpha * \beta = 0$ .

## Application of the Collatz algorithm :

The Collatz algorithm  $(3^* + 1)$  is applied **simultaneously** to  $x$  and  $(x + V)$ .

**The coefficient  $a$  is power of 2, the algorithm is applied to the odd part  $y$  of  $x := a^*(y)$  giving a sequence of Syracuse  $S(x_0)$  and the odd part  $z$  of  $V := b^*(z)$  is multiplied by 3 plus an adjustment.**

In operation  $3^* + 1$ ,  $x := a^*(3*y + 1) = a'^*(y')$ ,  $x$  is increased by  $(a - 1)$  to subtract from  $V$  and we have for  $V$  in  $x + V$  :  $V := b^*(3*z) - (a-1) = b'^*(z')$ .

$a'$  and  $b'$  are power of 2,  $y'$  and  $z'$  are odd integer variables.

So we have the equality

**$a^*(3*y+1) + b^*(3*z) - (a-1) = a^*(3*y) + 1 + b^*(3*z) = 3 * (a^*(y) + b^*(z)) + 1$ ,**

**giving  $3^*(x + V) + 1$ , with  $x$  and  $V$  of before the operation  $3^* + 1$ , according to the rule 2 of the algorithm.**

The rule 2 and adjustment give :

$$x := 2^{\alpha*}(3*y+1), V := 2^{\beta*}(3*z) - (2^\alpha - 1) \quad (2)$$

We deduce the rule:

$$(x := 2^{\alpha*}(3*y + 1)) \wedge (V := 2^{\beta*}(3*z) - (2^\alpha - 1)) \implies V < x$$

**In the line  $a'^*(y') + b'^*(z')$ ,  $a'$  and  $b'$  are divided by  $\gcd(a', b')$  according to the rule 1 of the algorithm.**

If  $\gcd(a', b') = 1$  then division by 2 is deferred and then we have :

$$x := 2^{\alpha'}*(y'), V := 2^{\beta'}*(z') \text{ and } \alpha' * \beta' = 0.$$

## Evaluation of variable of adjustment $V$ :

When  $x$  is multiplied by 3 then  $+ 1$ ,  $V$  is multiplied by 3.

When  $x$  is divided by 2,  $V$  is divided by 2.

When  $x = a(3*y+1)$ ,  $x$  is increased by  $(a - 1)$ ,  $V$  is decreased by  $(a - 1)$ .

**We deduce that  $V$  is always less than  $x$ .**

We deduce the rule :

$$(V < x) \implies (x + V < 2*x)$$

This rule shows that sequence  $S(x_0+2)$  is bounded because it is upper bounded by sequence of Syracuse  $S(2*x_0) = 2*S(x_0)$ .

By hypothesis  $S(x_0)$  is a sequence of Syracuse.

Let  $R(x_0)$  the sequence of Syracuse without context of  $(x + V)$  **(2)**, division by 2 is done and  $R(x_0) \subset S(x_0)$ .

Power of 2 is a neutral factor in evolution of sign of  $(x + V)$ , application of Collatz algorithm **without division by 2** gives with  $x=y_0$  and  $V=V_0=2$  (initial data):

$$\begin{aligned}(x + V) &= 1(y_0) + 2(1) > 0 \\ &= 1(3y_0 + 1) + 2(3) - (1 - 1) > 0 \\ &= 1(3^2y_0 + 3 + 1) + 2(3^2) > 0\end{aligned}$$

.....

This shows  $(x + V)$  is always positive and therefore the sequence  $S(x_0+2)$  is lower bounded by 0 :

$$(x + V) > 0$$

We deduce the rule :

$$(V < x) \wedge (x + V > 0) \implies (0 < x + V < 2*x)$$

**Conclusion :**

The **bounded sequence  $S(x_0+2)$**  and the sequence of Syracuse  $S(2*x_0) = 2*S(x_0)$  – **upper bound** - converge to the only trivial cycle : [4, 2, 1].

So by recurrence, every positive integer gives a sequence of Syracuse.

\*\*

Generation of sequences of Syracuse  $S[x]$  and  $S[x + 2]$  :

Application of Collatz algorithm to generate sequences  $S[x_0]$  :

Generation of sequence of Syracuse  $S[17]$  :  $x_0 = 17$

1(17), 1(52), 4(13), 1(13), 1(40), 8(5), 1(5), 1(16), 16(1), 1(1)

Generation of sequence of Syracuse  $S[19]$  :  $x_0 = 19$

1(19), 1(58), 2(29), 1(29), 1(88), 8(11), 1(11), 1(34), 2(17),  
1(17), 1(52), 4(13), 1(13), 1(40), 8(5), 1(5), 1(16), 16(1), 1(1)

Application of Collatz algorithm simultaneously to  $x$  and  $(x + V)$  to generate sequences of Syracuse  $S[x_0]$  and  $S[x_0 + V_0]$  :

Generation of sequences of Syracuse  $S[17]$  and  $S[17 + 2]=S[19]$  :

$$x_0 = 17 \text{ et } x_0 + V_0 = 17 + 2 = 19$$

$S[17] =$

$S[17 + 2] = S[19] =$

$$1(\mathbf{17}) + 2(1) = 19 = 1(\mathbf{19})$$

$$1(\mathbf{52}) + 2(3) = 58 = 2(\mathbf{29})$$

$$4(\mathbf{13}) + 2(3) = 58 = 2(\mathbf{29})$$

$$2(13) + 1(3) = 29 = 1(\mathbf{29})$$

$$2(\mathbf{40}) + 1(9) - 1 = 88 = 8(\mathbf{11})$$

$$16(\mathbf{5}) + 8(1) = 88 = 8(\mathbf{11})$$

$$2(5) + 1(1) = 11 = 1(\mathbf{11})$$

$$2(\mathbf{16}) + 1(3) - 1 = 34 = 2(\mathbf{17})$$

$$32(\mathbf{1}) + 2(1) = 34 = 2(\mathbf{17})$$

$$16(1) + 1(1) = 17 = 1(\mathbf{17})$$

$$16(4) + 1(3) - 15 = 52 = 4(\mathbf{13})$$

$$64(1) + 4(-3) = 52 = 4(\mathbf{13})$$

$$16(1) + 1(-3) = 13 = 1(\mathbf{13})$$

$$16(4) + 1(-9) - 15 = 40 = 8(\mathbf{5})$$

$$64(1) + 8(-3) = 40 = 8(\mathbf{5})$$

$$8(1) + 1(-3) = 5 = 1(\mathbf{5})$$

$$8(4) + 1(-9) - 7 = 16 = 16(\mathbf{1})$$

$$32(1) + 16(-1) = 16 = 16(\mathbf{1})$$

$$2(1) + 1(-1) = 1 = 1(\mathbf{1})$$

$$2(4) + 1(-3) - 1 = 4 = 4(\mathbf{1})$$

$$8(1) + 4(-1) = 4 = 4(\mathbf{1})$$

$$2(1) + 1(-1) = 1 = 1(\mathbf{1})$$