

1 Nature is something that people do exploring or just enjoy. But
 2 what are people? Humans are the ones who study nature. The last
 3 three sentences are circular reasoning: Nature \rightarrow People \rightarrow Nature.
 4 But the planet has the following disbelief: “Any circular reasoning is a
 5 logical fallacy because it is circular reasoning.” I noticed a tautology
 6 in the last sentence. Because we can’t keep our thoughts open all the
 7 time, they must be linked together. For example, some dictionaries
 8 explain meanings of the words: “A car is a machine that is driven.”
 9 But what is a machine? But what is “driven”? So, circular reasoning
 10 (explanations) arise, and they must occur: God is Mr. Love, and Mr.
 11 Love is God.

12

2. SECOND PROOF

13 The total amount H of prime numbers is infinite:

$$(2) \quad H = \infty .$$

14 Therefore, H cannot be any finite number. This means that $H \neq 1$,
 15 $H \neq 2$, $H \neq 3$, and so on. I see that the number on the right-hand
 16 side grows indefinitely, so I have the right to write the final record:

$$(3) \quad H \neq \infty .$$

17 But recall Eq. (2). Therefore, after inserting this equation into the left-
 18 hand side of Eq. (3), I have $\infty \neq \infty$ and $\infty - \infty \neq 0$. The equations
 19 (2) and (3) are not in mutual contradiction because $\infty - \infty$ is a type
 20 of mathematical uncertainty.

21 A “counter-example” is a situation in which the zero of the zeta
 22 function does not belong to $x = 1/2$. The total number V of such
 23 counter-examples is still unknown but cannot be a finite number [1].
 24 Therefore, $V \neq 1$, $V \neq 2$, $V \neq 3$, and upto infinity:

$$(4) \quad V \neq \infty .$$

25 By inserting the definition of V into the left-hand side of Eq. (4), I am
 26 reading from it: the unknown number of counter-examples cannot be
 27 infinite.

28

3. THIRD PROOF

29 Suppose that Riemann Hypothesis fails. Then [2]

$$(5) \quad \lambda_n \leq \frac{\ln(\ln(N_k^{Y_k}))}{\ln(\ln(n_k))} = \frac{\ln Y_k + \ln(\ln(N_k))}{\ln(\ln(n_k))} ,$$

1 where $N_k = \text{rad}(n_k) \leq n_k$ is the radical of n_k , $Y_k = Y_k(p_k) \geq 1$ is a
 2 function of the largest prime factor of N_k , and

$$(6) \quad \lambda_n = \prod_{i=1}^k \frac{p_i^{a_i+1}}{p_i^{a_i+1} - 1} \geq \frac{p_v^{a_v+1}}{p_v^{a_v+1} - 1} \geq 1,$$

3 where p_i are the prime factors of n_k and a_i are the powers of those.
 4 From Eqs. (5) and (6), one has

$$(7) \quad \frac{N_k^{Y_k}}{n_k} \geq 1.$$

5 Y_k tends to 1, as $p_k \rightarrow \infty$ during $n_k \rightarrow \infty$. The $n_k \geq (N_k)^h$ holds,
 6 where h is defined as a fixed constant, e.g., $h = 1.3$. Therefore, Eq. (7)
 7 will be violated which proves Riemann's Hypothesis.

8 If the only choice for h is $h = 1$, this means that for some n_k one
 9 has $n_k = N_k$, i.e., all $a_i = 1$. The latter contradicts the property of
 10 being p-adic. The p-adic property is seen from Eq. (6). Why? Because
 11 Eq. (5) with $\lambda_n \geq 1$, $Y_k \rightarrow 1$, and $N_k \leq n_k$ means $\lambda_n \rightarrow 1$. The latter
 12 combined with Eq. (6) means that all $a_v \rightarrow \infty$, where $1 \leq v < k$.

13 By the way, the p-adic property implies $p_k \rightarrow \infty$ for $n_k \rightarrow \infty$. Why?
 14 See Eq. (5) with $\lambda_n \rightarrow 1$. The latter means $N_k \rightarrow \infty$ which again
 15 means that $p_k \rightarrow \infty$.

16 4. FOURTH PROOF

17 Let within the first N non-trivial zeroes of the Zeta Function happen
 18 to be X counter-examples, which are the zeroes outside the critical line.
 19 Is known that $X/N = 0$ at the limit $N \rightarrow \infty$ from Ref. [3]. However,
 20 that result has zero importance because any distribution of counter-
 21 example is allowed. For example, none of the counter-examples within
 22 $N < 10^{1000000000000000}$. However, the result must have meaning because
 23 it is based on a logical endeavor. That is only possible if there are none
 24 of the counter-examples at all because the result has the title: "100 %
 25 of the zeros of $\zeta(s)$ are on the critical line."

26 5. FIFTH PROOF

27 The number $N(T) = \Omega(T) + S(T)$ of zeroes of Zeta function has
 28 jumps only when $S(T)$ has a jump $\Delta S(T) = S(T + \delta T) - S(T) = 1$
 29 if $\delta T \rightarrow 0$, see Ref. [4], where $0 < x < 1$, $0 < y \leq T + \delta T$ area was
 30 studied. Therefore, $\Delta N(T) = N(T + \delta T) - N(T) = 1$. However,
 31 there are at least two counter-examples at a given y : $x_0 + iy$ and
 32 $1 - x_0 + iy$ due to Dr. Riemann's original paper (or the introductory

1 part of the Sixth Proof in this paper). But $\Delta N(T) = 1 < 2$. From
 2 this contradiction, there cannot be counter-examples.

3 6. SIXTH PROOF

4 The Dirichlet's Eta and Landau's Xi functions have the same zeroes
 5 $s_0 = x + iy$ as the Zeta function in the critical strip. As well as their
 6 complex-conjugate versions. The Xi function has $\xi(s) = \xi(1-s)$, hence,
 7 $\eta(s_0) = \eta(1-s_0)$. All this means that

$$(8) \quad \sum_{n=1}^{\infty} (-1)^n (z^x - z^{1-x}) \sin(y \ln z) = 0,$$

8 where $z = 1/n$. It is the equation $x = x(y)$. Taking the ν -th order
 9 y -derivative of both sides, I obtain a system where the unknowns are
 10 the derivatives

$$(9) \quad L(\mu) = \frac{d^\mu x}{dy^\mu},$$

11 where $\mu = 1, 2, 3, \dots, \nu$. The necessary condition for all $L(\mu)$ to be
 12 zero is

$$(10) \quad \sum_{n=1}^{\infty} (-1)^n (z^x - z^{1-x}) (\ln z)^\nu \cos(y \ln z) = 0,$$

13 if ν is odd, and

$$(11) \quad \sum_{n=1}^{\infty} (-1)^n (z^x - z^{1-x}) (\ln z)^\nu \sin(y \ln z) = 0,$$

14 if ν is even because if one inserts $L(\mu) = 0$ into the equations, they
 15 do not hold true unless Eqs. (10), (11) are holding. There are infin-
 16 itely many independent equations for the unknown x because $\nu =$
 17 $1, 2, 3, \dots, \infty$. However, the value $x = 1/2$ is the obvious solution of
 18 all these equations. Hence, no other values of x exist.

19 REFERENCES

- 20 [1] Guy Robin, "Grandes valeurs de la fonction somme des diviseurs et hypothèse
 21 de Riemann." *J. Math. pures appl.*, 63(2): 187–213 (1984).
 22 [2] F. Vega, "Robin's criterion on divisibility." *Ramanujan J* 59, 745–755 (2022).
 23 <https://doi.org/10.1007/s11139-022-00574-4>
 24 [3] Shekhar Suman, 100 % of the zeros of $\zeta(s)$ are on the critical line.
 25 <https://doi.org/10.48550/arXiv.2205.00811>
 26 [4] Timothy S. Trudgian, "An improved upper bound for the argument of the
 27 Riemann zeta function on the critical line II." *J. Number Theory*, 134: 280–292
 28 (2014).