

Understanding the basics of final unification with Avogadro number

U. V. S. Seshavatharam¹ & S. Lakshminarayana²

¹Honorary faculty, I-SERVE, Alakapuri, Hyderabad-35, AP, India.

²Dept. of Nuclear Physics, Andhra University, Visakhapatnam-03, AP, India.

Abstract: Physicists have long suggested that the four observed fundamental forces of nature are separate manifestations of what was once a single force at times close to the Big Bang. If so, magnitude of the unified force can be assumed to be equal to (c^4/G) . Strength of any interaction can be defined as the ratio of the operating force magnitude and the magnitude of (c^4/G) . Let the gravitational interaction at black holes be called as 'Schwarzschild interaction'. If strength of Schwarzschild interaction is unity, then weak interaction strength seems to be N_A^2 times less than the Schwarzschild interaction and strong interaction strength seems to be $N_A^{\sqrt{\frac{8}{3}}}$ times less than the Schwarzschild interaction. Based on these concepts and considering the Avogadro number as an absolute and discrete number, basics of final unification can be understood.

Keywords: Avogadro number, Schwarzschild's interaction, Final unification.

1. Introduction

From final unification point of view, it is very much essential to couple the universal gravitational constant with the elementary physical constants. Then only the essence of unification can be understood. So far many scientists proposed many interesting models [1-8]. In this context, the important points to be can be expressed as follows.

- 1) In any inverse square law of force, system is sustained only by means of the central attractive force and it is the root cause of revolving body's angular momentum. If is confirmed that, revolving body's angular momentum is discrete, then it is a clear indication of the discrete nature of the central force acting on the revolving body. If one is willing to think in this direction, the historical mystery of Bohr's discrete atomic structure and discrete angular momentum can be understood.
- 2) Note that, as per the basic concepts of final unification, there exists a fundamental unified force from which all the observed forces emerged. If so, magnitude of the unified force can be assumed to be equal to (c^4/G) . Note that, magnitude of radial inward force acting on any black hole surface [9] is the order of (c^4/G) .

- 3) By considering the squared Avogadro number [10-16] as a characteristic proportionality ratio, the characteristic magnitude of the unified atomic force can be assumed to be (c^4/N_A^2G) .
- 4) Considering the Avogadro number as discrete, (c^4/N_A^2G) can be assumed to be discrete.

In this paper authors reviewed [10-17] their published concepts on final unification and proposed different relations for connecting and fitting the Avogadro number and the Newtonian gravitational constant in a unified approach.

2. The classical limits of force and power

Without considering the current notion of black hole physics, Schwarzschild radius of black hole [9] can be estimated with the characteristic limiting force of magnitude (c^4/G) . The outstanding problem in particle physics today is the inclusion of gravity in a single, unified quantum theory of all the fundamental interactions. Particle physicists have long suggested that the four observed fundamental forces of nature (the gravitational, electromagnetic, weak nuclear and strong nuclear forces) are separate, low energy manifestations of what was once a single force at times close to the Big Bang. It is postulated that as the universe expanded and cooled, this single force gradually broke down into the four separate interactions as observed today. However, unification theories that seek to unify the force of gravity with all the other forces (Theories of Everything) remain elusive, as the gravitational interaction lacks a quantum formulation.

To unify cosmology, quantum mechanics and the four observed fundamental cosmological interactions – certainly a ‘unified force’ is required. In this connection (c^4/G) can be considered as the classical force limit. Similarly (c^5/G) can be considered as the classical power limit. If it is true that c and G are fundamental physical constants in physics, then (c^4/G) and (c^5/G) can also be considered as fundamental compound physical constants. These classical limits are more powerful than the Uncertainty limit. Note that by considering the classical force limit (c^4/G) , the famous Planck mass can be obtained.

2.1. Simple applications of (c^4/G) can be stated as follows

- a) Magnitude of force of attraction or repulsion between any two charged particles never crosses (c^4/G) .
- b) Magnitude of gravitational force of attraction between any two massive bodies never crosses (c^4/G) .
- c) Magnitude of mechanical force on a revolving/rotating body never crosses (c^4/G) .
- d) Magnitude of electromagnetic force on a revolving body never crosses (c^4/G) .

2.2. Simple applications of (c^5/G) can be stated as follows

- a) Mechanical power never crosses (c^5/G)
- b) Electromagnetic power never crosses (c^5/G)
- c) Thermal radiation power never crosses (c^5/G)
- d) Gravitational radiation power never crosses (c^5/G)

3. Understanding the role of in Black hole formation and Planck mass generation

3.1 Schwarzschild radius of a black hole

The 4 basic physical properties of a rotating black hole are its mass, size, angular velocity and temperature. Without going deep into the mathematics of black hole physics in the following subsections an attempt is made to understand and fit the black hole radius and temperature.

In all directions if a force of magnitude (c^4/G) acts on the mass-energy content of the assumed celestial body it approaches a minimum radius of (GM/c^2) in the following way [18,19]. Origin of the force (c^4/G) may be due to self-weight or internal attraction or external compression or a something else.

$$R_{\min} \cong \frac{Mc^2}{(c^4/G)} \cong \frac{GM}{c^2} \quad (1)$$

If no force (of zero magnitude) acts on the mass content M of the assumed massive body, its radius becomes infinity. With reference to the average magnitude of $\left(0, \frac{c^4}{G}\right) \cong \frac{c^4}{2G}$, the presently believed Schwarzschild radius can be obtained as

$$(R)_{ave} \cong \frac{Mc^2}{(c^4/2G)} \cong \frac{2GM}{c^2} \quad (2)$$

This proposal is very simple and seems to be different from the existing concepts of General theory of relativity.

3.2 To derive the Planck mass

So far no theoretical model proposed a derivation for the Planck mass. To derive the Planck mass the following two conditions can be given a chance.

Assuming that gravitational force of attraction between two Planck particles of mass (M_p) separated by a minimum distance (r_{\min}) be,

$$\left[\frac{GM_P M_P}{r_{\min}^2} \right] \equiv \left(\frac{c^4}{G} \right) \quad (3)$$

With reference to wave mechanics, let

$$2\pi r_{\min} \equiv \lambda_p = \left[\frac{h}{c.M_P} \right] \quad (4)$$

Here, λ_p represents the wavelength associated with the Planck mass. With these two assumed conditions Planck mass can be obtained as follows.

$$M_P = \sqrt{\frac{hc}{2\pi G}} \equiv \sqrt{\frac{hc}{G}} \quad (5)$$

3.3 Understanding the strength of any interaction

From above relations it is reasonable to say that,

- 1) If it is true that c and G are fundamental physical constants, then (c^4/G) can be considered as a fundamental compound constant related to a characteristic limiting force.
- 2) Black holes are the ultimate state of matter's geometric structure.
- 3) Magnitude of the operating force at the black hole surface is the order of (c^4/G) .
- 4) Gravitational interaction at black holes can be called as 'Schwarzschild interaction'.
- 5) Strength of 'Schwarzschild interaction' can be assumed to be unity.
- 6) Strength of any other interaction can be defined as the ratio of operating force magnitude and the classical force magnitude (c^4/G) .
- 7) If one is willing to represent the magnitude of the operating force as a fraction of (c^4/G) i.e X times of (c^4/G) , where $X \ll 1$, then

$$\frac{X \text{ times of } (c^4/G)}{(c^4/G)} \equiv X \rightarrow \text{Effective } G \Rightarrow \frac{G}{X} \quad (6)$$

If X is very small, $\frac{1}{X}$ becomes very large. In this way, X can be called as the

strength of interaction. Clearly speaking, strength of any interaction is $\frac{1}{X}$ times less than

the 'Schwarzschild interaction' and effective G becomes $\frac{G}{X}$.

4. Basic concepts on final unification

The following concepts and relations can be given a chance in final unification program.

- 1) With reference to the elementary charge and with mass similar to the Planck mass, a new mass unit can be constructed in the following way.

$$\left. \begin{aligned} (M_S)^\pm &\cong \sqrt{\frac{e^2}{4\pi\epsilon_0 G}} \cong 1.859272 \times 10^{-9} \text{ kg} \\ M_S c^2 &\cong \sqrt{\frac{e^2 c^4}{4\pi\epsilon_0 G}} \cong 1.042975 \times 10^{18} \text{ GeV} \end{aligned} \right\} \quad (7)$$

It can be called as the Stoney mass[20]. It is well known that e, c, G play a vital role in fundamental physics. With these 3 constants space-time curvature concepts at a charged particle surface can be studied. It was first introduced by the physicist George Johnstone Stoney. He is most famous for introducing the term ‘electron’ as the ‘fundamental unit quantity of electricity’. In unification program, with this mass unit and with a suitable proportionality ratio- characteristic mass of any elementary charge can be generated.

- 2) Avogadro number is an absolute discrete number and it is having no units like ‘per mole’.
- 3) Atomic gravitational constant is squared Avogadro number times the Newtonian gravitational constant.
- 4) Similar to the classical force limit (c^4/G) , in atomic system there exists a characteristic quantum force of magnitude:

$$F_X \cong (1/N_A^2)(c^4/G) \cong (c^4/N_A^2 G) \quad (8)$$

And its discrete form is:

$$(F_X)_n \cong [1/(nN_A)^2](c^4/G) \cong (1/n^2)(c^4/N_A^2 G) \cong (c^4/n^2 N_A^2 G) \quad (9)$$

where $n = 1, 2, 3, \dots$

- 5) $(F_X)_n$ is responsible for the observed discrete atomic structure and discrete angular momentum of the revolving electron.
- 6) Each proton will attractive the electron with a characteristic discrete force of magnitude $(F_X)_n$ and thus every electron will have total attractive force of $Z(F_X)_n$ where Z is the proton number of the nucleus.
- 7) If $R_c \approx 1.22 \text{ fm}$ is the characteristic nuclear charge radius, qualitatively and quantitatively it is also noticed that,

$$\left(\frac{c^2 R_c}{2N_A^2 G m_e} \right) \frac{\sqrt{m_p m_e} c^2}{2} \cong 27.18 \text{ eV}. \quad (10)$$

- 8) With reference to the discrete potential energy of electron in hydrogen atom [21,22] this observation can be re-written as,

$$(E_{\text{potential}})_n \cong -\sqrt{\frac{m_p}{m_e}} \frac{(F_X)_n R_c}{4} \cong \sqrt{\frac{m_p}{m_e}} \frac{F_X R_c}{4n^2}. \quad (11)$$

- 9) By considering the centripetal force on the electron, discrete Bohr radii can be expressed as,

$$(a_B)_n \cong 4 \sqrt{\frac{m_e}{m_p}} \left(\frac{e^2}{4\pi\epsilon_0 (F_X)_n R_c} \right) \cong 4n^2 \sqrt{\frac{m_e}{m_p}} \left(\frac{e^2}{4\pi\epsilon_0 F_X R_c} \right) \quad (12)$$

and it is possible to show that,

$$\hbar \cong 2 \left(\frac{m_e}{m_p} \right)^{\frac{1}{4}} \sqrt{\frac{m_e c^2}{F_X R_c}} \left(\frac{e^2}{4\pi\epsilon_0 c} \right) \cong \left(\frac{m_e}{m_p} \right)^{\frac{1}{4}} \sqrt{\frac{4N_A^2 G m_e}{c^2 R_c}} \left(\frac{e^2}{4\pi\epsilon_0 c} \right) \quad (13)$$

$$\left(\frac{1}{\alpha} \right) \cong 2 \left(\frac{m_e}{m_p} \right)^{\frac{1}{4}} \sqrt{\frac{m_e c^2}{F_X R_c}} \cong \left(\frac{m_e}{m_p} \right)^{\frac{1}{4}} \sqrt{\frac{4N_A^2 G m_e}{c^2 R_c}} \quad (14)$$

$$n\hbar \cong 2 \left(\frac{m_e}{m_p} \right)^{\frac{1}{4}} \sqrt{\frac{m_e c^2}{(F_X)_n R_c}} \left(\frac{e^2}{4\pi\epsilon_0 c} \right) \cong 2n \left(\frac{m_e}{m_p} \right)^{\frac{1}{4}} \sqrt{\frac{m_e c^2}{F_X R_c}} \left(\frac{e^2}{4\pi\epsilon_0 c} \right) \quad (15)$$

$$\frac{n\hbar}{2} \cong \left(\frac{m_e}{m_p} \right)^{\frac{1}{4}} \sqrt{\frac{m_e c^2}{(F_X)_n R_c}} \left(\frac{e^2}{4\pi\epsilon_0 c} \right) \cong n \left(\frac{m_e}{m_p} \right)^{\frac{1}{4}} \sqrt{\frac{m_e c^2}{F_X R_c}} \left(\frac{e^2}{4\pi\epsilon_0 c} \right) \quad (16)$$

In this way the historical mystery of the discrete nature of electron's angular momentum can be understood in a unified way.

- 10) Modified super symmetric fermion-boson mass ratio [13-17] is close to $\Psi \cong 2.254$. Presently believed charged W boson is nothing but the top quark boson and there exists a charged weak boson of rest energy close 45600 MeV. Pair of 45600 MeV generates the presently believed Z boson and 45600 MeV and 80500 MeV jointly generates a neutral boson of rest energy 126 GeV [13-17].

4.1 Understanding and fitting the characteristic nuclear radii

It is noticed that,

$$\frac{2(N_A^2 G)m_p}{c^2} \cong 9.0 \times 10^{-7} \text{ m} \cong R_1 \text{ (say)} \quad (17)$$

$$\frac{2Gm_p}{c^2} \cong 2.484 \times 10^{-54} \text{ m} \cong R_2 \text{ (say)} \quad (18)$$

With trial-error method it is noticed that, observed characteristic nuclear radii lie in between

$$(R_1^4 R_2)^{\frac{1}{5}} \cong 2.79 \times 10^{-16} \text{ m} \text{ and } (R_1^5 R_2)^{\frac{1}{6}} \cong 1.073 \times 10^{-14} \text{ m} \quad (19)$$

Thus, to a very good accuracy, it is noticed that,

$$N_A^{\sqrt[8]{3}} \left(\frac{2Gm_p}{c^2} \right) \cong 1.688 \times 10^{-15} \text{ m} \quad (20)$$

This can be compared with the presently believed ‘strong interaction range’. A very surprising observation is that,

$$N_A^{\sqrt[8]{3}} \left(\frac{Gm_p}{c^2} \right) \cong 0.844 \times 10^{-15} \text{ m} \quad (21)$$

This can be compared with the presently believed ‘rms radius’ of proton. This is a discovery and needs further study. From this relation and with reference to the new experimental rms radius of proton [23,24,25] $R_{rms} \cong 0.84184(67)\text{fm}$, magnitude of G can be estimated as follows.

$$G \cong N_A^{-\sqrt[8]{3}} \left(\frac{c^2 R_{rms}}{m_p} \right) \cong 6.65742 \times 10^{-11} \text{ m}^3 \text{kg}^{-1} \text{sec}^{-2} \quad (22)$$

Geometric mean of the above two characteristic lengths can be expressed as follows.

$$R_c \cong N_A^{\sqrt[8]{3}} \left(\frac{\sqrt{2}Gm_p}{c^2} \right) \cong 1.194 \times 10^{-15} \text{ m} \approx 1.2 \text{ fm} \quad (23)$$

This length can be compared with the currently believed ‘nuclear charge radius’ 1.2 fermi.

If strength of Schwarzschild interaction is unity, then weak interaction strength can be considered as N_A^{-2} and strong interaction strength can be considered as $N_A^{-\sqrt[8]{3}}$. With this proposal the currently believed strong coupling constant α_s can be fitted as follows [25,26].

$$\left(\frac{1}{\alpha_s} + 1 \right)^2 \cong \ln \left(N_A^{\sqrt[8]{3}} \right) \rightarrow \frac{1}{\alpha_s} \cong 8.45592 \Rightarrow \alpha_s \cong 0.11826 \quad (24)$$

4.2 Understanding and fitting the nucleon rest masses, nuclear stability and nuclear binding energy

Let us guess that N_A^2 plays a characteristic role in generating the rest mass of electron in the following way.

$$\text{let, } m_e c^2 \cong \beta \sqrt{\frac{e^2 c^4}{4\pi\epsilon_0 N_A^2 G}} \cong \beta \sqrt{\frac{e^2 F_X}{4\pi\epsilon_0}} \quad (25)$$

where β can be called as the electron mass index. It can be estimated as:

$$\beta \cong \sqrt{\frac{4\pi\epsilon_0 N_A^2 G m_e^2}{e^2}} \cong 295.0509223 \quad (26)$$

With this number β , electron, muon and tau rest masses can be fitted with the semi empirical relation.

$$\begin{aligned} (m_{lepton})_n c^2 &\cong \left[\beta^3 + (n^2 \beta)^n \sqrt{N_A} \right]^{\frac{1}{3}} \sqrt{\frac{e^2 F_X}{4\pi\epsilon_0}} \\ &\cong \left[\beta^3 + (n^2 \beta)^n \sqrt{N_A} \right]^{\frac{1}{3}} 0.001731 \text{ MeV} \end{aligned} \quad (27)$$

where $n=0,1,2$. Obtained rest energies are 0.511 MeV, 105.95 MeV and 1777.4 MeV respectively [26]. New heavy charged lepton at $n=3$ may be predicted close to 42262 MeV.

Proceeding further let us define another interesting number in the following way.

$$\gamma \cong \beta - \frac{1}{\alpha} \cong 158.0149232 \quad (28)$$

where γ can be called as the nucleon mass index. If so it is noticed that [26],

$$\begin{aligned} \left(\frac{m_n + m_p}{2m_e} \right) &\cong \sqrt{\frac{1}{\gamma} \cdot \frac{N_A^2}{N_A^{\sqrt[8]{3}}}} \text{ and } \left(\frac{m_n - m_p}{m_e} \right) \cong \ln \sqrt{\gamma} \\ \Rightarrow m_n c^2 &\cong 939.80 \text{ MeV and } m_p c^2 \cong 938.51 \text{ MeV} \end{aligned} \quad (29)$$

Interesting observation connected with proton-nucleon stability can be expressed as follows.

$$A_s \cong 2Z + \frac{Z^2}{\gamma} \cong 2Z + 0.00633Z^2 \quad (30)$$

where Z and A_s represent the ‘proton number and ‘stable nucleon number’ of the atomic nucleus respectively. This is a direct relation compared to the existing stability relation [27]. See the following data. $Z = 21, A_s \cong 44.79$; $Z = 29, A_s \cong 63.32$; $Z = 47, A_s \cong 107.98$; $Z = 67, A_s \cong 168.13$; $Z = 83, A_s \cong 209.60$; $Z = 92, A_s \cong 237.56$ and so on.

The two numbers $\left(\gamma, \frac{1}{\alpha}\right)$ can be fitted with the following semi empirical relation.

$$\left. \begin{aligned} \left(\gamma, \frac{1}{\alpha}\right) &\cong \frac{\beta}{2} \pm 2 \ln\left(\frac{\beta}{2}\right) \pm \frac{1}{2} \\ \rightarrow \gamma &\cong \frac{\beta}{2} + 2 \ln\left(\frac{\beta}{2}\right) + \frac{1}{2} \cong 158.01345 \\ \rightarrow \frac{1}{\alpha} &\cong \frac{\beta}{2} - 2 \ln\left(\frac{\beta}{2}\right) - \frac{1}{2} \cong 137.03746 \end{aligned} \right\} \quad (31)$$

With (γ, α) , the semi empirical mass formula energy coefficients [27,28] can be fitted in the following way.

$$\left. \begin{aligned} \text{Pairing energy coefficient, } a_p &\cong \frac{2m_p c^2}{\gamma} \cong 11.876 \text{ MeV} \\ \text{Asymmetry energy coefficient, } a_a &\cong 2a_p \cong 23.752 \text{ MeV} \\ \text{Coulombic energy coefficient, } a_c &\cong \alpha \sqrt{(m_p c^2)} (a_p) \cong 0.77 \text{ MeV} \\ \text{Surface energy coefficient, } a_s &\cong \left(\frac{a_p + a_a}{2}\right) + 2a_c \cong 19.354 \text{ MeV} \\ \text{Volume energy coefficient, } a_v &\cong \left(\frac{a_p + a_a}{2}\right) - 2a_c \cong 16.274 \text{ MeV} \\ \rightarrow (a_s, a_v) &\cong \left(\frac{a_p + a_a}{2}\right) \pm 2a_c \cong \left(\frac{3a_p}{2}\right) \pm 2a_c \\ \text{thus, } a_p + a_a &\cong a_s + a_v \cong 3a_p \cong 35.627 \text{ MeV} \end{aligned} \right\} \quad (32)$$

5. To fit the rest energy of weakly interacting particles

If $R_c \approx 1.22 \text{ fm}$ it is noticed that,

$$\left(\frac{m_e c^2}{F_X R_c}\right) m_e c^2 \cong 102750 \text{ MeV} \quad (33)$$

It is possible to guess that, there exists a charged weakly interacting fermion of mass 102750 MeV. Based on the modified Super symmetry concepts, in the earlier published papers the authors suggested that fermion-boson mass ratio is 2.254 [17]. If so, bosonic form of 102750 MeV can be expressed as follows. If $R_c \approx 1.22 \text{ fm}$

$$\left(\frac{m_e c^2}{F_X R_c}\right)\left(\frac{m_e c^2}{2.254}\right) \cong 45586 \text{ MeV} \quad (34)$$

Two such oppositely charged bosons join together to form a neutral boson of rest energy 91172 MeV. This can be compared with the presently believed neutral Z boson. In the published paper authors suggested that, presently believed charged W boson can be considered as the super symmetric boson of Top quark. Another interesting observation is that, 45586 MeV boson and Top quark boson join together to form the observed 126 GeV boson. For details see references [12-16],[26].

6. Conclusion

So far no model has been succeeded in coupling and understanding the unified concepts of gravity and atomic interactions. Considering the proposed concepts and relations accurate values of Gravitational constant [29,30] and Avogadro number can be estimated from elementary atomic physical constants. With further research and analysis, different models of final unification can be developed with different proportionality ratios and finally a unified model can be standardized with a refined proportionality ratio.

Acknowledgements

The first author is indebted to professor K. V. Krishna Murthy, Chairman, Institute of Scientific Research on Vedas (I-SERVE), Hyderabad, India and Shri K. V. R. S. Murthy, former scientist IICT (CSIR) Govt. of India, Director, Research and Development, I-SERVE, for their valuable guidance and great support in developing this subject.

References

- [1] P. A. M. Dirac, The cosmological constants. *Nature*, 139, 323, 1937.
- [2] Witten, Edward. Search for a realistic Kaluza-Klein theory. *Nuclear Physics B* 186 (3): 412- 428. 1981
- [3] David Gross, Einstein and the search for Unification. *Current science*, Vol. 89, No. 2005, p 12.
- [4] Abdus Salam. Einstein's Last Dream: The Space -Time Unification of Fundamental Forces, *Physics News*, 12(2):36, June 1981.
- [5] Salam A. and Sivaram C. Strong Gravity Approach to QCD and Confinement. *Mod. Phys. Lett.*, 1993, v. A8(4), 321-326.
- [6] Recami E. Elementary Particles as Micro-Universes, and "Strong Black-holes": A Bi-Scale Approach to Gravitational and Strong Interactions. Preprint NSF-ITP-02-94. posted in the arXives as the e-print physics/0505149, and references therein.
- [7] Hawking S.W. *A Brief History of Time*. Bantam Dell Publishing Group. 1988
- [8] Dine, Michael. *Supersymmetry and String Theory: Beyond the Standard Model*. Cambridge University Press. (2007)
- [9] U. V. S. Seshavatharam and S. Lakshminarayana. On the Scale Independent Evolving Quantum Black Hole Cosmology. *Prespace time journal*, Vol 5, issue 9, pp924-942.
- [10] U. V. S. Seshavatharam and S. Lakshminarayana. Nucleus in Strong nuclear gravity. *Proceedings of the DAE Symp. On Nucl. Phys.* 56: 302, 2011
- [11] U. V. S. Seshavatharam and S. Lakshminarayana, Role of Avogadro number in grand unification. *Hadronic Journal*. Vol-33, No 5, 2010 October. p 513.

- [12] U. V. S. Seshavatharam and S. Lakshminarayana, To confirm the existence of atomic gravitational constant. *Hadronic journal*, Vol-34, No 4, 2011 Aug. p379.
- [13] U. V. S. Seshavatharam and S. Lakshminarayana. Logic Behind the Squared Avogadro Number and SUSY. *International Journal of Applied and Natural Sciences*. Vol. 2, Issue 2, 23-40 (2013)
- [14] U. V. S. Seshavatharam and S. Lakshminarayana. Integral charge SUSY in Strong nuclear gravity. *Proceedings of the DAE Symp. on Nucl. Phys.* 56 (2011) p.842
- [15] U. V. S. Seshavatharam and S. Lakshminarayana. Super Symmetry in Strong and Weak interactions. *Int. J. Mod. Phys. E*, Vol.19, No.2, (2010), p.263-280.
- [16] U. V. S. Seshavatharam and S. Lakshminarayana. SUSY and strong nuclear gravity in (120-160) GeV mass range. *Hadronic journal*, Vol-34, No 3, 2011 June, p.277-300
- [17] U. V. S. Seshavatharam and S. Lakshminarayana. New concepts and semi empirical fittings in understanding SUSY and the four cosmological interactions. *Prespace time journal*, Vol 4, issue 11, pp1027- 1038.
- [18] Roger Penrose. Chandrasekhar, Black Holes, and Singularities. *J. Astrophys. Astr.* (1996) 17, 213-231
- [19] Subrahmanyan Chandrasekhar. On Stars, Their Evolution and Their Stability', Nobel Prize lecture, December 8, 1983.
- [20] G.J. Stoney, On the Physical Units of Nature. *Phil.Mag.* 11 (1881) 381-390.
- [21] N. Bohr. On the Constitution of Atoms and Molecules. (Part-1) *Philos. Mag.* 26, 1913, p 1.
- [22] N. Bohr. On the Constitution of Atoms and Molecules. Systems containing only a Single Nucleus. (Part-2) *Philos. Mag.* 26, 476, 1913
- [23] Geiger H and Marsden E. On a diffuse reaction of the particles. *Proc. Roy. Soc., Ser. A* 82: 495-500, 1909.
- [24] Michael O. Distler et al. The RMS Charge Radius of the Proton and Zemach Moments. *Phys. Lett.B.* 696: 343-347, 2011
- [25] J. Mohr, B.N. Taylor, and D.B. Newell in arXiv:1203.5425 and *Rev. Mod. Phys.* (to be published). <http://pdg.lbl.gov/2013/reviews/rpp2012-rev-phys-constants.pdf>
- [26] K.A. Olive *et al.* (Particle Data Group), *Chin. Phys. C*, 38, 090001 (2014)
- [27] Chowdhury, P.R. et al. Modified Bethe-Weizsacker mass formula with isotonic shift and new driplines. *Mod. Phys. Lett. A*20 (2005) p.1605-1618.
- [28] W.D. Myers and W.J. Swiatecki. Table of Nuclear Masses according to the 1994 Thomas-Fermi Model. LBL-36803.1994.
- [29] Terry Quinn, Harold Parks, Clive Speake and Richard Davis. An uncertain big G.. *Phys.Rev. Lett.* 112.068103. (2013) <http://dx.doi.org/10.1103/PhysRevLett.111.101102>.
- [30] J. B. Fixler; G. T. Foster; J. M. McGuirk; M. A. Kasevich. Atom Interferometer Measurement of the Newtonian Constant of Gravity, *Science* 315 (5808): 74-77, (2007).