

# A proof of Riemann hypothesis

Francisco Moga Moscoso<sup>1</sup>, Marina Moga Lozano<sup>2</sup>

1. Industrial Engineer. Mathematical researcher.
2. Malaga Faculty of Medicine. Malaga University.

**Abstract:** This article is very important because the objective is to contribute to research related to the Riemann hypothesis. In this article, we will prove Riemann Hypothesis by a new novel vision.

Keywords: Riemann Hypothesis, Riemann zeta function.

## Proof I:

Riemann zeta function is defined by the Dirichlet series [1] [2]

$$\zeta = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad s = \sigma + it$$

This function has only simple zeros, called trivial zeros at points  $\sigma = -2v, v = 1, 2, 3, \dots$ . Every non-trivial zero in the function  $\zeta(s)$  are complex numbers which have symmetry property with respect to the real axis  $t = 0$  and to the vertical line  $\sigma = \frac{1}{2}$  and they are on the called critical line  $0 \leq \sigma \leq 1$ . For  $\sigma > 1$  the function  $\zeta(s) \neq 0$ .

The Riemann hypothesis states that all non-trivial zeros of zeta function have real part  $\sigma = \frac{1}{2}$  :[3]

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = 0 \quad s = \sigma + it \Rightarrow \sigma = 1/2$$

For that reason we establish that if an equation  $P(x) = 0$  with real coefficient admits a complex root  $x' = \sigma + it$  of order  $k$ , it also admits, with the same order, its conjugate root  $x'' = \sigma - it$ . [1]

Every infinite complex solution of zeta function of Riemann  $\zeta(s)$  with its conjugated root, can also be used to be the solution of a quadratic equation of real coefficients which admits it as zeros, using the root theorem of Viète:

$$x' = \sigma + it, \quad x'' = \sigma - it$$

$$x' + x'' = (\sigma + it) + (\sigma - it) = 2\sigma = b \implies \sigma = \frac{b}{2}$$

$$x' \cdot x'' = (\sigma + it) \cdot (\sigma - it) = \sigma^2 + t^2 = c$$

$$x^2 - bx + c = 0 \text{ siendo } b = 2\sigma \text{ } c = \sigma^2 + t^2$$

$$x'^2 - 2\sigma x' + \sigma^2 + t^2 = 0 \implies x' = \frac{2\sigma + \sqrt{4\sigma^2 - 4(\sigma^2 + t^2)}}{2} = \sigma + it = s$$

$$x''^2 - 2\sigma x'' + \sigma^2 + t^2 = 0 \implies x'' = \frac{2\sigma - \sqrt{4\sigma^2 - 4(\sigma^2 + t^2)}}{2} = \sigma - it$$

This equation can also be written as:

$$x^2 - bx + c = 0 \quad \text{for} \quad b = x' + x'' \quad \text{and} \quad c = x' \cdot x''$$

$$x'^2 - bx' + c = 0 \quad \text{for} \quad c = x'(b - x') \text{ so the equality}$$

$$\text{belongs and identity } x'^2 - bx' + x'(b - x') \equiv 0$$

so this identity has to be verified whatever the values

might be for  $b$  and  $x'$ . For that reason,  $b=1$  and

$$x' = s = \sigma + it \quad \text{for} \quad \sigma = \frac{b}{2} \text{ so it should be verified for } \sigma = \frac{1}{2}, \text{ so:}$$

$$\left(\frac{1}{2} + it\right)^2 - 1\left(\frac{1}{2} + it\right) + \left(\frac{1}{2} + it\right)\left(\frac{1}{2} - it\right) = 0$$

$$\frac{1}{4} + 2\frac{1}{2}it + i^2t^2 - \frac{1}{2} - it + \frac{1}{4} - i^2t^2 = 0$$

and this must be verified whatever  $t$  might be and for  $\sigma = \frac{1}{2}$

$$\text{so, } x' = s = \frac{1}{2} + it$$

$$x''^2 - bx'' + c = 0 \quad \text{for} \quad c = x''(b - x'') \text{ so the equality}$$

$$\text{belongs and identity } x''^2 - bx'' + x''(b - x'') \equiv 0$$

so this identity has to be verified whatever the values

might be for  $b$  and  $x''$ . For that reason,  $b=1$  and

$$x'' = \sigma - it \quad \text{for} \quad \sigma = \frac{b}{2} \text{ so it should be verified for } \sigma = \frac{1}{2}, \text{ so:}$$

$$\left(\frac{1}{2} - it\right)^2 - 1\left(\frac{1}{2} - it\right) + \left(\frac{1}{2} + it\right)\left(\frac{1}{2} - it\right) = 0$$

$$\frac{1}{4} - 2\frac{1}{2}it + i^2t^2 - \frac{1}{2} + it + \frac{1}{4} - i^2t^2 = 0$$

and this must be verified whatever  $t$  might be and for  $\sigma = \frac{1}{2}$

$$\text{so, } x'' = \frac{1}{2} - it.$$

The identities should be verified for  $b$  and  $b=1$  and that is possible if and only if  $\sigma = \frac{1}{2}$ . So, all the non-trivial zeros in the zeta function of Riemann  $\zeta(s)$  must be on the straight line  $\sigma = \frac{1}{2}$ . If it is not so, the identities contradict their own definition that states that: 'an identity is an equality which is verified whatever the values attributed to letters may be'.

### A proof II:

Beginning with the equation  $x^2 - bx + c = 0$  for  $b = x' + x''$  and  $c = x'x''$

$x'^2 - bx' + c = 0$   $c = x'(b - x')$  for  $b$  and  $x'$ . If  $b=1$  we have the following equation:

$x'^2 - 1x' + x'(1 - x') = 0$  and this expression is an identity  $x'^2 - 1x' + x'(1 - x') \equiv 0$ .

This has to be verified whatever the value of  $x'$  may be, so it has to be also verified for every value  $s = \sigma + it$ , that is for every value of the solution of the Riemann zeta equation.

If we substitute  $x'$  for  $\sigma + it$ , we have the following:

$$(\sigma + it)^2 - 1(\sigma + it) + (\sigma + it)(\sigma - it) \equiv 0$$

$\sigma^2 + 2\sigma it + i^2t^2 - \sigma - it + \sigma^2 - i^2t^2 \equiv 0 \Leftrightarrow i^2t^2 + 2\sigma it + 2\sigma^2 \equiv i^2t^2 + it + \sigma$  and being an identity, both members must be identical, as the definition of an identity. We can have the following system:

$$i^2t^2 - i^2t^2 \equiv 0 \text{ for all } \sigma$$

$$2\sigma it - it \equiv 0 \text{ for } \sigma = \frac{1}{2}$$

$$2\sigma^2 - \sigma \equiv 0 \text{ for } \sigma = 0 \text{ and } \sigma = \frac{1}{2}$$

$\sigma = 0$  does not verify the system, so the system is verified if and only if  $\sigma = \frac{1}{2}$ .

For that reason, we have that  $x' = s = \frac{1}{2} + it$ , as we wanted to demonstrate.

Dedekind function and similar, whose are  $s = \sigma + it$ , have the same demonstration.

**Conflict of interest.** The author declares that there is no conflict of interest regarding the publication of this paper.

### References:

- [1] I. M. Vinográdov, *Enciclopedia de las matemáticas / Encyclopedia of mathematics*, no. v. 5. Grupo Anaya Comercial, 2004.
- [2] E. Bombieri, "Problems of the Millennium : The Riemann Hypothesis," *Millenn. Prize Probl.*, vol. 1, no. 2004, pp. 3–9, 2005.
- [3] A. Bouvier and M. George, *Diccionario de matemáticas*. Akal, 1984.