

Proof of Goldbach conjecture

BY :MOHAMMED ZOHAL

mohammedzohal2016@gmail.com

University Hassan II

Abstract .

For any real number $x > 0$, let $\lfloor x \rfloor$ be the largest integer not exceeding x

and $N_{\lfloor \sqrt{x} \rfloor} = \prod_{p \leq \lfloor \sqrt{x} \rfloor, p \in \mathcal{P}} p$ is the product of all primes not exceeding $\lfloor \sqrt{x} \rfloor$

with \mathcal{P} is the set of primes .

let $2n \geq 4$ a positive integer and $R(2n) = \text{card}\{(p,q) / p+q=2n, p,q \in \mathcal{P}^2\}$ denotes the number of prime couples (p,q) such that $p+q=2n$.

In paper we will prove that there is two constants $C_n = \prod_{p/2n, p \neq 2} \frac{p-1}{p-2}$ and $B(n)$

such that $R(2n) \geq \frac{C_n}{2}(n - \sqrt{2n}) \prod_{p/\frac{N_n}{2}} (1 - \frac{2}{p}) - B(n)$ for any integer $n \geq 2$.

This would help us to prove that the Goldbach conjecture is true .

Introduction .

In the letter sent by Goldbach to Euler in 1742 (Christian, 1742) he stated that “its seems that every odd number greater than 2 can be expressed as the sum of three primes”. As reformulated by Euler, an equivalent form of this conjecture called the “strong” or “binary” Goldbach conjecture states that all positive even integers greater or equal to 4 can be expressed as the sum of two primes which are sometimes called a Goldbach partition. Jorg (2000) and Matti (1993) have verified it up to 4.1014. Chen (1973) has shown that all large enough even numbers are the sum of a prime and the product of at most two primes...

The majority of mathematicians believe that Goldbach’s conjecture is true, especially on statistical considerations ,on the subject we give the proof of Goldbach’s strong conjecture whose veracity is based on a clear and simple approach.

Respectively.

Theorem A .

[The Goldbach conjecture is true .](#)

Lemma 1. For any real number $x > 0$, let $\lfloor x \rfloor$ be the largest integer

not exceeding x and $N_{\lfloor \sqrt{x} \rfloor} = \prod_{p \leq \lfloor \sqrt{x} \rfloor, p \in \mathcal{P}} p$ is the

product of all primes not exceeding $\lfloor \sqrt{x} \rfloor$, with \mathcal{P} is the set of primes

$\mathcal{P} = \{2, 3, 5, 7, \dots\}$ and let $\text{gcd}(a,b)$ denotes the greatest

common divisor of the elements (a,b)

then if $\lfloor \sqrt{x} \rfloor + 1 \leq n \leq x$ and $\text{gcd}(n, N_{\lfloor \sqrt{x} \rfloor}) = 1 \Rightarrow n$ is a prime

Proof of Lemma 1. let $N_{\lfloor \sqrt{x} \rfloor} = \prod_{p \leq \lfloor \sqrt{x} \rfloor, p \in \mathcal{P}} p$.

we suppose that $\text{gcd}(n, N_{\lfloor \sqrt{n} \rfloor}) = 1$.

let d be a prime divisor of $n \implies 1 < d \leq \lfloor \sqrt{x} \rfloor$
 $\implies d/N_{\lfloor \sqrt{x} \rfloor}$
 $\implies \gcd(n, N_z) \neq 1$ **Absurd**
then n is a prime

Lemma 2 . (see [01]) let μ denotes the Mobius function then .

$$\sum_{d'|\gcd(n,d)} \mu(d') = \begin{cases} 1 & \text{if } \gcd(n, d) = 1 \\ 0 & \text{if not} \end{cases}$$

Lemma 3 . (see [01])

let f be a multiplicative function then $\sum_{d|n} f(d)$ is also multiplicative .

Lemma 4 .(see[04])

$$\prod_{p \leq x, p \neq 2} \left(1 - \frac{2}{p}\right) \sim \frac{1}{\log(x)^2}, \text{ for all sufficiently large } x$$

Lemma 5 .(see [05])

Let a, b and c , any given integers and let $ax+by=c$ be a diophantine equation, then $ax+by=c$ has a solution **iff** $\gcd(a,b)|c$.

And if (x_0, y_0) is a particular solution of $ax+by=c$ then there exists an integer k such that $(x_0 + \frac{kb}{\gcd(a,b)}, y_0 - \frac{ka}{\gcd(a,b)})$ is the set of solutions .

Lemma 6.

$$\text{let } N_{\lfloor \sqrt{x} \rfloor} = \prod_{p \leq \lfloor \sqrt{x} \rfloor, p \in \mathcal{P}} p \text{ and } d_1/N_{\lfloor \sqrt{x} \rfloor}.$$

$$\text{then } d_2/N_{\lfloor \sqrt{x} \rfloor}, d_1 \wedge d_2 = 1 \iff d_2/\frac{N_{\lfloor \sqrt{x} \rfloor}}{d_1},$$

Proof of Lemma 6.

$$\text{let } N_{\lfloor \sqrt{x} \rfloor} = \prod_{p \leq \lfloor \sqrt{x} \rfloor, p \in \mathcal{P}} p \text{ and } d_1/N_{\lfloor \sqrt{x} \rfloor}.$$

$$1- \text{ we suppose that } d_2/\frac{N_{\lfloor \sqrt{x} \rfloor}}{d_1}.$$

$$\text{we have } d_2/\frac{N_{\lfloor \sqrt{x} \rfloor}}{d_1} \Rightarrow d_2 d_1/N_{\lfloor \sqrt{x} \rfloor} \Rightarrow d_2/N_{\lfloor \sqrt{x} \rfloor}$$

$$\text{and since } N_{\lfloor \sqrt{x} \rfloor} \text{ is squarefree and } d_1 d_2/N_{\lfloor \sqrt{x} \rfloor} \text{ then } d_1 \wedge d_2 = 1$$

$$\text{this means that } d_2/\frac{N_{\lfloor \sqrt{x} \rfloor}}{d_1} \Rightarrow d_2/N_{\lfloor \sqrt{x} \rfloor} \text{ and } d_1 \wedge d_2 = 1$$

$$2- \text{ we suppose that } d_2/N_{\lfloor \sqrt{x} \rfloor}, d_1 \wedge d_2 = 1 .$$

$$\text{we have } d_2/N_{\lfloor \sqrt{x} \rfloor}, d_1/N_{\lfloor \sqrt{x} \rfloor} \text{ and } d_1 \wedge d_2 = 1 \Rightarrow d_2 d_1/N_{\lfloor \sqrt{x} \rfloor}$$

$$\Rightarrow d_2/\frac{N_{\lfloor \sqrt{x} \rfloor}}{d_1}$$

then from 1 and 2 we obtain the equivalence .

$$d_2/N_{\lfloor\sqrt{x}\rfloor}, d_1 \wedge d_2 = 1 \Leftrightarrow d_2/\frac{N_{\lfloor\sqrt{x}\rfloor}}{d_1}$$

Lemma 7 . (see [06])

Let $\tau(n)$ denotes the number of divisors of n .

for all $\varepsilon > 0$, $\tau(n)=o(n^\varepsilon)$

Proof of Theorem A .

let $n \geq 2$, \mathcal{P} denotes the set of primes $R(2n)=\text{card}\{(p,q) / p+q=2n \quad p,q \in \mathcal{P}^2\}$

denotes the number of couple of primes (p,q) such that $p+q=2n$.

For any real number $x > 0$, let $\lfloor x \rfloor$ be the largest integer not exceeding x .

Let $R'(2n)=\text{card}\{(p,q)/p+q=2n, \lfloor\sqrt{2n}\rfloor < p \leq n, n \leq q < 2n-\lfloor\sqrt{2n}\rfloor, p,q \in \mathcal{P}^2\}$ denotes the

number of couple of primes, (p,q) such that $\lfloor\sqrt{2n}\rfloor < p \leq n, n \leq q < 2n-\lfloor\sqrt{2n}\rfloor$ and $p+q=2n$

and $R''(2n)=\text{card}\{(p,q)/p+q=2n, 1 < p \leq \lfloor\sqrt{2n}\rfloor, 2n-\lfloor\sqrt{2n}\rfloor \leq q < 2n, p,q \in \mathcal{P}^2\}$

from the definitions of $R(2n), R'(2n)$ and $R''(2n)$ we can easily prove that

$$R(2n)=R'(2n)+R''(2n) .$$

let $z=\lfloor\sqrt{2n}\rfloor$ and $N_{\lfloor\sqrt{2n}\rfloor}=N_z=\prod_{p \leq \lfloor\sqrt{x}\rfloor, p \in \mathcal{P}} p$

By **Lemma 1** we have .

$$\begin{aligned} R'(2n) &= \text{card}\{(p,q)/p+q=2n, \lfloor\sqrt{2n}\rfloor < p \leq n, n \leq q < 2n-\lfloor\sqrt{2n}\rfloor, \text{gcd}(p,N_z)=1, \text{gcd}(q,N_z)=1\} \\ &= \sum_{p \wedge N_z=1, z < p \leq n} \sum_{q \wedge N_z=1, n \leq q < 2n-z} \sum_{p+q=2n} 1 \\ &= \sum_{p \wedge N_z=1, q \wedge N_z=1, p+q=2n, z < p \leq n} 1 \end{aligned}$$

We apply **Lemma 2** on $\sum_{p \wedge N_z=1, q \wedge N_z=1, p+q=2n, z < p \leq n} 1$.

$$\begin{aligned} R'(2n) &= \sum_{d_1/p \wedge N_z, d_2/q \wedge N_z, p+q=2n, z < p \leq n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) \\ &= \sum_{d_1/p, d_1/N_z \text{ and } d_2/q, d_2/N_z, p+q=2n, z < p \leq n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) \\ &= \sum_{d_1/N_z, d_2/N_z} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) \sum_{d_1/p, d_2/q, p+q=2n, z < p \leq n} 1 \end{aligned}$$

but we have the equivalence .

$$d_1/p, d_2/p+2n \Leftrightarrow \exists j, k \in \mathcal{N}^{*2} \text{ such that } p=jd_1 \text{ et } p+2n=kd_2$$

$$\begin{aligned} \text{Then } R'(2n) &= \sum_{d_1/N_z, d_2/N_z} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) \sum_{p=jd_1, q=kd_2, p+q=2n, z < p \leq n} 1 \\ &= \sum_{d_1/N_z, d_2/N_z} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) \sum_{jd_1+kd_2=2n, z < p=jd_1 \leq n} 1 \end{aligned}$$

Problem 1 .

if we want to give an explicit formula to $R'(2n)$ we would have to

calculate the sum $\sum_{jd_1+kd_2=2n, z < p=jd_1 \leq n} 1$

In fact we will find that if $\text{gcd}(d_1, d_2)/2n$ then .

$$\sum_{jd_1+kd_2=2n, z < p=jd_1 \leq n} 1 = \frac{2n - \lfloor\sqrt{2n}\rfloor}{d_1 d_2} \text{gcd}(d_1, d_2) + O(1)$$

Proof of Probeme 1.

Remark 1.

We remark that the sum $\sum_{jd_1+kd_2=2n, z < p=jd_1 \leq n} 1$, depends only on diophantine equation $d_1+kd_2=2n$, with j and k are the variables .

1 if the equation $jd_1 + kd_2 = 2n$ has a solution

we set $\delta(j, k) = \{$

0 if not

1 if $\gcd(d_1, d_2)/2n$

based on [Lemma 5](#) we have $\delta(j, k) = \{$

0 if not

$$\begin{aligned} \text{then , } \sum_{jd_1+kd_2=2n, z < p=jd_1 \leq n} 1 &= \sum_{z < p=jd_1 \leq n} \delta(j, k) \\ &= \sum_{z < p=jd_1 \leq n, \gcd(d_1, d_2)/2n} 1 \\ &= \sum_{\substack{z \\ d_1 < j \leq \frac{n}{d_1}, \gcd(d_1, d_2)/2n, j \in \mathcal{N}^*}} 1 \end{aligned}$$

We suppose that $\gcd(d_1, d_2)/2n$.

$$\begin{aligned} \text{By [Lemma 5](#) , we have } \sum_{jd_1+kd_2=2n, z < p=jd_1 \leq n} 1 &= \sum_{\substack{z \\ d_1 < j \leq \frac{n}{d_1}, \gcd(d_1, d_2)/2n, j \in \mathcal{N}^*}} 1 \\ &= \sum_{\substack{z \\ d_1 < j=j_0 + \frac{td_2}{\gcd(d_1, d_2)} \leq \frac{n}{d_1}, j \in \mathcal{N}^*}} 1 \\ &= \sum_{\substack{z \\ d_1 - j_0 < \frac{td_2}{\gcd(d_1, d_2)} \leq \frac{n}{d_1} - j_0, t \in \mathcal{Z}}}} 1 \\ &= \sum_{\substack{z \\ d_1 - j_0 < \frac{td_2}{\gcd(d_1, d_2)} \leq \frac{n}{d_1} - j_0, t \in \mathcal{Z}}} 1 \\ &= \sum_{\substack{z \\ \frac{d_1 - j_0}{d_2} \gcd(d_1, d_2) < t \leq \frac{\frac{n}{d_1} - j_0}{d_2} \gcd(d_1, d_2), t \in \mathcal{Z}}} 1 \\ &= \lfloor \frac{\frac{n}{d_1} - j_0}{d_2} \gcd(d_1, d_2) \rfloor - \lfloor \frac{z - j_0}{d_2} \gcd(d_1, d_2) \rfloor + 1 - 1 \\ &= \lfloor \frac{\frac{n}{d_1} - j_0}{d_2} \gcd(d_1, d_2) \rfloor - \lfloor \frac{z - j_0}{d_2} \gcd(d_1, d_2) \rfloor \\ &= \frac{\frac{n}{d_1} - j_0}{d_2} \gcd(d_1, d_2) - \frac{z - j_0}{d_2} \gcd(d_1, d_2) + O(1) \\ &= \frac{\frac{n}{d_1} - \frac{z}{d_1}}{d_2} \gcd(d_1, d_2) + O(1) \\ &= \frac{n - z}{d_2 d_1} \gcd(d_1, d_2) + O(1) \end{aligned}$$

Then if $\gcd(d_1, d_2)/2n$, we obtain .

$$\sum_{jd_1+kd_2=2n, z < p=jd_1 \leq n} 1 = \frac{n - \lfloor \sqrt{2n} \rfloor}{d_1 d_2} \gcd(d_1, d_2) + O(1)$$

Let us now return to calculate $R'(2n)$.

By [Problem 1](#) we have.

$$\begin{aligned}
R'(2n) &= \sum_{d_1/N_z, d_2/N_z} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) \sum_{j d_1 + k d_2 = 2n, z < p = j d_1 \leq n} 1 \\
&= \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) \left(\frac{n-z}{d_1 d_2} \gcd(d_1, d_2) + O(1) \right) \\
&= \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) \frac{n-z}{d_1 d_2} \gcd(d_1, d_2) + \\
&\quad \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) O(1) \\
&= (n-z) \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \frac{\boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2)}{d_1 d_2} \gcd(d_1, d_2) + \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) O(1)
\end{aligned}$$

Problem 2 . let $\tau(n) = \sum_{d|n} 1$ denotes the number of divisors of n .

then the error term $\sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) O(1)$

is equal to $O(\tau(\text{rad}(2n)))$

Proof of Problem 2 .

we set $F = \{d = d_1 \wedge d_2 / d_1 / N_z, d_2 / N_z \text{ and } d / 2n\}$

then we will obtain .

$$\begin{aligned}
L &= \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) O(1) \\
&= \sum_{d \in F} \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2) = d} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) O(1) \\
&= \sum_{d \in F} \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2) = d} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) O(1) \\
&= \sum_{d \in F} \sum_{d_1/N_z} \boldsymbol{\mu}(d_1) \sum_{d_2/N_z, \gcd(d_1, d_2) = d} \boldsymbol{\mu}(d_2) O(1) \\
&= \sum_{d \in F} \sum_{\frac{d_1}{d} / \frac{N_z}{d}} \boldsymbol{\mu}(d_1) \sum_{\frac{d_2}{d} / \frac{N_z}{d}, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right) = 1} \boldsymbol{\mu}(d_2) O(1)
\end{aligned}$$

Let $|x|$ denotes the absolute value of x , then we have.

$$\begin{aligned}
\sum_{d \in F} \sum_{\frac{d_1}{d} / \frac{N_z}{d}} \boldsymbol{\mu}(d_1) \sum_{\frac{d_2}{d} / \frac{N_z}{d}, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right) = 1} \boldsymbol{\mu}(d_2) &\leq \left| \sum_{d \in F} \sum_{\frac{d_1}{d} / \frac{N_z}{d}} \boldsymbol{\mu}(d_1) \sum_{\frac{d_2}{d} / \frac{N_z}{d}, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right) = 1} \boldsymbol{\mu}(d_2) \right| \\
&\leq \sum_{d \in F} \left| \sum_{\frac{d_1}{d} / \frac{N_z}{d}} \boldsymbol{\mu}(d_1) \sum_{\frac{d_2}{d} / \frac{N_z}{d}, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right) = 1} \boldsymbol{\mu}(d_2) \right| \\
&\leq \sum_{d \in F} \sum_{\frac{d_1}{d} / \frac{N_z}{d}} |\boldsymbol{\mu}(d_1)| \left| \sum_{\frac{d_2}{d} / \frac{N_z}{d}, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right) = 1} \boldsymbol{\mu}(d_2) \right| \\
&\leq \sum_{d \in F} \sum_{\frac{d_1}{d} / \frac{N_z}{d}} \left| \sum_{\frac{d_2}{d} / \frac{N_z}{d}, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right) = 1} \boldsymbol{\mu}(d_2) \right|
\end{aligned}$$

By [Lemma 6](#) , we have $\sum_{\frac{d_2}{d} / \frac{N_z}{d}, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right) = 1} \boldsymbol{\mu}(d_2) = \sum_{\frac{d_2}{d} / \frac{N_z}{d}} \boldsymbol{\mu}(d_2)$.

$$\begin{aligned}
\text{Then } \sum_{d \in F} \sum_{\frac{d_1}{d} / \frac{N_z}{d}} \boldsymbol{\mu}(d_1) \sum_{\frac{d_2}{d} / \frac{N_z}{d}, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right) = 1} \boldsymbol{\mu}(d_2) &\leq \sum_{d \in F} \sum_{\frac{d_1}{d} / \frac{N_z}{d}} \left| \sum_{\frac{d_2}{d} / \frac{N_z}{d}} \boldsymbol{\mu}(d_2) \right| \\
&\leq \sum_{d \in F} \sum_{\frac{d_1}{d} / \frac{N_z}{d}} \left| \sum_{\frac{d_2}{d} / \frac{N_z}{d}} \boldsymbol{\mu}\left(\frac{d_2}{d} d\right) \right|
\end{aligned}$$

since d_2 is a squarefree then , $\gcd(\frac{d_2}{d}, d)=1$ then , $\boldsymbol{\mu}\left(\frac{d_2}{d}d\right)=\boldsymbol{\mu}(d)\boldsymbol{\mu}\left(\frac{d_2}{d}\right)$

$$\begin{aligned} \text{Then } \sum_{d \in F} \sum_{\frac{d_1}{d}/\frac{N_z}{d}} \boldsymbol{\mu}(d_1) \sum_{\frac{d_2}{d}/\frac{N_z}{d}, \gcd(\frac{d_1}{d}, \frac{d_2}{d})=1} \boldsymbol{\mu}(d_2) &\leq \sum_{d \in F} \sum_{\frac{d_1}{d}/\frac{N_z}{d}} \left| \sum_{\frac{d_2}{d}/\frac{N_z}{d}} \boldsymbol{\mu}\left(\frac{d_2}{d}\right) \boldsymbol{\mu}(d) \right| \\ &\leq \sum_{d \in F} \sum_{\frac{d_1}{d}/\frac{N_z}{d}} \left| \sum_{\frac{d_2}{d}/\frac{N_z}{d}} \boldsymbol{\mu}\left(\frac{d_2}{d}\right) \right| \end{aligned}$$

By [Lemma 2](#) we have $\sum_{\frac{d_1}{d}/\frac{N_z}{d}} \left| \sum_{\frac{d_2}{d}/\frac{N_z}{d}} \boldsymbol{\mu}\left(\frac{d_2}{d}\right) \right| = \left| \boldsymbol{\mu}\left(\frac{N_z}{d}\right) \right|$

Let $d \in F$,then by the definition of $F = \{d=d_1 \wedge d_2/, d_1/N_z, d_2/N_z, d/2n\}$,
 d will be a squarefree ,then $\frac{N_z}{d}$ is also a squarefree

$$\text{Then } , \sum_{\frac{d_1}{d}/\frac{N_z}{d}} \left| \sum_{\frac{d_2}{d}/\frac{N_z}{d}} \boldsymbol{\mu}(d_2) \right| = \left| \boldsymbol{\mu}\left(\frac{N_z}{d}\right) \right| = 1$$

$$\text{Then, } \sum_{d \in F} \sum_{\frac{d_1}{d}/\frac{N_z}{d}} \boldsymbol{\mu}(d_1) \sum_{\frac{d_2}{d}/\frac{N_z}{d}, \gcd(\frac{d_1}{d}, \frac{d_2}{d})=1} \boldsymbol{\mu}(d_2) \leq \sum_{d \in F} 1$$

Remark 2.

$$\begin{aligned} \text{since } N_z \text{ is squarefree then , } F &= \{d=d_1 \wedge d_2/, d_1/N_z, d_2/N_z, d/\text{rad}(2n)\} \\ &= \{d/\text{rad}(2n) / d \leq \lfloor \sqrt{2n} \rfloor\} \end{aligned}$$

$$\begin{aligned} \text{We have } \tau(\text{rad}(2n)) &= \text{card}\{d/\text{rad}(2n)\} \\ &= \text{card}\{\{d/\text{rad}(2n) / d \leq \lfloor \sqrt{2n} \rfloor\} \cup \{d/\text{rad}(2n) / d \geq \lfloor \sqrt{x} \rfloor\}\} \\ &= \text{card}\{F \cup \{d/\text{rad}(2n) / d \geq \lfloor \sqrt{x} \rfloor\}\} \end{aligned}$$

this means that , $F \subset \{d/\text{rad}(2n)\}$

then we can deduce that, $\text{rad}(F) \leq \tau(\text{rad}(2n))$

We have $\sum_{d \in F} 1 = \text{rad}(F)$,

By [Remark 2](#) , we have $\text{rad}(F) \leq \tau(\text{rad}(2n))$

Then , $\sum_{d \in F} 1 \leq \tau(\text{rad}(2n))$

$$\text{Then } , \sum_{d \in F} \sum_{\frac{d_1}{d}/\frac{N_z}{d}} \boldsymbol{\mu}(d_1) \sum_{\frac{d_2}{d}/\frac{N_z}{d}, \gcd(\frac{d_1}{d}, \frac{d_2}{d})=1} \boldsymbol{\mu}(d_2) \leq \tau(\text{rad}(2n))$$

Which means that $\sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \boldsymbol{\mu}(d_1) \boldsymbol{\mu}(d_2) O(1) = O(\tau(\text{rad}(2n)))$

Result 1.

The error term of $R'(2n)$ is equal to $O(\tau(\text{rad}(2n)))$, this is the most important result in this paper ,because in the sections we will prove that the main part of $R'(2n)$ is much more greater than the error term $O(\tau(\text{rad}(2n)))$

Let us return to calculate $R'(2n)$.

$$R'(2n) = (n-z) \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \frac{\mu(d_1)\mu(d_2)}{d_1 d_2} \gcd(d_1, d_2) + \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \mu(d_1)\mu(d_2) O(1)$$

By the [Problem 2](#) we have .

$$\begin{aligned} R'(2n) &= (n-z) \sum_{d_1/N_z, d_2/N_z, \gcd(d_1, d_2)/2n} \frac{\mu(d_1)\mu(d_2)}{d_1 d_2} \gcd(d_1, d_2) + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d_1/N_z} \frac{\mu(d_1)}{d_1} \sum_{d_2/N_z, \gcd(d_1, d_2)/2n} \frac{\mu(d_2)}{d_2} \gcd(d_1, d_2) + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d \in F} \sum_{d_1/N_z} \frac{\mu(d_1)}{d_1} \sum_{d_2/N_z, \gcd(d_1, d_2)=d} \frac{\mu(d_2)}{d_2} d + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d \in F} \sum_{d_1/N_z} \frac{\mu(d_1)}{d_1} \sum_{d_2/N_z, \gcd(d_1, d_2)=d} \frac{\mu(d_2)}{d_2} d + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d \in F} \sum_{\frac{d_1}{d}/N_z} \frac{\mu\left(\frac{d_1}{d}\right)}{\frac{d_1}{d}} \sum_{\frac{d_2}{d}/N_z, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right)=1} \frac{\mu\left(\frac{d_2}{d}\right)}{\frac{d_2}{d}} d + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d \in F} \frac{1}{d} \sum_{\frac{d_1}{d}/N_z} \frac{\mu\left(\frac{d_1}{d}\right)}{\frac{d_1}{d}} \sum_{\frac{d_2}{d}/N_z, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right)=1} \frac{\mu\left(\frac{d_2}{d}\right)}{\frac{d_2}{d}} + O(\tau(\text{rad}(2n))) \end{aligned}$$

since $\gcd\left(\frac{d_1}{d}, d\right) = 1$ and $\gcd\left(\frac{d_2}{d}, d\right) = 1$ then $\mu\left(\frac{d_1}{d}\right) = \mu(d)\mu\left(\frac{d_1}{d}\right)$ and

$$\mu\left(\frac{d_2}{d}\right) = \mu(d)\mu\left(\frac{d_2}{d}\right).$$

$$R'(2n) = (n-z) \sum_{d \in F} \frac{1}{d} \sum_{\frac{d_1}{d}/N_z} \frac{\mu\left(\frac{d_1}{d}\right)}{\frac{d_1}{d}} \sum_{\frac{d_2}{d}/N_z, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right)=1} \frac{\mu\left(\frac{d_2}{d}\right)}{\frac{d_2}{d}} \mu(d)^2 + O(\tau(\text{rad}(2n)))$$

since d is a squarefree then $\mu(d)^2 = 1$.

$$R'(2n) = (n-z) \sum_{d \in F} \frac{1}{d} \sum_{\frac{d_1}{d}/N_z} \frac{\mu\left(\frac{d_1}{d}\right)}{\frac{d_1}{d}} \sum_{\frac{d_2}{d}/N_z, \gcd\left(\frac{d_1}{d}, \frac{d_2}{d}\right)=1} \frac{\mu\left(\frac{d_2}{d}\right)}{\frac{d_2}{d}} + O(\tau(\text{rad}(2n)))$$

By [Lemma 6](#) we have .

$$R'(2n) = (n-z) \sum_{d \in F} \frac{1}{d} \sum_{\frac{d_1}{d}/N_z} \frac{\mu\left(\frac{d_1}{d}\right)}{\frac{d_1}{d}} \sum_{\frac{d_2}{d}/\frac{N_z}{\frac{d_1}{d}}} \frac{\mu\left(\frac{d_2}{d}\right)}{\frac{d_2}{d}} + O(\tau(\text{rad}(2n)))$$

Since $\frac{\mu\left(\frac{d_2}{d}\right)}{\frac{d_2}{d}}$ is multiplicative then by [Lemma 3](#), $\sum_{\frac{d_2}{d}/\frac{N_z}{\frac{d_1}{d}}} \frac{\mu\left(\frac{d_2}{d}\right)}{\frac{d_2}{d}}$ is also multiplicative

$$\begin{aligned} \text{and } \sum_{\frac{d_2}{d}/\frac{N_z}{\frac{d_1}{d}}} \frac{\mu\left(\frac{d_2}{d}\right)}{\frac{d_2}{d}} &= \prod_{p/\frac{N_z}{\frac{d_1}{d}}} \left(1 - \frac{1}{p}\right) \\ &= \frac{\prod_{p/N_z} \left(1 - \frac{1}{p}\right)}{\prod_{p/\frac{d_1}{d}} \left(1 - \frac{1}{p}\right)} \end{aligned}$$

$$\begin{aligned} R'(2n) &= (n-z) \sum_{d \in F} \frac{1}{d} \sum_{\frac{d_1}{d}/N_z} \frac{\mu\left(\frac{d_1}{d}\right)}{\frac{d_1}{d}} \frac{\prod_{p/N_z} \left(1 - \frac{1}{p}\right)}{\prod_{p/\frac{d_1}{d}} \left(1 - \frac{1}{p}\right)} + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d \in F} \frac{\prod_{p/N_z} \left(1 - \frac{1}{p}\right)}{d} \sum_{\frac{d_1}{d}/N_z} \frac{\mu\left(\frac{d_1}{d}\right)}{\frac{d_1}{d} \prod_{p/\frac{d_1}{d}} \left(1 - \frac{1}{p}\right)} + O(\tau(\text{rad}(2n))) \end{aligned}$$

We apply again [Lemma 3](#) on $\sum_{\frac{d_1}{d}/\frac{N_z}{d}} \frac{\mu(\frac{d_1}{d})}{\frac{d_1}{d} \prod_{p/\frac{d_1}{d}} (1-\frac{1}{p})}$ we obtain .

$$\begin{aligned} \sum_{\frac{d_1}{d}/\frac{N_z}{d}} \frac{\mu(\frac{d_1}{d})}{\frac{d_1}{d} \prod_{p/\frac{d_1}{d}} (1-\frac{1}{p})} &= \prod_{p/\frac{N_z}{d}} \left(1-\frac{1}{p(1-\frac{1}{p})}\right) \\ &= \prod_{p/\frac{N_z}{d}} \left(1-\frac{1}{p-1}\right) \\ &= \prod_{p/\frac{N_z}{d}} \left(\frac{p-2}{p-1}\right) \end{aligned}$$

$$\begin{aligned} \text{Then } R'(2n) &= (n-z) \sum_{d \in F} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1-\frac{1}{p}\right) \prod_{p/\frac{N_z}{d}} \left(\frac{p-2}{p-1}\right) + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d \in F} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(\frac{p-1}{p}\right) \left(\frac{p-2}{p-1}\right) + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d \in F} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(\frac{p-2}{p}\right) + O(\tau(\text{rad}(2n))) \\ &= (n-z) \sum_{d \in F} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right) + O(\tau(\text{rad}(2n))) \end{aligned}$$

$$\begin{aligned} \text{We have } F &= \{d=d_1 \wedge d_2/, d_1/N_z, d_2/N_z, d/\text{rad}(2n)\} \\ &= \{d/\text{rad}(2n) / d \leq \lfloor \sqrt{x} \rfloor\} \end{aligned}$$

then $F \subset \{d/\text{rad}(2n)\}$.

$$\text{which means that, } \sum_{d \in F} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right) \geq \sum_{d/\text{rad}(2n)} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right)$$

$$\begin{aligned} \text{we have } \sum_{d/\text{rad}(2n)} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right) &= \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right) \sum_{d/\text{rad}(2n)} \frac{\prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right)}{d \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right)} \\ &= \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right) \sum_{d/\text{rad}(2n)} \frac{\prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right)}{d \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right)} \end{aligned}$$

if $\gcd(d,2) \neq 2$ then $\frac{N_z}{d}$ is even and $2/\frac{N_z}{d}$ which means that $\prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right) = 0$.

$$\begin{aligned} \text{Then } \sum_{d/\text{rad}(2n)} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right) &= \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right) \sum_{d/\text{rad}(2n), \gcd(d,2)=2} \frac{\prod_{p/\frac{N_z}{d}} \left(1-\frac{2}{p}\right)}{d \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right)} \\ &= \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right) \sum_{\frac{d}{2}/\frac{\text{rad}(2n)}{2}} \frac{\prod_{p/\frac{N_z}{\frac{d}{2}}} \left(1-\frac{2}{p}\right)}{d \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right)} \\ &= \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right) \sum_{\frac{d}{2}/\frac{\text{rad}(2n)}{2}} \frac{\prod_{p/\frac{N_z}{\frac{d}{2}}} \left(1-\frac{2}{p}\right)}{d \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right)} \\ &= \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right) \sum_{\frac{d}{2}/\frac{\text{rad}(2n)}{2}} \frac{\prod_{p/\frac{N_z}{\frac{d}{2}}} \left(1-\frac{2}{p}\right)}{d \prod_{p/\frac{N_z}{2}} \left(1-\frac{2}{p}\right) \prod_{p/\frac{d}{2}} \left(1-\frac{2}{p}\right)} \end{aligned}$$

$$\begin{aligned}
&= \prod_{p/\frac{N_z}{2}} \left(1 - \frac{2}{p}\right) \sum_{\frac{d}{2}/\frac{\text{rad}(2n)}{2}} \frac{1}{d \prod_{p/\frac{d}{2}} \left(1 - \frac{2}{p}\right)} \\
&= \frac{1}{2} \prod_{p/\frac{N_z}{2}} \left(1 - \frac{2}{p}\right) \sum_{\frac{d}{2}/\frac{\text{rad}(2n)}{2}} \frac{2}{d \prod_{p/\frac{d}{2}} \left(1 - \frac{2}{p}\right)} \\
&= \frac{1}{2} \prod_{p/\frac{N_z}{2}} \left(1 - \frac{2}{p}\right) \sum_{\frac{d}{2}/\frac{\text{rad}(2n)}{2}} \frac{1}{\frac{d}{2} \prod_{p/\frac{d}{2}} \left(1 - \frac{2}{p}\right)}
\end{aligned}$$

since $\frac{1}{\frac{d}{2} \prod_{p/\frac{d}{2}} \left(1 - \frac{2}{p}\right)}$ is multiplicative then by [Lemma 3](#) we have $\sum_{\frac{d}{2}/\frac{\text{rad}(2n)}{2}} \frac{1}{\frac{d}{2} \prod_{p/\frac{d}{2}} \left(1 - \frac{2}{p}\right)}$

$$\begin{aligned}
&\text{is also multiplicative and } \sum_{\frac{d}{2}/\frac{\text{rad}(2n)}{2}} \frac{1}{\frac{d}{2} \prod_{p/\frac{d}{2}} \left(1 - \frac{2}{p}\right)} = \prod_{p/\frac{\text{rad}(2n)}{2}} \left(1 + \frac{1}{p \left(1 - \frac{2}{p}\right)}\right) \\
&= \prod_{p/\frac{\text{rad}(2n)}{2}} \left(1 + \frac{1}{p-2}\right) \\
&= \prod_{p/\frac{\text{rad}(2n)}{2}} \frac{p-1}{p-2} \\
&= \prod_{p/2n, p \neq 2} \frac{p-1}{p-2}
\end{aligned}$$

$$\text{Then } \sum_{d/\text{rad}(2n)} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1 - \frac{2}{p}\right) = \frac{\prod_{p/2n, p \neq 2} \frac{p-1}{p-2}}{2} \prod_{p/\frac{N_z}{2}} \left(1 - \frac{2}{p}\right)$$

$$\text{We set } C_n = \prod_{p/2n, p \neq 2} \frac{p-1}{p-2}$$

$$\text{Then } \sum_{d/\text{rad}(2n)} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1 - \frac{2}{p}\right) = \frac{C_n}{2} \prod_{p/\frac{N_z}{2}} \left(1 - \frac{2}{p}\right)$$

$$\text{Then } \sum_{d \in F} \frac{1}{d} \prod_{p/\frac{N_z}{d}} \left(1 - \frac{2}{p}\right) \geq \frac{C_n}{2} \prod_{p/\frac{N_z}{2}} \left(1 - \frac{2}{p}\right)$$

$$\text{Which means that } R'(2n) \geq \frac{C_n}{2} (n-z) \prod_{p/\frac{N_z}{2}} \left(1 - \frac{2}{p}\right) - \tau(\text{rad}(2n)) \text{ such that } C_n = \prod_{p/2n, p \neq 2} \frac{p-1}{p-2}.$$

We know that , $R(2n) \geq R'(2n)$.

$$\text{Then } R(2n) \geq \frac{C_n}{2} (n - \lfloor \sqrt{2n} \rfloor) \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right) - B(n), \text{ such that } B(n) = \tau(\text{rad}(2n))$$

$$\text{and } C_n = \prod_{p/2n, p \neq 2} \frac{p-1}{p-2} .$$

By [Lemma 4](#) we have $\prod_{p \leq x, p \neq 2} \left(1 - \frac{2}{p}\right) \sim \frac{1}{\log(x)^2}$, for all sufficiently large x .

By [Lemma 7](#) we have for $\varepsilon = \frac{1}{2}$, $B(n) = \tau(\text{rad}(2n)) = o((\text{rad}(2n))^{\frac{1}{2}}) = o((2n)^{\frac{1}{2}})$

Then .

$$\begin{aligned} \frac{C_n}{2}(n - \lfloor \sqrt{2n} \rfloor) \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right) - B(n) &= \frac{C_n}{2}(n - \lfloor \sqrt{2n} \rfloor) \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right) - o\left((2n)^{\frac{1}{2}}\right) \\ &= \frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right) \left(1 - \frac{1}{\sqrt{2n}} - o\left(\frac{(2n)^{\frac{1}{2}}}{\frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right)}\right)\right) \end{aligned}$$

We have

$$\begin{aligned} \frac{(2n)^{\frac{1}{2}}}{\frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right)} &= \frac{\sqrt{2}}{\frac{C_n}{2} \sqrt{n} \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right)} \\ &\sim \frac{\sqrt{2} \log(\sqrt{2n})^2}{\frac{C_n}{2} \sqrt{n}} \\ &\sim \frac{\sqrt{2} \log(2n)^2}{2C_n \sqrt{n}} \end{aligned}$$

Since $\frac{\sqrt{2} \log(2n)^2}{2C_n \sqrt{n}} \rightarrow 0$ when $x \rightarrow \infty$, then $\frac{(2n)^{\frac{1}{2}}}{\frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right)} = o(1)$

Then $1 - \frac{1}{\sqrt{2n}} - o\left(\frac{(2n)^{\frac{1}{2}}}{\frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right)}\right) = 1 + o(1)$

Then $\frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right) \left(1 - \frac{1}{\sqrt{2n}} - o\left(\frac{(2n)^{\frac{1}{2}}}{\frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right)}\right)\right) = \frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right) (1 + o(1))$

Then $\frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right) \left(1 - \frac{1}{\sqrt{2n}} - o\left(\frac{(2n)^{\frac{1}{2}}}{\frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right)}\right)\right) \sim \frac{C_n}{2} n \prod_{p \leq \lfloor \sqrt{2n} \rfloor, p \neq 2} \left(1 - \frac{2}{p}\right)$

$$\begin{aligned} &\sim \frac{C_n}{2} \frac{4n}{\log(2n)^2} \\ &\sim 2C_n \frac{n}{\log(2n)^2} \end{aligned}$$

This means that $R(2n) \rightarrow \infty$ when $n \rightarrow \infty$

[This confirm The Goldbach conjecture is true](#)

Acknowledgments .

I would like to thank the referee for his/her detailed comments.

References .

- [01] Elementary Methods in Number Theory - Nathanson M.B
- chapter 6.4 ; multiplicative functions .
- [02] H. Davenport. Multiplicative Number Theory, volume 74 of
Graduate Texts in Mathematics. Springer-Verlag, New York, 2nd edition, 1980.
- [03] J.-P. Serre. A Course in Arithmetic, volume 7 of Graduate

Texts in Mathematics. Springer-Verlag, New York, 1973.

- [04] Les Nombres premiers, entre l'ordre et le chaos .Dunad -Gérald Tenenbaum
-Michel Mendès France (page 41)
- [05] Diophantine equations - en.wikipedia.org
- [06] Divisor function -en.wikipedia.org .
- [07] Yuan Wang, Goldbach Conjecture, 2nd edn., World Scientific Publishing,
New Jersey, Singapore (2000).
- [08] Richard K. Guy, Unsolved problems in number theory, Third Edition,
Springer-Verlag, Berlin (2004).
- [09] Nikolai G. Chudakov, On the Goldbach problem, Doklady Akademii Nauk SSSR, 17 (1937),
335-338. [10] T. Estermann, On Goldbach's problem: proof that