

A Result of Even & Prime

Shan Jian Wang

wangjianshan@oakmon.cn

Abstract

Conjecture:

Any even number greater than 2 can be written as the sum of two prime numbers.
Does the prime pair exist universally? If does, is the prime pair unique relatively? If not, how many prime pairs are there in an even?

Method:

Triangular lattice

Result:

The number of prime pairs in an even can be expressed analytically and graphically

Conclusion:

Any even number greater than 2 can be written as the sum of two prime numbers

Keywords

Goldbach; Euler; even; prime

Catalogue

Preface

1. Structures

1.1. Concepts

1.2. Discussions

2. Functions

2.1. Function: $N(a)$

2.2. Function: $P(a)$

3. Analysis

3.1. $U(a)-T(a) = S(a)-N(a)$

3.2. $H(a)=Z(a)/Y(a)$

3.3.1. $H(a)\sim J(a)/(J(a)+K(a)), a>0.$

3.3.2. $T(a)\sim (Y(a)-Y(a-2))*J(a)/(J(a)+K(a))+X(a)-X(a-2)-S(a)+N(a), a>0.$

Postscript

Conflict of interest statement

Data availability statement

References

Preface

Let $x > 0$, and $\pi(x)$ represents the number of prime numbers not exceeding x .

$$\pi(x) \sim x / \ln(x)$$

1. Structures

1.1. Concepts

Set of natural numbers is recorded as N , $N = \{n\}$. $n = 0, 1, 2, \dots$

One variable belongs to N , mark it as n ;

Two variables belong to N , mark them as n_1 and n_2 .

Set of positive integral numbers is recorded as P , $P = \{p\}$. $p = 1, 2, 3, \dots$

One variable belongs to P , mark it as p ;

Two variables belong to P , mark them as p_1 and p_2 .

Set of even numbers is recorded as A , $A = \{a | a = 2 * n\}$. $a = 0, 2, 4, \dots$

One variable belongs to A , mark it as a .

Two variables belong to A , mark them as a_1 and a_2 .

Set of odd numbers is recorded as B , $B = \{b | b = 2 * n + 1\}$. $b = 1, 3, 5, \dots$

One variable belongs to B , mark it as b .

Two variables belong to B , mark them as b_1 and b_2 .

Set of odd composite numbers is recorded as C , $C = \{c | c = (2 * p_1 + 1) * (2 * p_2 + 1)\}$. $c = 9, 15, 21, \dots$

One variable belongs to C , mark it as c .

Two variables belong to C , mark them as c_1 and c_2 .

Set of one is recorded as Q , $Q = \{1\}$.

Set of two is recorded as R , $R = \{2\}$.

Set of prime numbers is recorded as D ,

$D = \{d | d \text{ belongs to } B \text{ and } R, d \text{ does not belong to } C \text{ or } Q\}$. $d = 2, 3, 5, \dots$

One variable belongs to D , mark it as d .

Two variables belong to D , mark them as d_1 and d_2 .

1.2. Discussions

Let $a > 0$, and $T(a)$ represents the number of prime pairs in the even a ,

Let $a > 0$, and $N(a)$ represents the number of odd pairs in the even a ;

In the same a , $T(a)$ does not exceed $N(a)$.

2. Functions

2.1. Function: $N(a)$

$a = a/2 + a/2$, $a > 0$.

If $a/2$ belongs to A , $a = [(a/2 - 1) - 2n] + [(a/2 + 1) + 2n]$.

Shan Jian Wang

$(a/2+1)-2n$ is noted as bL , $(a/2+1)+2n$ is noted as bR .

$n < (a-2)/4$, $\text{Card}(n) = a/4$.

If $a/2$ belongs to B , $a = (a/2 - 2n) + (a/2 + 2n)$.

$a/2 - 2n$ is noted as bL , $a/2 + 2n$ is noted as bR .

$n < a/4$, $\text{Card}(n) = (a+2)/4$.

Three functions: bL , bR ; $N(a)$.

$bL = (a/2 + 1) - 2n$, $a/2$ belongs to A ; $bL = a/2 - 2n$, $a/2$ belongs to B .

$bR = (a/2 + 1) + 2n$, $a/2$ belongs to A ; $bR = a/2 + 2n$, $a/2$ belongs to B .

$\text{Card}(n)$ is noted as $N(a)$:

$N(a) = a/4$, $a/2$ belongs to A ; $N(a) = (a+2)/4$, $a/2$ belongs to B .

$N(a) \sim a/4$, $a > 0$.

Let $P(a) = bR - bL$:


The vertical axis represents positive even numbers, the horizontal axis represents $P(a)$. Any intersection corresponds to $(a, P(a))$ and (bL, bR) .

	0	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36	38	40	42
2	1.1																					
4		1.3																				
6	3.3		1.5																			
8		3.5		1.7																		
10	5.5		3.7		1.9																	
12		5.7		3.9		1.11																
14	7.7		5.9		3.11		1.13															
16		7.9		5.11		3.13		1.15														
18	9.9		7.11		5.13		3.15		1.17													
20		9.11		7.13		5.15		3.17		1.19												
22	11.11		9.13		7.15		5.17		3.19		1.21											
24		11.13		9.15		7.17		5.19		3.21		1.23										
26	13.13		11.15		9.17		7.19		5.21		3.23		1.25									
28		13.15		11.17		9.19		7.21		5.23		3.25		1.27								
30	15.15		13.17		11.19		9.21		7.23		5.25		3.27		1.29							
32		15.17		13.19		11.21		9.23		7.25		5.27		3.29		1.31						
34	17.17		15.19		13.21		11.23		9.25		7.27		5.29		3.31		1.33					
36		17.19		15.21		13.23		11.25		9.27		7.29		5.31		3.33		1.35				
38	19.19		17.21		15.23		13.25		11.27		9.29		7.31		5.33		3.35		1.37			
40		19.21		17.23		15.25		13.27		11.29		9.31		7.33		5.35		3.37		1.39		
42	21.21		19.23		17.25		15.27		13.29		11.31		9.33		7.35		5.37		3.41		1.41	
44		21.23		19.25		17.27		15.29		13.31		11.33		9.35		7.37		5.39		3.41		1.43

Delete pairs in cells and put $P(a)$ in, then there is the triangular lattice:

2	0																					
4		2																				
6	0		4																			
8		2		6																		
10	0		4		8																	
12		2		6		10																
14	0		4		8		12															
16		2		6		10		14														
18	0		4		8		12		16													
20		2		6		10		14		18												
22	0		4		8		12		16		20											
24		2		6		10		14		18		22										
26	0		4		8		12		16		20		24									
28		2		6		10		14		18		22		26								
30	0		4		8		12		16		20		24		28							
32		2		6		10		14		18		22		26		30						
34	0		4		8		12		16		20		24		28		32					
36		2		6		10		14		18		22		26		30		34				
38	0		4		8		12		16		20		24		28		32		36			
40		2		6		10		14		18		22		26		30		34		38		
42	0		4		8		12		16		20		24		28		32		36		40	
44		2		6		10		14		18		22		26		30		34		38		42

2.2. Function: $P(a)$




If f belongs to A , then $\{(a, P(a)) | a=f\}$ is noted as $\{L=f\}$.

If g belongs to B , then $G = \{(bL, bR) | bL=g \text{ or } bR=g\}$ is noted as $\{R=g\}$.


Any odd composite number belongs to $(0, a)$ corresponds to one cell in $\{L=a\}$;

$$P(a)=|(a-g)-g|, a>g.$$


$$P(a)=|(a-1)-1|, a>1.$$



$$P(a)=|(a-3)-3|, a>3.$$



$$P(a)=|(a-5)-5|, a>5.$$




...

3. Analysis

3.1. $U(a)-T(a)=S(a)-N(a)$

Number in blank named cell, white the cells.



3.3.1. $H(a) \sim J(a)/(J(a)+K(a)), a > 0.$

$J = \{(cL, cR) | cL \text{ belongs to } (0, a/4] \text{ and } cR \text{ belongs to } (a/2, 3*a/4]\};$

Card(cL, cR) is noted as J(a), $J(a) = S(a/4) * (S(3*a/4) - S(a/2)).$

$K = \{(cL, cR) | cL \text{ belongs to } (a/4, a/2] \text{ and } cR \text{ belongs to } (3*a/4, a]\};$


Card(cL, cR) is noted as K(a), $K(a) = (S(a/2) - S(a/4)) * (S(a) - S(3*a/4)).$

$J(a)/(J(a)+K(a)) \sim (((a/4)/2 - (a/4)/\ln(a/4)) * (((3*a/4)/2 - (3*a/4)/\ln(3*a/4)) - ((a/2)/2 - (a/2)/\ln(a/2)))) / (((a/4)/2 - (a/4)/\ln(a/4)) * (((3*a/4)/2 - (3*a/4)/\ln(3*a/4)) - ((a/2)/2 - (a/2)/\ln(a/2))) + (((a/2)/2 - (a/2)/\ln(a/2)) - ((a/4)/2 - (a/4)/\ln(a/4))) * ((a/2 - a/\ln(a)) - ((3*a/4)/2 - (3*a/4)/\ln(3*a/4)))).$

3.3.2. $T(a) \sim (Y(a) - Y(a-2)) * J(a)/(J(a)+K(a)) + X(a) - X(a-2) - S(a) + N(a), a > 0.$

$T(a) \sim (Y(a) - Y(a-2)) * J(a)/(J(a)+K(a)) + X(a) - X(a-2) - S(a) + N(a);$

$T(a) \sim (((a/2)/2 - (a/2)/\ln(a/2)) * (((a-1)/2 - (a-1)/\ln(a-1)) - ((a/2-1)/2 - (a/2-1)/\ln(a/2-1))) - ((a/2-1)/2 - (a/2-1)/\ln(a/2-1)) * (((a-3)/2 - (a-3)/\ln(a-3)) - ((a/2-2)/2 - (a/2-2)/\ln(a/2-2))) * (((a/4)/2 - (a/4)/\ln(a/4)) * (((3*a/4)/2 - (3*a/4)/\ln(3*a/4)) - ((a/2)/2 - (a/2)/\ln(a/2)))) / (((a/4)/2 - (a/4)/\ln(a/4)) * (((3*a/4)/2 - (3*a/4)/\ln(3*a/4)) - ((a/2)/2 - (a/2)/\ln(a/2))) + (((a/2)/2 - (a/2)/\ln(a/2)) - ((a/4)/2 - (a/4)/\ln(a/4))) * ((a/2 - a/\ln(a)) - ((3*a/4)/2 - (3*a/4)/\ln(3*a/4)))) + ((a/2-1)/2 - (a/2-1)/\ln(a/2-1)) * ((a/2-1)/2 - (a/2-1)/\ln(a/2-1) + 1)/2 - ((a/2-2)/2 - (a/2-2)/\ln(a/2-2)) * ((a/2-2)/2 - (a/2-2)/\ln(a/2-2) + 1)/2 + a/\ln(a) - a/4.$



Question: How many prime pairs are there in an even?

Answer: $T(a)$, what has been explained above is the analytical approximation to $T(a)$ and its image.

Postscript

Prime number theorem shows that the number of prime numbers has an approaching, and the above explanation shows that if the number of prime numbers has an approaching then the number of prime pairs in an even has an approaching. Sequence of $\pi(x)$ is not strictly monotonic, at the same time $T(a)$ is not strictly monotonically increasing in the interval $\{a > 17\}$.

Conclusion: Any even number greater than 2 can be written as the sum of two prime numbers.

Conflict of interest statement:

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Data availability statement:

My manuscript has no associated data.

References:

- [1] Gronwall H T .Theory of Prime Numbers[J].Bulletin of the American Mathematical Society,1914,20(7):368-376.
- [2] Dickson E L .Review: D. N. Lehmer, List of Prime Numbers from 1 to 10,006,721[J].Bulletin of the American Mathematical Society,1915,21(7):355-356.
- [3] Atle S .An Elementary Proof of the Prime-Number Theorem for Arithmetic Progressions[J].Canadian Journal of Mathematics,1950,266-78.
- [4] Schoenfeld L .Review: T. Estermann, Introduction to modern prime number theory[J].Bulletin of the American Mathematical Society,1953,59(3):255-258.
- [5] Bays C ,Hudson H R .Zeroes of Dirichlet L-Functions and Irregularities in the Distribution of Primes[J].Mathematics of Computation,2000,69(230):861-866.
- [6] R P C .A pseudo zeta function and the distribution of primes.[J].Proceedings of the National Academy of Sciences of the United States of America,2000,97(14):7697-9.
- [7] Green B ,Tao T .The Primes Contain Arbitrarily Long Arithmetic Progressions[J].Annals of Mathematics,2008,167(2):481-547.
- [8] Tao T ,Ziegler T .The primes contain arbitrarily long polynomial progressions[J].Acta Mathematica,2008,201(2):213-305.
- [9] Kumchev V A ,Liu Y J .Sums of primes and squares of primes in short intervals[J].Monatshefte für Mathematik,2009,157(4):335-363.
- [10] Friedlander B J ,Iwaniec H .Hyperbolic prime number theorem[J].Acta Mathematica,2009,202(1):1-19.
- [11] Li X .A spectral interpretation for the explicit formula in the theory of prime numbers[J].Science China Mathematics,2010,53(3):791-812.
- [12] Tao ,T.,Ziegler , et al.Erratum to The primes contain arbitrarily long polynomial progressions (Acta Math., (2008), 201, (213-305), 10.1007/s11511-008-0032-5)[J].Acta Mathematica,2013,210(2):403-404.
- [13] Bounded gaps between primes[J].Annals of Mathematics,2014,179(3):1121-1174.
- [14] Pollack P .Bounded gaps between primes with a given primitive root[J].Algebra Number Theory,2014,8(7):1769-1786.
- [15] Jiamin L ,Jianya L .Triples of almost primes[J].Science China Mathematics,2023,66(12):2779-2794.