

Proof of Fermat's Last Theorem for odd primes

Minho Baek

ABSTRACT. It was already proved right that $x^n + y^n = z^n$, ($n > 2$) has no solutions in positive integers which we called Fermat's Last Theorem (FLT) by Andrew Wiles. But his proof would be impossible in the 17th century. Since Fermat showed he proved n =even numbers by leaving proof for $n=4$, many people have tried to prove the odd prime. I took the idea from Euler proof and proved in case of n =odd primes by simple method.

1. Introduction

Pierre de Fermat claimed he had proof that no three positive integer x , y and z satisfy the equation $x^n + y^n = z^n$ for n greater than 2 which we called Fermat's Last Theorem (FLT). For about 3 century, many people have tried to prove FLT. Finally, FLT was proved right by Andrew Wiles in 1995. However, the proof of Wiles is a modern math that is difficult to understand and complex. So, some people, who think Fermat prove FLT by himself, still believe elementary method exist. We know that FLT can be proved by proving n =odd primes because case of n =even proved by Fermat himself. I got an idea to solve for n =odd primes from Euler's proof. In this paper, I proved the case of n =odd primes number by simple method.

2. Proof for n =odd primes

$$x^n + y^n = z^n, (x < y < z)$$

Where x, y and z = positive integer, relatively prime.

This equation can be classified into three categories as follows.

Case 1. $(x, y, z) = (\text{even}, \text{odd}, \text{odd})$

Case 2. $(x, y, z) = (\text{odd}, \text{even}, \text{odd})$

Case 3. $(x, y, z) = (\text{odd}, \text{odd}, \text{even})$

Case 1. $(x, y, z) = (\text{even}, \text{odd}, \text{odd})$

Let $y=(u-v)$, $z=(u+v)$.

Assume u and v are not relatively prime.

Let $u=fU$, $v=fV$.

$$y = f(U - V)$$

$$z = f(U + V)$$

But this contradicts because y and z are relatively prime.

So, u and v are relatively prime.

Also u and v are opposite parity because y and z are odd.

$$x^n = (u + v)^n - (u - v)^n$$

$$x^n = (u^n + C_1 u^{n-1} v^1 + C_2 u^{n-2} v^2 + \dots + C_{n-2} u^2 v^{n-2} + C_{n-1} u^1 v^{n-1} + v^n) - (u^n - C_1 u^{n-1} v^1 + C_2 u^{n-2} v^2 + \dots - C_{n-2} u^2 v^{n-2} + C_{n-1} u^1 v^{n-1} - v^n)$$

Where $C = \{C_1, C_2, \dots, C_{n-2}, C_{n-1}\} = \left\{ \frac{n}{1}, \frac{n(n-1)}{1 \cdot 2}, \dots, \frac{n(n-1) \dots 4 \cdot 3}{1 \cdot 2 \dots (n-3)(n-2)}, \frac{n(n-1) \dots 2 \cdot 1}{1 \cdot 2 \dots (n-2)(n-1)} \right\}$

$$x^n = 2v(v^{n-1} + n(u^{n-1} + C'_3 u^{n-3} v^2 + \dots + C'_{n-2} u^2 v^{n-3}))$$

Where $C' = \{C'_1, C'_2, \dots, C'_{n-2}, C'_{n-1}\} = \left\{ \frac{1}{1}, \frac{(n-1)}{1 \cdot 2}, \dots, \frac{(n-1) \dots 4 \cdot 3}{1 \cdot 2 \dots (n-3)(n-2)}, \frac{(n-1) \dots 2 \cdot 1}{1 \cdot 2 \dots (n-2)(n-1)} \right\}$

$$x^n = 2v(v^{n-1} + n(u^{n-1} + m))$$

Where $m = C'_3 u^{n-3} v^2 + C'_5 u^{n-5} v^4 + \dots + C'_{n-4} u^4 v^{n-5} + C'_{n-2} u^2 v^{n-3}$

Assume u=even, v=odd.

$$v^{n-1} + n(u^{n-1} + m) \text{ is odd.}$$

Assume u=odd, v=even.

$$v^{n-1} + n(u^{n-1} + m) \text{ is odd.}$$

So, $v^{n-1} + n(u^{n-1} + m)$ is always odd.

Thus the greatest common factor of $2v$ and $v^{n-1} + n(u^{n-1} + m)$ is odd.

Assume the common factor is odd except 1 and n.

Let $v = fV$ and $v^{n-1} + n(u^{n-1} + m) = fN$.

$$f^{n-1} V^{n-1} + n(u^{n-1} + fM) = fN$$

Where $fM = f(C'_3 u^{n-3} f^1 V^2 + C'_5 u^{n-5} f^3 V^4 + \dots + C'_{n-4} u^4 f^{n-6} V^{n-5} + C'_{n-2} u^2 f^{n-4} V^{n-3})$

$$nu^{n-1} = f(N - f^{n-2} V^{n-1} - nM)$$

u and v have common factor of f.

But this contradicts because u and v are relatively prime.

So, the greatest common factor of $2v$ and $v^{n-1} + n(u^{n-1} + m)$ is either 1 or n.

Assume the greatest common factor of $2v$ and $v^{n-1} + n(u^{n-1} + m)$ is 1.

It is possible that $2v = p^n$ and $v^{n-1} + n(u^{n-1} + m) = q^n$.

Where p and q are relatively prime

$$x^n = p^n q^n$$

Assume the greatest common factor of $2v$ and $v^{n-1} + n(u^{n-1} + m)$ is n .

Let $v=nr$.

u and r are relatively prime because u and v are relatively prime.

$$x^n = 2nr(n^{n-1}r^{n-1} + n(u^{n-1} + m))$$

$$x^n = 2n^2r(n^{n-2}r^{n-1} + (u^{n-1} + m))$$

The greatest common factor of $2n^2r$ and $n^{n-2}r^{n-1} + (u^{n-1} + m)$ must be 1.

It is possible that $2n^2r = p'^n$ and $n^{n-2}r^{n-1} + (u^{n-1} + m) = q'^n$.

$$x^n = p'^n q'^n$$

Let $z=y+i$.

From $y=(u-v)$, $z=(u+v)$,

$$z - y = (u + v) - (u - v) = 2v$$

$$2v = i$$

Let $y=k+i$.

k is odd because y is odd and i is even.

Assume y and i are not relatively prime.

Let $y=fY$, $i=fI$

$$z = y + i$$

$$z = f(Y + I)$$

But this contradicts because y and z are relatively prime.

So, y and i are relatively prime.

Assume k and y are not relatively prime.

Let $k=fK$, $y=fY$.

$$i = y - k$$

$$i = f(Y - K)$$

But this contradicts because y and i are relatively prime.

So, k and y are relatively prime.

Let $k = (u' - v)$, $y = (u' + v)$.

Where $u' < u$

Assume u' and v are not relatively prime.

Let $u' = fU', v = fV$

$$k = f(U' - V)$$

$$y = f(U' + V)$$

But this contradicts because k and y are relatively prime.

So, u' and v are relatively prime.

Also u' and v are opposite parity because k and y are odd.

$$y^n - k^n = (u' + v)^n - (u' - v)^n$$

$$y^n - k^n = 2v(v^{n-1} + n(u'^{n-1} + m'))$$

Where $m' = C'_3 u'^{n-3} v^2 + C'_5 u'^{n-5} v^4 + \dots + C'_{n-4} u'^4 v^{n-5} + C'_{n-2} u'^2 v^{n-3}$

Assume $u' = \text{even}, v = \text{odd}$.

$v^{n-1} + n(u'^{n-1} + m')$ is odd.

Assume $u' = \text{odd}, v = \text{even}$.

$v^{n-1} + n(u'^{n-1} + m')$ is odd.

So, $v^{n-1} + n(u'^{n-1} + m')$ is always odd.

Thus the greatest common factor of $2v$ and $v^{n-1} + n(u'^{n-1} + m')$ is odd.

Assume the common factor is odd number except 1 and n .

Let $v = fV$ and $v^{n-1} + n(u'^{n-1} + m') = fN$.

$$f^{n-1} V^{n-1} + n(u'^{n-1} + fM') = fN$$

Where $fM' = f(C'_3 u'^{n-3} f^1 V^2 + C'_5 u'^{n-5} f^3 V^4 + \dots + C'_{n-4} u'^4 f^{n-6} V^{n-5} + C'_{n-2} u'^2 f^{n-4} V^{n-3})$

$$nu'^{n-1} = f(N - f^{n-2} V^{n-1} - nM')$$

u' and v have common factor of f .

But this contradicts because u' and v are relatively prime.

So, the greatest common factor of $2v$ and $v^{n-1} + n(u'^{n-1} + m')$ is either 1 or n .

Assume the greatest common factor of $2v$ and $v^{n-1} + n(u'^{n-1} + m')$ is 1.

It is possible that $2v = p^n$ and $v^{n-1} + n(u'^{n-1} + m') = \gamma$.

$$y^n - k^n = \gamma p^n$$

Assume $y^n = \gamma y'^n$, $k^n = \gamma k'^n$ or $y^n = \alpha p^n$, $k^n = \beta p^n$.

But this contradicts because k and y are relatively prime.

Assume $\gamma = s^n$.

$$y^n - k^n = s^n p^n$$

Where s and p are relatively prime

Let $y^n = c^n$, $k^n = b^n$, $s^n p^n = a^n$

$$a^n + b^n = c^n$$

But this equation contradicts by the method of infinite descent.

Assume $\gamma \neq s^n$.

$$y^n - k^n = \gamma p^n$$

Where $\gamma \neq s^n$

Let $k = t + i$.

t is odd because k is odd and i is even.

Assume k and i are not relatively prime.

Let $k = fK$, $i = fI$

$$y = k + i$$

$$y = f(K + I)$$

But this contradicts because k and y are relatively prime.

So, k and i are relatively prime.

Assume t and k are not relatively prime.

Let $t = fT$, $k = fK$.

$$i = k - t$$

$$i = f(K - T)$$

But this contradicts because k and i are relatively prime.

So, t and k are relatively prime.

Let $t = (u'' - v)$, $k = (u'' + v)$.

Assume u'' and v are not relatively prime.

Let $u'' = fU'', v = fV$.

$$t = f(U'' - V)$$

$$k = f(U'' + V)$$

But this contradicts because t and k are relatively prime.

So, u'' and v are relatively prime.

Also u'' and v are opposite parity because t and k are odd.

$$k^n - t^n = (u'' + v)^n - (u'' - v)^n$$

$$k^n - t^n = 2v(v^{n-1} + n(u''^{n-1} + m''))$$

Where $m'' = C'_3 u''^{n-3} v^2 + C'_5 u''^{n-5} v^4 + \dots + C'_{n-4} u''^4 v^{n-5} + C'_{n-2} u''^2 v^{n-3}$

Assume $u'' = \text{even}, v = \text{odd}$.

$v^{n-1} + n(u''^{n-1} + m'')$ is odd.

Assume $u'' = \text{odd}, v = \text{even}$.

$v^{n-1} + n(u''^{n-1} + m'')$ is odd.

So, $v^{n-1} + n(u''^{n-1} + m'')$ is always odd.

Thus the greatest common factor of $2v$ and $v^{n-1} + n(u''^{n-1} + m'')$ is odd.

Assume the common factor is odd number except 1 and n .

Let $v = fV$ and $v^{n-1} + n(u''^{n-1} + m'') = fN$.

$$f^{n-1} V^{n-1} + n(u''^{n-1} + fM'') = fN$$

Where $fM'' = f(C'_3 u''^{n-3} f^1 V^2 + C'_5 u''^{n-5} f^3 V^4 + \dots + C'_{n-4} u''^4 f^{n-6} V^{n-5} + C'_{n-2} u''^2 f^{n-4} V^{n-3})$

$$nu''^{n-1} = f(N - f^{n-2} V^{n-1} - nM'')$$

u'' and v have common factor of f .

But this contradicts because u'' and v are relatively prime.

So, the greatest common factor of $2v$ and $v^{n-1} + n(u''^{n-1} + m'')$ is either 1 or n .

Assume the greatest common factor of $2v$ and $v^{n-1} + n(u''^{n-1} + m'')$ is 1.

It is possible that $2v = p^n$ and $v^{n-1} + n(u''^{n-1} + m'') = \gamma'$.

$$k^n - t^n = \gamma' p^n$$

Assume $k^n = \gamma' y'^n$, $t^n = \gamma' k'^n$ or $k^n = \alpha' p^n$, $t^n = \beta' p^n$.

But this contradicts because t and k are relatively prime.

Assume $\gamma' = s'^n$.

$$k^n - t^n = s'^n p^n$$

Where s' and p are relatively prime

Let $k^n = c'^n, t^n = b'^n, s'^n p^n = a'^n$

$$a'^n + b'^n = c'^n$$

But this equation contradicts by the method of infinite descent.

Assume $\gamma' \neq s'^n$.

$$k^n - t^n = \gamma' p^n$$

Where $\gamma \neq s^n$

But this equation also contradicts by the method of infinite descent.

Assume the greatest common factor of $2v$ and $v^{n-1} + n(u''^{n-1} + m'')$ is n.

Let $v = nr$.

u'' and r are relatively prime because u'' and v are relatively prime.

$$k^n - t^n = 2nr(v^{n-1} + n(u''^{n-1} + m''))$$

$$k^n - t^n = 2n^2r(n^{n-2}r^{n-1} + (u''^{n-1} + m''))$$

The greatest common factor of $2n^2r$ and $n^{n-2}r^{n-1} + (u''^{n-1} + m'')$ must be 1.

It is possible that $2n^2r = p'^n$ and $n^{n-2}r^{n-1} + (u''^{n-1} + m'') = \delta'$.

$$k^n - t^n = \delta' p'^n$$

Assume $y^n = \delta' y''^n, k^n = \delta' k''^n$ or $y^n = \alpha''' p'^n, k^n = \beta''' p'^n$

But this contradicts because k and t are relatively prime.

Assume $\delta' = S'^n$.

$$k^n - t^n = S'^n p'^n$$

Where S and p are relatively prime

Let $y^n = c''''^n, k^n = b''''^n, S'^n p'^n = a''''^n$

$$a''''^n + b''''^n = c''''^n$$

But this equation contradicts by the method of infinite descent.

Assume $\delta' \neq S^n$.

$$k^n - t^n = \delta' p'^n$$

Where $\delta' \neq S^n$

But this equation also contradicts by the method of infinite descent.

Assume the greatest common factor of $2v$ and $v^{n-1} + n(u'^{n-1} + m')$ is n .

Let $v=nr$.

u' and r are relatively prime because u' and v are relatively prime.

$$y^n - k^n = 2nr(v^{n-1} + n(u'^{n-1} + m'))$$

$$y^n - k^n = 2n^2r(n^{n-2}r^{n-1} + (u'^{n-1} + m'))$$

The greatest common factor of $2n^2r$ and $n^{n-2}r^{n-1} + (u'^{n-1} + m')$ must be 1.

It is possible that $2n^2r = p'^n$ and $n^{n-2}r^{n-1} + (u'^{n-1} + m') = \delta$.

$$y^n - k^n = \delta p'^n$$

Assume $y^n = \delta y'^n$, $k^n = \delta k'^n$ or $y^n = \alpha'' p'^n$, $k^n = \beta'' p'^n$

But this contradicts because k and y are relatively prime.

Assume $\delta = S^n$.

$$y^n - k^n = S^n p'^n$$

Where S and p' are relatively prime

Let $y^n = c''^n$, $k^n = b''^n$, $S^n p'^n = a''^n$

$$a''^n + b''^n = c''^n$$

But this equation contradicts by the method of infinite descent.

Assume $\delta \neq S^n$.

$$y^n - k^n = \delta p'^n$$

Where $\delta \neq S^n$

Let $k=t+i$.

t is odd because k is odd and i is even.

Assume k and i are not relatively prime.

Let $k=fK, i=fl$

$$y = k + i$$

$$y = f(K + I)$$

But this contradicts because k and y are relatively prime.

So, k and i are relatively prime.

Assume t and k are not relatively prime.

Let $t=fT, k=fK$.

$$i = k - t$$

$$i = f(K - T)$$

But this contradicts because k and i are relatively prime.

So, t and k are relatively prime.

Let $t = (u'' - v), k = (u'' + v)$.

Assume u'' and v are not relatively prime.

Let $u'' = fU'', v = fV$.

$$t = f(U'' - V)$$

$$k = f(U'' + V)$$

But this contradicts because t and k are relatively prime.

So, u'' and v are relatively prime.

Also u'' and v are opposite parity because t and k are odd.

$$k^n - t^n = (u'' + v)^n - (u'' - v)^n$$

$$k^n - t^n = 2v(v^{n-1} + n(u''^{n-1} + m''))$$

Where $m'' = C'_3 u''^{n-3} v^2 + C'_5 u''^{n-5} v^4 + \dots + C'_{n-4} u''^4 v^{n-5} + C'_{n-2} u''^2 v^{n-3}$

Assume $u''=even, v=odd$.

$v^{n-1} + n(u''^{n-1} + m'')$ is odd.

Assume $u''=odd, v=even$.

$v^{n-1} + n(u''^{n-1} + m'')$ is odd.

So, $v^{n-1} + n(u''^{n-1} + m'')$ is always odd.

Thus the greatest common factor of $2v$ and $v^{n-1} + n(u''^{n-1} + m'')$ is odd.

Assume the common factor is odd number except 1 and n .

Let $v = fV$ and $v^{n-1} + n(u''^{n-1} + m'') = fN$.

$$f^{n-1}V^{n-1} + n(u''^{n-1} + fM'') = fN$$

Where $fM'' = f(C'_3 u''^{n-3} f^1 V^2 + C'_5 u''^{n-5} f^3 V^4 + \dots + C'_{n-4} u''^4 f^{n-6} V^{n-5} + C'_{n-2} u''^2 f^{n-4} V^{n-3})$

$$nu''^{n-1} = f(N - f^{n-2}V^{n-1} - nM'')$$

u'' and v have common factor of f .

But this contradicts because u'' and v are relatively prime.

So, the greatest common factor of $2v$ and $v^{n-1} + n(u''^{n-1} + m'')$ is either 1 or n .

Assume the greatest common factor of $2v$ and $v^{n-1} + n(u''^{n-1} + m'')$ is 1.

It is possible that $2v = p^n$ and $v^{n-1} + n(u''^{n-1} + m'') = \gamma'$.

$$k^n - t^n = \gamma' p^n$$

Assume $k^n = \gamma' y'^n$, $t^n = \gamma' k'^n$ or $k^n = \alpha' p^n$, $t^n = \beta' p^n$.

But this contradicts because t and k are relatively prime.

Assume $\gamma' = s'^n$.

$$k^n - t^n = s'^n p^n$$

Where s' and p are relatively prime

Let $k^n = c'^n$, $t^n = b'^n$, $s'^n p^n = a'^n$

$$a'^n + b'^n = c'^n$$

But this equation contradicts by the method of infinite descent.

Assume $\gamma' \neq s'^n$.

$$k^n - t^n = \gamma' p^n$$

Where $\gamma' \neq s'^n$

But this equation also contradicts by the method of infinite descent.

Assume the greatest common factor of $2v$ and $v^{n-1} + n(u''^{n-1} + m'')$ is n .

Let $v = nr$.

u'' and r are relatively prime because u'' and v are relatively prime.

$$k^n - t^n = 2nr(v^{n-1} + n(u''^{n-1} + m''))$$

$$k^n - t^n = 2n^2r(n^{n-2}r^{n-1} + (u''^{n-1} + m''))$$

The greatest common factor of $2n^2r$ and $n^{n-2}r^{n-1} + (u''^{n-1} + m'')$ must be 1.

It is possible that $2n^2r = p'^n$ and $n^{n-2}r^{n-1} + (u''^{n-1} + m'') = \delta'$.

$$k^n - t^n = \delta'p'^n$$

Assume $y^n = \delta'y''^n$, $k^n = \delta'k''^n$ or $y^n = \alpha'''p'^n$, $k^n = \beta'''p'^n$

But this contradicts because k and t are relatively prime.

Assume $\delta' = S'^n$.

$$k^n - t^n = S'^n p'^n$$

Where S and p are relatively prime

Let $y^n = c''''^n$, $k^n = b''''^n$, $S'^n p'^n = a''''^n$

$$a''''^n + b''''^n = c''''^n$$

But this equation contradicts by the method of infinite descent.

Assume $\delta' \neq S'^n$.

$$k^n - t^n = \delta'p'^n$$

Where $\delta' \neq S'^n$

But this equation also contradicts by the method of infinite descent.

Case 2. (x, y, z) = (odd, even, odd)

Case 2 has no solution because it can be proved in the same way as Case 1.

Case 3. (x, y, z) = (odd, odd, even)

Let $x=(u-v)$, $y=(u+v)$.

Assume u and v are not relatively prime.

Let $u=fU$, $v=fV$.

$$x = f(U - V)$$

$$y = f(U + V)$$

But this contradicts because x and y are relatively prime.

So, u and v are relatively prime.

Also u and v are opposite parity because x and y are odd.

$$z^n = (u - v)^n + (u + v)^n$$

$$z^n = (u^n - C_1 u^{n-1} v^1 + C_2 u^{n-2} v^2 + \dots - C_{n-2} u^2 v^{n-2} + C_{n-1} u^1 v^{n-1} - v^n) + (u^n + C_1 u^{n-1} v^1 + C_2 u^{n-2} v^2 + \dots + C_{n-2} u^2 v^{n-2} + C_{n-1} u^1 v^{n-1} + v^n)$$

$$\text{Where } C = \{C_1, C_2, \dots, C_{n-2}, C_{n-1}\} = \left\{ \frac{n}{1}, \frac{n(n-1)}{1 \cdot 2}, \dots, \frac{n(n-1) \dots 4 \cdot 3}{1 \cdot 2 \dots (n-3)(n-2)}, \frac{n(n-1) \dots 2 \cdot 1}{1 \cdot 2 \dots (n-2)(n-1)} \right\}$$

$$z^n = 2u(u^{n-1} + n(v^{n-1} + C'_{n-3} v^{n-3} u^2 + \dots + C'_2 v^2 u^{n-3}))$$

$$\text{Where } C' = \{C'_1, C'_2, \dots, C'_{n-2}, C'_{n-1}\} = \left\{ \frac{1}{1}, \frac{(n-1)}{1 \cdot 2}, \dots, \frac{(n-1) \dots 4 \cdot 3}{1 \cdot 2 \dots (n-3)(n-2)}, \frac{(n-1) \dots 2 \cdot 1}{1 \cdot 2 \dots (n-2)(n-1)} \right\}$$

$$z^n = 2u(u^{n-1} + n(v^{n-1} + m))$$

$$\text{Where } m = C'_{n-3} v^{n-3} u^2 + C'_{n-5} v^{n-5} u^4 + \dots + C'_4 v^4 u^{n-5} + C'_2 v^2 u^{n-3}$$

Assume u=even, v=odd.

$$u^{n-1} + n(v^{n-1} + m) \text{ is odd.}$$

Assume u=odd, v=even.

$$u^{n-1} + n(v^{n-1} + m) \text{ is odd.}$$

So, $u^{n-1} + n(v^{n-1} + m)$ is always odd.

Thus the greatest common factor of $2u$ and $u^{n-1} + n(v^{n-1} + m)$ is odd.

Assume the common factor is odd except 1 and n.

$$\text{Let } u = fU \text{ and } u^{n-1} + n(v^{n-1} + m) = fN.$$

$$f^{n-1} U^{n-1} + n(u^{n-1} + fM) = fN$$

$$\text{Where } fM = f(C'_{n-3} v^{n-3} f^1 U^2 + C'_{n-5} v^{n-5} f^3 U^4 + \dots + C'_4 v^4 f^{n-6} U^{n-5} + C'_2 v^2 f^{n-4} U^{n-3})$$

$$nu^{n-1} = f(N - f^{n-2} U^{n-1} - nM)$$

u and v have common factor of f.

But this contradicts because u and v are relatively prime.

So, the greatest common factor of $2u$ and $u^{n-1} + n(v^{n-1} + m)$ is either 1 or n.

Assume the greatest common factor of $2u$ and $u^{n-1} + n(v^{n-1} + m)$ is 1.

$$\text{It is possible that } 2u = p^n \text{ and } u^{n-1} + n(v^{n-1} + m) = q^n.$$

Where p and q are relatively prime

$$z^n = p^n q^n$$

Assume the greatest common factor of $2u$ and $u^{n-1} + n(v^{n-1} + m)$ is n.

Let $u=nr$.

u and r are relatively prime because u and v are relatively prime.

$$z^n = 2nr(u^{n-1} + n(v^{n-1} + m))$$

$$z^n = 2n^2r(n^{n-2}r^{n-1} + (v^{n-1} + m))$$

The greatest common factor of $2n^2r$ and $n^{n-2}r^{n-1} + (v^{n-1} + m)$ must be 1.

It is possible that $2n^2r = p'^n$ and $n^{n-2}r^{n-1} + (v^{n-1} + m) = q'^n$.

$$z^n = p'^n q'^n$$

Let $k = u - v'$, $t = u + v'$

Where $v'=nv$

Assume u and v' are not relatively prime.

Let $u=fU$, $v'=f$ $V'=fnV$.

$$x = f(U - V'/n)$$

$$y = f(U + V'/n)$$

But this contradicts because x and y are relatively prime.

So, u and v' are relatively prime.

Assume k and t are not relatively prime.

Let $u=fU$, $v=fV$.

$$k = f(U - V')$$

$$t = f(U + V')$$

But this contradicts because u and v' are relatively prime.

So, k and t are relatively prime.

Also u and v' are opposite parity because x and y are odd.

$$k^n + t^n = (u - v')^n + (u + v')^n$$

$$k^n + t^n = 2u(u^{n-1} + n(v'^{n-1} + m'))$$

Where $m' = C'_{n-3}v'^{n-3}u^2 + C'_{n-5}v'^{n-5}u^4 + \dots + C'_4v'^4u^{n-5} + C'_2v'^2u^{n-3}$

Assume u =even, v' =odd.

$u^{n-1} + n(v'^{n-1} + m')$ is odd.

Assume $u=\text{odd}$, $v'=\text{even}$.

$u^{n-1} + n(v'^{n-1} + m')$ is odd.

So, $u^{n-1} + n(v'^{n-1} + m')$ is always odd.

Thus the greatest common factor of $2u$ and $u^{n-1} + n(v'^{n-1} + m')$ is odd.

Assume the common factor is odd number except 1 and n .

Let $u = fU$, $u^{n-1} + n(v'^{n-1} + m') = fN$.

$$f^{n-1}U^{n-1} + n(v'^{n-1} + fM') = fN$$

Where $fM' = f(C'_{n-3}v'^{n-3}f^1U^2 + C'_{n-5}v'^{n-5}f^3U^4 + \dots + C'_4v'^4f^{n-6}U^{n-5} + C'_2v'^2f^{n-4}U^{n-3})$

$$nv'^{n-1} = f(N - f^{n-2}U^{n-1} - nM')$$

u and v' have common factor of f .

But this contradicts because u and v' are relatively prime.

So, the greatest common factor of $2u$ and $u^{n-1} + n(v'^{n-1} + m')$ is either 1 or n .

Assume the greatest common factor of $2u$ and $u^{n-1} + n(v'^{n-1} + m')$ is 1.

It is possible that $2u = p^n$ and $u^{n-1} + n(v'^{n-1} + m') = \gamma$.

$$k^n + t^n = \gamma p^n$$

Assume $k^n = \gamma k'^n$, $t^n = \gamma t'^n$ or $k^n = \alpha p^n$, $t^n = \beta p^n$.

But this contradicts because k and t are relatively prime.

Assume $\gamma = s^n$.

$$x^n + k^n = s^n p^n$$

Where s and p are relatively prime

Let $x^n = a^n$, $k^n = b^n$, $s^n p^n = c^n$

$$a^n + b^n = c^n$$

But this equation contradicts by the method of infinite descent.

Assume $\gamma \neq s^n$.

$$k^n + t^n = \gamma p^n$$

Where $\gamma \neq s^n$

Let $w = u - v''$, $o = u + v''$

Where $v'' = n v'$

Assume u and v'' are not relatively prime.

Let $u = fU$, $v'' = fV'' = fn V'$.

$$k = f(U - V''/n)$$

$$t = f(U + V''/n)$$

But this contradicts because k and t are relatively prime.

So, u and v'' are relatively prime.

Assume w and o are not relatively prime.

Let $u = fU$, $v = fV$.

$$w = f(U - V')$$

$$o = f(U + V')$$

But this contradicts because u and v'' are relatively prime.

So, w and o are relatively prime.

Also u and v'' are opposite parity because k and t are odd.

$$w^n + o^n = (u - v'')^n + (u + v'')^n$$

$$w^n + o^n = 2u(u^{n-1} + n(v''^{n-1} + m''))$$

Where $m'' = C'_{n-3}v''^{n-3}u^2 + C'_{n-5}v''^{n-5}u^4 + \dots + C'_4v''^4u^{n-5} + C'_2v''^2u^{n-3}$

Assume $u = \text{even}$, $v'' = \text{odd}$.

$u^{n-1} + n(v''^{n-1} + m'')$ is odd.

Assume $u = \text{odd}$, $v'' = \text{even}$.

$u^{n-1} + n(v''^{n-1} + m'')$ is odd.

So, $u^{n-1} + n(v''^{n-1} + m'')$ is always odd.

Thus the greatest common factor of $2u$ and $u^{n-1} + n(v''^{n-1} + m'')$ is odd.

Assume the common factor is odd number except 1 and n .

Let $u = fU$ and $u^{n-1} + n(v''^{n-1} + m'') = fN$.

$$f^{n-1}U^{n-1} + n(v''^{n-1} + fM'') = fN$$

Where $fM'' = f(C'_{n-3}v''^{n-3}f^1U^2 + C'_{n-5}v''^{n-5}f^3U^4 + \dots + C'_4v''^4f^{n-6}U^{n-5} + C'_2v''^2f^{n-4}U^{n-3})$

$$nv''^{n-1} = f(N - f^{n-2}U^{n-1} - nM'')$$

u and v'' have common factor of f.

But this contradicts because u and v'' are relatively prime.

So, the greatest common factor of $2u$ and $u^{n-1} + n(v''^{n-1} + m'')$ is either 1 or n.

Assume the greatest common factor of $2u$ and $u^{n-1} + n(v''^{n-1} + m'')$ is 1.

It is possible that $2u = p^n$ and $u^{n-1} + n(v''^{n-1} + m'') = \gamma'$.

$$w^n + o^n = \gamma'p^n$$

Assume $w^n = \gamma'w'^n$, $o^n = \gamma'o'^n$ or $w^n = \alpha'p^n$, $o^n = \beta'p^n$.

But this contradicts because k and t are relatively prime.

Assume $\gamma' = s'^n$.

$$w^n + o^n = s'^np^n$$

Where s' and p are relatively prime

Let $w^n = a'^n$, $o^n = b'^n$, $s'^np^n = c'^n$

$$a'^n + b'^n = c'^n$$

But this equation contradicts by the method of infinite descent.

Assume $\gamma' \neq s'^n$.

$$w^n + o^n = \gamma'p^n$$

Where $\gamma' \neq s'^n$

Assume the greatest common factor of $2u$ and $u^{n-1} + n(v''^{n-1} + m'')$ is n.

Let $u=nr$.

u and r are relatively prime because u and v'' are relatively prime.

$$w^n + o^n = 2nr(u^{n-1} + n(v''^{n-1} + m''))$$

$$w^n + o^n = 2n^2r(n^{n-2}r^{n-1} + (v''^{n-1} + m''))$$

The greatest common factor of $2n^2r$ and $n^{n-2}r^{n-1} + (v''^{n-1} + m'')$ must be 1.

It is possible that $2n^2r = p'^n$ and $n^{n-2}r^{n-1} + (v'^{n-1} + l) = \delta'$.

$$w^n + o^n = \delta'p'^n$$

Assume $w^n = \delta'w'^n$, $o^n = \delta'o'^n$ or $w^n = \alpha''p'^n$, $o^n = \beta''p'^n$.

But this contradicts because w and o are relatively prime.

Assume $\delta' = S'^n$.

$$w^n + o^n = S'^n p'^n$$

Where S' and p' are relatively prime

Let $w^n = a''''^n$, $o^n = b''''^n$, $S'^n p'^n = c''''^n$

$$a''''^n + b''''^n = c''''^n$$

But this equation contradicts by the method of infinite descent.

Assume $\delta' \neq S'^n$.

$$w^n + o^n = \delta'p'^n$$

Where $\delta' \neq S'^n$

But this equation contradicts by the method of infinite descent.

Assume the greatest common factor of $2u$ and $u^{n-1} + n(v'^{n-1} + m')$ is n .

Let $u = nr$.

u and r are relatively prime because u and v' are relatively prime.

$$k^n + t^n = 2nr(u^{n-1} + n(v'^{n-1} + m'))$$

$$k^n + t^n = 2n^2r(n^{n-2}r^{n-1} + (v'^{n-1} + m'))$$

The greatest common factor of $2n^2r$ and $n^{n-2}r^{n-1} + (v'^{n-1} + m')$ must be 1.

It is possible that $2n^2r = p'^n$ and $n^{n-2}r^{n-1} + (v'^{n-1} + l) = \delta$.

$$k^n + t^n = \delta p'^n$$

Assume $k^n = \delta k''^n$, $t^n = \delta t''^n$ or $k^n = \alpha''p'^n$, $t^n = \beta''p'^n$.

But this contradicts because w and o are relatively prime.

Assume $\delta' = s'^n$.

$$k^n + t^n = s'^n p'^n$$

Where s' and p are relatively prime

Let $w^n = a''''^n, o^n = b''''^n, S^n p^n = c''''^n$

$$a''''^n + b''''^n = c''''^n$$

But this equation contradicts by the method of infinite descent.

Assume $\delta \neq s'^n$.

$$k^n + t^n = \delta p'^n$$

Where $\delta \neq s'^n$

Let $w = u - v'', o = u + v''$

Where $v'' = n v'$

Assume u and v'' are not relatively prime.

Let $u = fU, v'' = fV, V'' = fn V'$.

$$k = f(U - V''/n)$$

$$t = f(U + V''/n)$$

But this contradicts because k and t are relatively prime.

So, u and v'' are relatively prime.

Assume w and o are not relatively prime.

Let $u = fU, v = fV$.

$$w = f(U - V')$$

$$o = f(U + V')$$

But this contradicts because u and v'' are relatively prime.

So, w and o are relatively prime.

Also u and v'' are opposite parity because k and t are odd.

$$w^n + o^n = (u - v'')^n + (u + v'')^n$$

$$w^n + o^n = 2u(u^{n-1} + n(v''^{n-1} + m''))$$

Where $m'' = C'_{n-3} v''^{n-3} u^2 + C'_{n-5} v''^{n-5} u^4 + \dots + C'_4 v''^4 u^{n-5} + C'_2 v''^2 u^{n-3}$

Assume $u = \text{even}, v'' = \text{odd}$.

$u^{n-1} + n(v''^{n-1} + m'')$ is odd.

Assume u =odd, v'' =even.

$u^{n-1} + n(v''^{n-1} + m'')$ is odd.

So, $u^{n-1} + n(v''^{n-1} + m'')$ is always odd.

Thus the greatest common factor of $2u$ and $u^{n-1} + n(v''^{n-1} + m'')$ is odd.

Assume the common factor is odd number except 1 and n .

Let $u = fU$ and $u^{n-1} + n(v''^{n-1} + m'') = fN$.

$$f^{n-1}U^{n-1} + n(v''^{n-1} + fM'') = fN$$

Where $fM'' = f(C'_{n-3}v''^{n-3}f^1U^2 + C'_{n-5}v''^{n-5}f^3U^4 + \dots + C'_4v''^4f^{n-6}U^{n-5} + C'_2v''^2f^{n-4}U^{n-3})$

$$nv''^{n-1} = f(N - f^{n-2}U^{n-1} - nM'')$$

u and v'' have common factor of f .

But this contradicts because u and v'' are relatively prime.

So, the greatest common factor of $2u$ and $u^{n-1} + n(v''^{n-1} + m'')$ is either 1 or n .

Assume the greatest common factor of $2u$ and $u^{n-1} + n(v''^{n-1} + m'')$ is 1.

It is possible that $2u = p^n$ and $u^{n-1} + n(v''^{n-1} + m'') = \gamma'$.

$$w^n + o^n = \gamma'p^n$$

Assume $w^n = \gamma'w'^n$, $o^n = \gamma'o'^n$ or $w^n = \alpha'p^n$, $o^n = \beta'p^n$.

But this contradicts because k and t are relatively prime.

Assume $\gamma' = s'^n$.

$$w^n + o^n = s'^np^n$$

Where s' and p are relatively prime

Let $w^n = a'^n$, $o^n = b'^n$, $s'^np^n = c'^n$

$$a'^n + b'^n = c'^n$$

But this equation contradicts by the method of infinite descent.

Assume $\gamma' \neq s'^n$.

$$w^n + o^n = \gamma'p^n$$

Where $\gamma' \neq s'^n$

Assume the greatest common factor of $2u$ and $u^{n-1} + n(v''^{n-1} + m'')$ is n .

Let $u=nr$.

u and r are relatively prime because u and v'' are relatively prime.

$$w^n + o^n = 2nr(u^{n-1} + n(v''^{n-1} + m''))$$

$$w^n + o^n = 2n^2r(n^{n-2}r^{n-1} + (v''^{n-1} + m''))$$

The greatest common factor of $2n^2r$ and $n^{n-2}r^{n-1} + (v''^{n-1} + m'')$ must be 1.

It is possible that $2n^2r = p'^n$ and $n^{n-2}r^{n-1} + (v''^{n-1} + m'') = \delta'$.

$$w^n + o^n = \delta'p'^n$$

Assume $w^n = \delta'w'^n$, $o^n = \delta'o'^n$ or $w^n = \alpha'''p'^n$, $o^n = \beta'''p'^n$.

But this contradicts because w and o are relatively prime.

Assume $\delta' = S'^n$.

$$w^n + o^n = S'^n p'^n$$

Where S' and p' are relatively prime

Let $w^n = a''''n$, $o^n = b''''n$, $S'^n p'^n = c''''n$

$$a''''n + b''''n = c''''n$$

But this equation contradicts by the method of infinite descent.

Assume $\delta' \neq S'^n$.

$$w^n + o^n = \delta'p'^n$$

Where $\delta' \neq S'^n$

But this equation contradicts by the method of infinite descent.

Therefore, there are no positive integers in case of n =odd prime number since all of Case 1, 2 and 3 are contradiction.

5. REFERENCES

[1] A. Wilds. Modular elliptic curves and Fermat's last theorem. Ann. of Math., 141:443-551, 1995.