

Resolution of the Abraham–Minkowski Controversy

Andrew T. Hamilton*

The Abraham–Minkowski controversy has arisen because the Minkowski stress-energy tensor for the electromagnetic field in polarized and magnetized matter is not symmetric. This note proposes an alternative that is both symmetric and traceless.

I. INTRODUCTION

A. Conventions Adopted

The Minkowski metric is $(-, +, +, +)$.

$\nabla \cdot \vec{\sigma} \equiv \nabla_i \sigma_{ij}$ so that a stress-energy (or stress-energy-momentum) tensor takes the form

$$\begin{pmatrix} u & \mathbf{c}\mathbf{g} \\ \mathbf{S}/c & -\vec{\sigma} \end{pmatrix} \quad (1)$$

B. Background

The Abraham–Minkowski controversy is a century-long dispute as to the true form of the electromagnetic stress-energy tensor in and, hence, the electromagnetic force on matter that is polarized and magnetized.

It has its origin in 1908, when Minkowski derived a stress-energy tensor for the electromagnetic field in polarized and magnetized matter [1, Eq. 75],

$$\Theta^{\mu\nu} + \eta^{\mu\beta} P_{\alpha\beta} F^{\alpha\nu} - \frac{1}{4} \eta^{\mu\nu} P_{\alpha\beta} F^{\alpha\beta} \quad (2)$$

where $\Theta^{\mu\nu}$ is the (symmetric) electromagnetic stress-energy tensor, $\eta^{\mu\nu}$ is the (Minkowski) metric tensor, $P^{\mu\nu}$ is the polarization-magnetization tensor [2, Eq. 21], and $F^{\mu\nu}$ is the electromagnetic field tensor.

The controversy started the next year, when Abraham [3] derived relations with ‘... *eine merkwürdige Symmetrieeigenschaft jenes Gleichungssystems, die sich in Minkowski’s Ansätzen nicht findet.*’ The underlying reason for this is the source of the controversy: Minkowski’s stress-energy tensor is not symmetric.

Despite having been ‘resolved’ numerous times [4, 5], the controversy persists to this day. This humble contribution will make no difference to that situation.

II. DERIVATION

A. Electromagnetic Force Density

The Lorentz force density on *free* charges and currents is

$$(\nabla \cdot \mathbf{D})\mathbf{E} + \left(\nabla \times \mathbf{H} - \frac{\partial \mathbf{D}}{\partial t} \right) \times \mathbf{B} \quad (3)$$

which is the same as

$$-\partial_\mu \Theta^{\mu j} - (\partial^\beta P_{\alpha\beta}) F^{\alpha j} \quad (4)$$

where the Latin index j denotes spatial components.

The Lorentz force density on *bound* charges and currents is [6, Eq. 64]

$$(\mathbf{P} \cdot \nabla)\mathbf{E} + \frac{\partial \mathbf{P}}{\partial t} \times \mathbf{B} + \mathbf{M} \times (\nabla \times \mathbf{B}) + (\mathbf{M} \cdot \nabla)\mathbf{B} \quad (5)$$

which is *almost* the same as

$$-P_{\alpha\beta} (\partial^\beta F^{\alpha j}) \quad (6)$$

the difference being that the latter expression includes an additional term

$$-\frac{\partial}{\partial t} (\mathbf{P} \times \mathbf{B}) \quad (7)$$

Omitting this term would break the Lorentz covariance of Expression 6, which constitutes compelling evidence that it is real. However, it does not contribute to the Lorentz force density, so how to account for it?

B. Hidden Momentum

That $\mathbf{P} \times \mathbf{B}$ has the form of a momentum density suggests treating it as a component of the momentum density of matter.

According to Newton’s second law of motion, the force density on matter is

$$\partial_\mu \Upsilon^{\mu j} \quad (8)$$

where $\Upsilon^{\mu\nu}$ is the stress-energy tensor for matter. This is the same as

$$\frac{\partial \mathbf{g}_\Upsilon}{\partial t} - \nabla \cdot \vec{\sigma}_\Upsilon \quad (9)$$

where \mathbf{g}_Υ is the momentum density of the matter and $\vec{\sigma}_\Upsilon$ its stress tensor. If Expression 7 is a real force density, then the electromagnetic force density on matter is equal to the sum of Expressions 4 and 6,

$$\begin{aligned} \partial_\mu \Upsilon^{\mu j} &= -\partial_\mu \Theta^{\mu j} - (\partial^\beta P_{\alpha\beta}) F^{\alpha j} - P_{\alpha\beta} (\partial^\beta F^{\alpha j}) \\ &= -\partial_\mu \Pi^{\mu j} \end{aligned} \quad (10)$$

where

$$\Pi^{\mu\nu} = \Theta^{\mu\nu} + \eta^{\mu\beta} P_{\alpha\beta} F^{\alpha\nu} \quad (11)$$

* andrew.hamilton@bullagreen.com

is Medina and Stephany's stress-energy tensor for the electromagnetic field [9, Eq. 132]. The similarity to Minkowski's stress-energy tensor (Expression 2) is apparent.

Consequently, the Lorentz force density (the sum of Expressions 3 and 5) is equal to

$$\frac{\partial}{\partial t}(\mathbf{g}_T + \mathbf{P} \times \mathbf{B}) - \nabla \cdot \vec{\sigma}_T \quad (12)$$

and therefore, the effective momentum of a body of finite extent is

$$\int_{\mathcal{V}} (\mathbf{g}_T + \mathbf{P} \times \mathbf{B}) d\tau \quad (13)$$

which implies that an electric dipole \mathbf{p} in a magnetic field has linear momentum $\mathbf{p} \times \mathbf{B}$ irrespective of its motion.

This is an example of hidden momentum, a 'discovery' of Shockley and James [7], who found that magnetic dipoles possess "hidden momentum" $G_l = -\epsilon_0 E \times \mu_0 m$ associated with energy flow in a current loop of magnetic dipole strength m situated in an electric field E . They examined only magnetic dipoles, but the situation for electric dipoles is analogous.

However, this creates a conundrum in that, unlike a magnetic dipole, an electric dipole has no moving parts (e.g. moving charges) and hence nowhere to store momentum. So, where is it hidden? Minkowski [1] hides it in plain sight by incorporating it in the momentum density of the electromagnetic field. Griffiths [8] locates it in the sources of the magnetic field. I defer this question until the end.

C. Resolution

$\Pi^{\mu\nu}$, like Minkowski's stress-energy tensor, is problematic in that it:

1. is not symmetric
2. does not account for the hidden momentum of magnetic dipoles described by Shockley and James.

The remedy is to introduce a new stress-energy tensor

$$\Delta^{\mu\nu} = \eta^{\mu\beta} M_{\alpha\beta} G^{\alpha\nu} \quad (14)$$

where $M^{\mu\nu}$ and $G^{\mu\nu}$ are the duals of $P^{\mu\nu}$ and $F^{\mu\nu}$ respectively. If $T^{\mu\nu}$ is defined such that

$$\Upsilon^{\mu\nu} = T^{\mu\nu} + \eta^{\mu\beta} M_{\alpha\beta} G^{\alpha\nu} \quad (15)$$

which mirrors Equation 11, then Equation 10 is equivalent to

$$\partial_\mu T^{\mu j} = -\partial_\mu V^{\mu j} \quad (16)$$

where

$$V^{\mu\nu} = \Theta^{\mu\nu} + \eta^{\mu\beta} (P_{\alpha\beta} F^{\alpha\nu} + M_{\alpha\beta} G^{\alpha\nu}) \quad (17)$$

is symmetric and traceless. The momentum density component of $\Delta^{\mu\nu}$ is

$$\frac{1}{c} \mathbf{M} \times \mathbf{E} = c \mathbf{g}_\Delta \quad (18)$$

so that the effective momentum of a body of finite extent becomes

$$\begin{aligned} & \int_{\mathcal{V}} (\mathbf{g}_T + \mathbf{g}_\Delta + \mathbf{P} \times \mathbf{B}) d\tau \\ &= \int_{\mathcal{V}} \left(\mathbf{g}_T + \frac{1}{c^2} \mathbf{M} \times \mathbf{E} + \mathbf{P} \times \mathbf{B} \right) d\tau \end{aligned} \quad (19)$$

where $c\mathbf{g}_T$ is the momentum density component of $T^{\mu\nu}$. This leads to the following interpretations.

1. That $\Upsilon^{\mu\nu}$ is the stress-energy tensor for matter means that $\Pi^{\mu\nu}$ must be the stress-energy tensor for the field.
2. $T^{\mu\nu}$ is the stress-energy tensor for matter prior to its being polarized or magnetized, so $\Delta^{\mu\nu}$ is the change in the stress-energy tensor for matter when it is polarized and magnetized.
3. Equation 16 is the (kinetic) electromagnetic force density.

Returning to the issue of the location of the hidden momentum of electric dipoles, by the first interpretation it is present in the momentum density of the field, which is $\mathbf{D} \times \mathbf{B}$, not that of matter.

III. CONCLUSION

Barnett [5] suggests that the '... resolution of the Abraham-Minkowski dilemma requires us to recognize that there are two distinct electromagnetic momenta, the kinetic momentum and the canonical momentum.'

In this he is correct. There are two distinct electromagnetic stress-energy tensors:

1. $V^{\mu\nu}$ is the kinetic stress-energy tensor, which is symmetric and traceless
2. $\Pi^{\mu\nu}$ is the canonical stress-energy tensor, which is the stress-energy tensor for the electromagnetic field.

In the absence of polarization and magnetization both are equal to the familiar electromagnetic stress-energy tensor, $\Theta^{\mu\nu}$.

ACKNOWLEDGEMENT

I thank David J. Griffiths for drawing my attention to his paper [8].

-
- [1] H. Minkowski, Die Grundgleichungen für die elektromagnetischen Vorgänge in bewegten Körpern, Nachrichten von der Königlichen Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse **1908**, 53.
- [2] D. J. Griffiths and V. Hnizdo, Mansuripur's paradox, Am. J. Phys. **81**, 570 (2013), arXiv:1303.0732 [physics.class-ph].
- [3] M. Abraham, Zur Elektrodynamik bewegter Körper, Rend. Circ. Matem. Palermo **28**, 1 (1909).
- [4] W. Shockley, A "Try Simplest Cases" resolution of the Abraham–Minkowski controversy on electromagnetic momentum in matter, Proc. Natl. Acad. Sci. U.S.A. **60**, 807 (1968); D. F. Nelson, Resolution of the problem of Minkowski and Abraham, in *Optical Society of America Annual Meeting* (Optica Publishing Group, 1990) p. FMM2; M. Mansuripur, Resolution of the Abraham–Minkowski controversy, Opt. Commun. **283**, 1997 (2010), arXiv:1208.0872 [physics.optics]; B. A. Kemp, Resolution of the Abraham–Minkowski debate: Implications for the electromagnetic wave theory of light in matter, J. Appl. Phys. **109**, 111101 (2011); J. Chen, Y. Dai, and Y. Xuanyuan, Possible solution of Abraham–Minkowski controversy by generalizing the principle of invariance of light speed, J. Opt. **49**, 127 (2020); A. S. Al-Ithawi, New theory of light and resolution of the Abraham–Minkowski controversy, Walailak J. Sci. Technol. **17**, 1060 (2020); G. Feng and J. Huang, A heuristic resolution of the Abraham–Minkowski controversy, Eur. Phys. J. Plus **136**, 520 (2021), arXiv:2101.01004 [physics.optics]; G.-X. Liang, The energy-momentum tensor of electromagnetic waves in media and the solution to the Abraham–Minkowski dilemma, arXiv:2309.03210 [physics.gen-ph] (2023).
- [5] S. M. Barnett, Resolution of the Abraham–Minkowski dilemma, Phys. Rev. Lett. **104**, 070401 (2010).
- [6] D. J. Griffiths and V. Hnizdo, What's the use of bound charge?, arXiv:1506.02590 [physics.class-ph] (2015).
- [7] W. Shockley and R. P. James, "Try Simplest Cases" discovery of "Hidden Momentum" forces on "Magnetic Currents", Phys. Rev. Lett. **18**, 876 (1967).
- [8] D. J. Griffiths, A catalogue of hidden momenta, Phil. Trans. R. Soc. A **376**, 10.1098/rsta.2018.0043 (2018).
- [9] R. Medina and J. Stephany, The energy-momentum tensor of electromagnetic fields in matter, arXiv:1703.02109 [physics.class-ph] (2017).