

# On The Dynamics Of The Gravitational Field And Higher Order Field Equations

D.E. Thomas

(Dated: May 17, 2025)

## Abstract

Einstein's General Relativity, while extraordinarily successful in explaining gravitational phenomena, faces challenges in reconciling with quantum mechanics and addressing certain cosmological observations. I present a higher order model of gravitation that reframes the gravitational field in terms of the connection, with the curvature tensor as the field strength. The action in this model is based on the squared norm of the field strength, mirroring the structure of other Standard Model interactions. This approach offers potential advantages for quantization and renormalizability. Through analysis of the Euler-Lagrange equations in the gravitational sector, particularly focusing on the metric equations, I demonstrate that Einstein's field equations emerge as a limiting case within a more comprehensive theoretical framework. This broader theory may provide explanations for observed experimental discrepancies in cosmology while maintaining consistency with established gravitational phenomena. This model thus offers a promising avenue for addressing current challenges in gravitational physics and cosmology.

## INTRODUCTION

Einstein’s field equations (EFEs), the cornerstone of General Relativity (GR), have stood as one of the most profound and successful descriptions of gravity for over a century. These elegant equations relate the geometry of spacetime to the distribution of matter and energy within it, providing a framework for understanding gravitational phenomena with a precision far exceeding that of any model previous to their conception [1, 2].

However, as our observational capabilities have advanced and our theoretical understanding has deepened, several critical challenges to the completeness of the EFEs have emerged. These challenges span both the cosmic and quantum scales, highlighting the need for a more comprehensive theory of gravity [3].

**The Cosmological Constant Problem:** A staggering discrepancy of significant orders of magnitude exists between quantum field theory predictions for vacuum energy and the observed cosmological constant. This discrepancy has been dubbed “the worst theoretical prediction in the history of physics”. It suggests a fundamental gap in our understanding of how vacuum energy interacts with gravity at cosmological scales [4, 5].

**Dark Matter and Dark Energy:** The necessity of dark matter to explain galactic rotation curves and dark energy to account for the universe’s accelerating expansion suggests that the EFEs may not fully describe gravitational phenomena on cosmological scales. These additional components, constituting most of the universe’s content, are not predicted by the original equations [6, 7]. This discrepancy points to a significant incompleteness in our current gravitational theory.

**Quantum Gravity:** The EFEs, while extraordinarily successful in describing gravity at macroscopic scales, are fundamentally classical in nature. They do not incorporate quantum mechanics, leading to breakdowns at very small scales and high energies, such as those found in black holes or the early universe. Moreover, GR has resisted attempts at quantization, suggesting that a more fundamental theory—one that is naturally amenable to quantization and renormalization—may be necessary [8–10].

In this paper, I present a mathematical framework that builds upon Einstein’s foundational work while addressing these critical challenges. My approach aims to reconcile the observed cosmological differences with theoretical predictions without resorting to the introduction of exotic forms of matter or energy. Furthermore, it lays the groundwork for a

theory that is more naturally suited to quantization, potentially bridging the gap between gravity and quantum mechanics.

In the following section, I will introduce the mathematical framework, which extends the EFEs in a way that naturally accommodates the observed cosmological phenomena. This approach paves the way for a more complete description of gravitational interactions across all scales—from the quantum realm to the largest structures in the universe.

## I. FROM EINSTEIN TO A MODERN FIELD THEORY

### Einstein Field Equations

When Einstein first wrote down the field equations of his general theory of relativity in action form, he proposed the Lagrangian [1, 11],

$$L = g^{\mu\nu} \begin{Bmatrix} \alpha \\ \mu\beta \end{Bmatrix} \begin{Bmatrix} \beta \\ \nu\alpha \end{Bmatrix} \quad (1)$$

where  $g^{\mu\nu}$  is the metric tensor and  $\begin{Bmatrix} \alpha \\ \mu\beta \end{Bmatrix}$  is the Christoffel symbol defined as,

$$\begin{Bmatrix} \alpha \\ \mu\beta \end{Bmatrix} = \frac{1}{2} g^{\alpha\sigma} (\partial_\beta g_{\sigma\mu} + \partial_\mu g_{\sigma\beta} - \partial_\sigma g_{\mu\beta}) \quad (2)$$

Later, Einstein, Hilbert, and others reformulated the action of the theory as [2, 12],

$$S = \int R \sqrt{-g} d^4x \quad (3)$$

where  $R = R_{\mu\beta\nu}^\alpha g_\alpha^\beta g^{\mu\nu}$ ,

$R_{\mu\beta\nu}^\alpha$  is the Riemann curvature tensor,

$R$  is the curvature scalar,

$g$  is the determinant of the metric,  $\sqrt{-g} d^4x$  the canonical invariant volume element on a manifold. Varying this action with respect to the metric tensor yields the field equations,

$$R_{\mu\beta\nu}^\alpha g_\alpha^\beta = R_{\mu\nu} = 0 \quad (4)$$

here,  $R_{\mu\nu}$  is called the Ricci curvature tensor.

This result, however, was derived assuming that the spacetime is torsion-free [1, 2, 11, 13]. Including torsion  $\mathcal{T}$ , the full affine connection is expressed as,

$$\Gamma_{\mu\nu}^{\alpha} = \frac{1}{2}g^{\alpha\sigma}(\partial_{\nu}g_{\sigma\mu} + \partial_{\mu}g_{\sigma\nu} - \partial_{\sigma}g_{\mu\nu} + \mathcal{T}_{\sigma\mu\nu} + \mathcal{T}_{\mu\nu\sigma} - \mathcal{T}_{\nu\sigma\mu}) \quad (5)$$

In any holonomic basis, the affine connection completely describes the covariant derivative up to metric compatibility [2]. Recognizing this, Cartan showed that varying the action of GR with respect to torsion yields the torsion tensor field equations [14],

$$\mathcal{T}_{\mu\nu}^{\alpha} + g_{\mu}^{\alpha}\mathcal{T}_{\nu\sigma}^{\sigma} - g_{\nu}^{\alpha}\mathcal{T}_{\mu\sigma}^{\sigma} = 0 \quad (6)$$

Here one must find themselves at a crossroads. The field equations of Einstein-Cartan theory (4 & 6) are found by varying with respect to the two parts of the connection independently, while treating the metric and torsion each as gravitational potential fields. One should ask: why not treat the entire affine connection as the gravitational potential and find an invariant action to vary with respect to the field represented by the affine connection?

An action can be constructed in a form similar to that seen in other field theories. Specifically, I take a first-order derivative of the dynamic field and square it appropriately so as to ensure invariance with respect to some local symmetry [15, 16], in this case, general coordinate transformations.

In examining equation (1), I notice it appears that Einstein may have initially approached his model with this formulation in some manner, considering he treated the metric as the gravitational potential. However, given the limited exploration and popularization of this approach at the time, it is understandable that he was persuaded to adopt a more minimal scalar quantity. While this historical interpretation is intriguing, it is purely speculative on my part and not well-documented.

### Construction of The Affine Field Equations

I will start with the affine connection and take a derivative as follows,

$$\mathcal{R}_{\mu\beta\nu}^{\alpha} = \partial_{\beta}\Gamma_{\nu\mu}^{\alpha} - \partial_{\nu}\Gamma_{\beta\mu}^{\alpha} + \Gamma_{\beta\sigma}^{\alpha}\Gamma_{\nu\mu}^{\sigma} - \Gamma_{\nu\sigma}^{\alpha}\Gamma_{\beta\mu}^{\sigma} + \Gamma_{\beta\nu}^{\sigma}\Gamma_{\sigma\mu}^{\alpha} - \Gamma_{\nu\beta}^{\sigma}\Gamma_{\sigma\mu}^{\alpha} \quad (7)$$

$\mathcal{R}_{\mu\beta\nu}^{\alpha}$  is the complete curvature tensor in the presence of torsion or any other antisymmetric contributions to the connection [13].

$$\mathcal{R}_{\mu\beta\nu}^{\alpha} = R_{\mu\beta\nu}^{\alpha} + \mathcal{T}_{\beta\nu}^{\sigma}\Gamma_{\sigma\mu}^{\alpha} \quad (8)$$

Let us define  $\mathcal{R}_{\mu\beta\nu}^\alpha$  as the gravitational field strength. The Lagrangian density will be built from the field strength in a manner similar to modern geometric field theories [15, 16], by taking the inner product of the field strength with itself,

$$S = \int \mathcal{R}_{\mu\beta\nu}^\alpha \mathcal{R}_\alpha^{\mu\beta\nu} D^4x \quad (9)$$

where  $D^4x$  is a generalized invariant volume form. This Lagrangian and its associated Euler-Lagrange equations with respect to the gravitational field  $\Gamma$  will yield the gravitational field equations,

$$\frac{\partial}{\partial \Gamma_{\nu\mu}^\alpha} (\mathcal{R}_{\mu\beta\nu}^\alpha \mathcal{R}_\alpha^{\mu\beta\nu}) - \partial_\rho \frac{\partial}{\partial (\partial_\rho \Gamma_{\nu\mu}^\alpha)} (\mathcal{R}_{\mu\beta\nu}^\alpha \mathcal{R}_\alpha^{\mu\beta\nu}) = 0 \quad (10)$$

the *field equations* are,

$$D_\beta \mathcal{R}_\alpha^{\mu\beta\nu} = 0 \quad (11)$$

In addition to the field equations, one must pair with *equations of motion* to provide a description of the motion of a test particle in the field solved for in the field equations. In Einstein's general theory of relativity, the geodesic equation serves the role of this equation of motion [17–19],

$$\frac{d^2x^\mu}{d\lambda^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\lambda} \frac{dx^\beta}{d\lambda} = 0 \quad (12)$$

This is found by varying the path interval with respect to the position of a free particle. In the model presented here, the path interval for a free particle is given by,

$$S = \int mc^2 D\lambda \quad (13)$$

where  $D\lambda$  is an invariant path element such that,

$$D\lambda \rightarrow \frac{D\lambda}{dx^\mu} dx^\mu \rightarrow d\lambda + \Gamma_{\mu\nu}^\mu \frac{dx^\nu}{d\lambda} d\lambda \quad (14)$$

For a discussion on how this is similar to the canonical invariant element ( $\sqrt{-g}d\lambda$ ), but more generalized, and a comparison with other models see Appendix A. Varying with respect to  $x^\mu$ , one finds,

$$\frac{d^2x^\alpha}{d\lambda^2} + R_{\beta\mu\nu}^\alpha \frac{dx^\beta}{d\lambda} \frac{dx^\mu}{d\lambda} X^\nu = 0 \quad (15)$$

where  $X$  is the separation vector between events on geodesics, this is canonically called the geodesic deviation [20, 21].

## II. METRIC FIELD EQUATIONS OF THE SQUARED NORM

### Field Equations of the Metric

The model presented establishes the gravitational field equations and equations of motion. However, in Einstein's GR he considers the metric field equations [1, 11]. I will examine the metric field equations in the presented model to compare with Einstein's equations and present any corrections to the Einstein model in the regime where it could be considered a valid approximation.

Using the action based on the field Lagrangian density  $\mathcal{R}_{\mu\beta\nu}^{\alpha}\mathcal{R}_{\alpha}^{\mu\beta\nu}$ , we can also derive the field equations of the metric,

$$2D_{[a}D_{b]}\mathcal{R}^{nmab} + \mathcal{R}_{ab}\mathcal{R}^{namb} = 0 \tag{16}$$

for a detailed derivation of these metric field equations using a variational principle approach see Appendix B.

In the rest of this paper I will generalize the notion of the curvautre tensor such that I refer to  $\mathcal{R}$  as the Riemann tensor, it's contraction the Ricci tensor, and the trace of this form of the Ricci tensor as the curvature scalar.

### The Einsteinian Limit

Under what conditions would one expect this to reduce to the Einstein equations, where we can consider the Einstein equations as an approximation? I expect that the Einstein equations would emerge as a weak field limit of (16), when the local curvature becomes negligible. I would not necessarily restrict the global curvature, however. These conditions correspond to a maximally symmetric space, i.e., where the curvature is isotropic and homogeneous everywhere in the universe [22–24]. This would imply that in regions where the gravitational field is weak, we can ignore local effects on the geometry and focus on the solutions for the global geometry [25], which I hypothesize can be reduced to Einstein's equations.

In a maximally symmetric space the metric is not a function of space, and in a nearly maximally symmetric space the metric can be decomposed into a maximally symmetric

background and a local part,

$$g_{ab}(x) = \eta_{ab} + h_{ab}(x) \quad (17)$$

where  $\eta$  is the maximally symmetric metric (this could be  $\eta^0$  the metric of a ‘flat’ Minkowski spacetime,  $\eta^+$  the metric of a ‘positively curved’ de Sitter space, or  $\eta^-$  the metric of a ‘negatively curved’ anti-de Sitter space). The global geometry of spacetime is captured in  $\eta$  and the local geometry is described by  $h(x)$ . This approach, the background field method, is commonly used in perturbative treatments of gravity and quantum field theory in curved spacetime [26, 27]. It allows us to separate the large-scale structure of spacetime from local fluctuations or perturbations. The term  $h(x)$  represents small deviations from the maximally symmetric background. In the context of GR, these deviations represent localized gravitational effects. Most commonly, these localized effects are used to describe gravitational waves [28]. I do not take that wave interpretation here since gravitational waves should be fully described by the gravitational field equations for  $\Gamma$ . Instead, I view this as local deviations from the background metric due to the local mass content in a region of spacetime.

The Riemann tensor can be decomposed in general [29],

$$\mathcal{R}_{abcd} = C_{abcd} + \frac{1}{2} (g_{a[c}\mathcal{R}_{d]b} - g_{b[c}\mathcal{R}_{d]a}) - \frac{1}{6}g_{a[c}g_{d]b}\mathcal{R} \quad (18)$$

where  $C_{abcd}$  is canonically called the Weyl tensor. The Weyl tensor vanishes in maximally symmetric spaces [30], so one obtains,

$$\mathcal{R}_{abcd}(\eta) = \frac{1}{2} (\eta_{a[c}\mathcal{R}_{d]b}(\eta) - \eta_{b[c}\mathcal{R}_{d]a}(\eta)) - \frac{1}{6}\eta_{a[c}\eta_{d]b}\mathcal{R} \quad (19)$$

furthermore, the Ricci tensor in a maximally symmetric space is trivially  $\mathcal{R}_{ab}(\eta) = \frac{1}{4}\eta_{ab}\mathcal{R}$ . Therefore, the Riemann tensor in a maximally symmetric space becomes,

$$\mathcal{R}_{abcd}(\eta) = \frac{1}{12}\eta_{a[c}\eta_{d]b}\mathcal{R} \quad (20)$$

thus, the field equations can be simplified as follows,

$$2D_{[a}D_{b]} \left( \frac{1}{12}\eta^{n[m}\eta^{a]b}\mathcal{R} \right) + \left( \frac{1}{4}\eta_{ab}\mathcal{R} \right) \left( \frac{1}{12}\eta^{n[a}\eta^{m]b}\mathcal{R} \right) = 0 \quad (21)$$

notice that neither  $\eta$  nor  $\mathcal{R}$  are functions of space in this context, and therefore the derivative term vanishes,

$$D_{[a}D_{b]} (\eta^{n[m}\eta^{a]b}\mathcal{R}) = 0$$

the remaining part of the field equations must then also vanish in this context,

$$\begin{aligned} \left(\frac{1}{4}\eta_{ab}\mathcal{R}\right)\left(\frac{1}{12}\eta^{n[a}\eta^{m]b}\mathcal{R}\right) &= 0 \\ \frac{1}{16}\eta^{nm}\mathcal{R}^2 &= 0 \end{aligned} \quad (22)$$

The proposal of Einstein's equations as a weak-field limit under these conditions would imply the following,

$$\frac{1}{16}\eta^{ab}\mathcal{R}^2 = G^{ab} + H^{ab} \quad (23)$$

where  $G^{ab}$  is the Einstein tensor, and  $H^{ab}$  represents a correction. The Einstein tensor in a maximally symmetric vacuum [31] is given by,

$$\begin{aligned} G^{ab}(\eta) &= \mathcal{R}^{ab}(\eta) \\ &= \frac{1}{4}\eta^{ab}\mathcal{R} \end{aligned} \quad (24)$$

from equations 23 and 24 one can conclude that,

$$\begin{aligned} H^{ab} &= \frac{1}{16}\eta^{ab}\mathcal{R}^2 - \frac{1}{4}\eta^{ab}\mathcal{R} \\ H^{ab} &= \frac{1}{4}\eta^{ab}\mathcal{R}\left(\frac{\mathcal{R}}{4} - 1\right) \\ H^{ab} &= \mathcal{R}^{ab}\left(\frac{\mathcal{R}}{4} - 1\right) \end{aligned} \quad (25)$$

when Einstein's equations do not require corrections  $H^{ab} \approx 0$  and the right hand side of (25) should also vanish,

$$\mathcal{R}^{ab}\left(\frac{\mathcal{R}}{4} - 1\right) \approx 0 \quad (26)$$

therefore either,

$$\mathcal{R}\eta^{ab} \approx 0$$

or

$$\frac{\mathcal{R}}{4} - 1 \approx 0$$

however, since the metric is not generally null either the scalar curvature is vanishing,  $\mathcal{R} \approx 0$ , or  $\mathcal{R} \approx 4$ , for Einstein's field equations to require no corrections.

### III. COMPARISON TO EMPIRICAL COSMOLOGY

#### Global Curvature and Dark Energy

On scales large enough that the maximally symmetric condition holds, one can examine the differences in the expected and observed average curvature of the universe in the form of the cosmological constant  $\Lambda$ . Since this is an observed quantity, slightly positive number near zero [6], I expect the correction  $H$  might be sufficient to account for the effects currently attributed to dark energy, potentially eliminating the need to invoke exotic explanations for cosmic expansion.

Consider the corrections  $H$  to be equivalent to  $\Lambda\eta$  when comparing the EFEs with a cosmological constant to the metric field equations. In this context, it is evident that the metric field equations do not require additional factors to account for the phenomena typically attributed to dark energy,

$$\begin{aligned}
 H_{ab} &\stackrel{!}{=} \Lambda\eta_{ab} \\
 \mathcal{R}_{ab}(\eta) \left( \frac{\mathcal{R}}{4} - 1 \right) &= \Lambda\eta_{ab} \\
 \frac{1}{4}\mathcal{R}\eta_{ab} \left( \frac{\mathcal{R}}{4} - 1 \right) &= \Lambda\eta_{ab} \\
 \mathcal{R}^2 - 4\mathcal{R} &= 16\Lambda
 \end{aligned} \tag{27}$$

solving for  $\mathcal{R}$ ,

$$\mathcal{R} \approx 2 \pm 2\sqrt{1 - 4\Lambda} \tag{28}$$

one can approximate  $\sqrt{1 - x} \approx 1 - \frac{x}{2}$  for small  $x$ , therefore,

$$\mathcal{R} \approx 2 \pm (2 - 4\Lambda) \Rightarrow \begin{cases} 4\Lambda \\ 4(1 - \Lambda) \end{cases} \tag{29}$$

A scalar curvature of  $\mathcal{R} \approx 4(1 - \Lambda) \approx 4$  corresponds to a mass density on the order of  $10^{10}kg/m^3$ , which is approximately that of a white dwarf star [32]. This represents an extremely large average curvature/mass density for a maximally symmetric universe and is therefore ruled out as non-physical. I now turn to the case of vanishing scalar curvature as the other solution where Einstein's equations require no correction.  $R \approx 4\Lambda \approx 0$  aligns with current observations [33]. Thus, this model naturally produces a term that behaves like the

cosmological constant, but replaces it with purely geometric terms. This could explain the phenomenon currently attributed to dark energy without the need to invoke exotic materia.

### Local Curvature and Dark Matter

The corrections due to  $h(x)$  from a specific mass will require an analysis of the perturbations in the field equations by considering a non-vanishing  $h$ . To begin this process, I will express the field equations in terms of the metric decomposition given in equation (17),

$$\mathcal{R}_{abcd} = \mathcal{R}_{abcd}(\eta) + \mathcal{R}_{abcd}(h) + \mathcal{R}_{abcd}(\mathcal{O}(h)) \quad (30)$$

where  $\mathcal{R}_{abcd}(\eta)$  is the curvature due to only the background metric (as found in equation 20),  $\mathcal{R}_{abcd}(h)$  is the curvature arising from the first-order corrections due to  $h$ , and  $\mathcal{R}_{abcd}(\mathcal{O}(h))$  represents the curvature corrections of second order or higher in  $h$ , which I will treat as small and negligible. Let's now write out all the terms of the connection to first order in  $h$ ,

$$\Gamma_{abc}(h) = \frac{1}{2} (\partial_b h_{ac} + \partial_c h_{ba} - \partial_a h_{bc}) \quad (31)$$

now the curvature to first order in  $h$ , neglecting the quadratic terms  $\Gamma\Gamma$  (since they are at minimum second order in  $h$  and therefore negligible for small  $h$ ), can be expressed as,

$$\begin{aligned} \mathcal{R}_{abcd}(h) &= \partial_c \Gamma_{abd}(h) - \partial_d \Gamma_{abc}(h) \\ \mathcal{R}_{abcd}(h) &= \frac{1}{2} (\partial_c \partial_b h_{ad} + \partial_c \partial_d h_{ba} - \partial_c \partial_a h_{bd} - \partial_d \partial_b h_{ac} - \partial_a \partial_c h_{ba} + \partial_d \partial_a h_{bc}) \end{aligned} \quad (32)$$

$$\mathcal{R}_{ab}(h) = \frac{1}{2} (\partial^n \partial_{(a} h_{b)n} - \square h_{ab} - \partial_b \partial_a tr(h)) \quad (33)$$

where  $\square$  is the d'Alembertian,  $\partial^n \partial_n$ , and  $tr(h)$  is the trace of  $h$ , i.e.  $h_n^n$ .

I am going to substitute this result (33) into the metric field equations (16), but because I am only considering very small  $h$  the derivatives  $D_{[a} D_{b]} \mathcal{R}^{nmab}(\eta)$  are negligible. So substituting (33) into just the second part of (16),

$$\begin{aligned} &\frac{1}{4} (\partial^n \partial_{(a} h_{b)n} - \square h_{ab} - \partial_b \partial_a tr(h)) \\ &\times (\partial^m \partial^a h^{nb} + \partial^m \partial^b h^{an} - \partial^m \partial^n h^{ab} - \partial^b \partial^a h^{nm} - \partial^b \partial^m h^{an} + \partial^b \partial^n h^{am}) = 0 \end{aligned} \quad (34)$$

When the first term in (34) is equal to zero, it is not novel because it arises in standard GR from the EFEs representing a well-established result consistent with the predictions of GR in weak-field limits or in maximally symmetric spaces [34]. Novel solutions to the metric will be found by looking at the second part of (34).

*Modified Schwarzschild Metric*

I want to find solutions to the metric that satisfy,

$$\partial^m \partial^a h^{nb} + \partial^m \partial^b h^{an} - \partial^m \partial^n h^{ab} - \partial^b \partial^a h^{nm} - \partial^b \partial^m h^{an} + \partial^b \partial^n h^{am} = 0 \quad (35)$$

where  $h$  vanishes at infinity. The general solution will take the form,

$$h^{ab} = e^{-\alpha r} A^{ab} \quad (36)$$

In the simplest case of the metric outside a spherically symmetric mass, the background metric is that of Schwarzschild [35] and the perturbation would give a correction to each part of the background,

$$h^{ab} = \eta^{ab} A e^{-\alpha r} \quad (37)$$

$$h = \begin{pmatrix} h_1(r) & 0 & 0 & 0 \\ 0 & h_2(r) & 0 & 0 \\ 0 & 0 & h_3(r) & 0 \\ 0 & 0 & 0 & h_4(r) \end{pmatrix} \quad (38)$$

the time and radial components,  $h_1$  and  $h_2$ ,

$$h_1(r) = -\frac{1}{h_2(r)} = A e^{-\alpha r} \quad (39)$$

Adding this to the Schwarzschild metric,

$$ds^2 = - \left( 1 - \frac{2GM}{c^2 r} - A e^{-\alpha r} \right) c^2 dt^2 + \left( 1 - \frac{2GM}{c^2 r} - A e^{-\alpha r} \right)^{-1} dr^2 + r^2 d\Omega^2 \quad (40)$$

$d\Omega$  captures the metric on the 2-sphere including any contributions that might arise from  $h_3$  and  $h_4$ .

*Modified Newtonian Potential*

Based on the corrections to the Schwarzschild metric found, the gravitational potential is modified with corrections to the Newtonian potential,

$$\Phi(r) = -\frac{GM}{r} - \frac{c^2}{2} A e^{-\alpha r} \quad (41)$$

observational limits on the value of  $A$  and  $\alpha$  in the perihelion of Mercury [36], orbiting binary pulsar timing [37, 38], deflection of light around the sun and other massive bodies [36], and gravitational wave propagation exist to constrain these parameters in order to agree with current data [39–41]. I wish to explain all the effects we see in cosmology without the need to invoke dark matter. This requires the exponential term dominates the potential for  $r$  greater than a characteristic distance  $r_d$ , where  $r_d$  is the distance where the transition from being dominated by the Newtonian to being dominated by the corrections occurs. Setting the terms equal at  $r = r_d$  to find the characteristic distance there the transition occurs,

$$\frac{GM}{r_d} = \frac{c^2}{2} A e^{-\alpha r_d} \quad (42)$$

$$r_d = \frac{2GM}{c^2 A} e^{\alpha r_d} \quad (43)$$

$$\ln(r_d) = \ln\left(\frac{2GM}{c^2 A}\right) + \alpha r_d \quad (44)$$

let  $A = \left(\frac{2GM}{c^2}\right)$  and solve for  $\alpha$  at the characteristic distance  $r_d$ ,

$$\alpha = \frac{\ln(r_d)}{r_d} \quad (45)$$

so,

$$\Phi(r) = -\frac{GM}{r} - GM e^{-\frac{\ln(r_d)}{r_d} r} \quad (46)$$

Notice for small masses at short distances the  $1/r$  term is a good approximation, but for large  $M$  the exponential term dominates at large distances. The characteristic distance  $r_d$  becomes an undetermined free parameter to be measured to satisfy experimental data.

### *Modified Keplerian Orbital Velocities*

One place the Newtonian potential is insufficient to explain the gravitational effects of the visible baryonic matter is in the orbital velocities in spiral galaxies [42, 43]. Modifying the Keplerian expectation based on the modified potential I found might help explain the difference in theory and measurement. The modified orbital velocity due to this modified potential are,

$$v = \sqrt{\frac{GM}{r} + GM r \frac{\ln(r_d)}{r_d} e^{-\frac{\ln(r_d)}{r_d} r}} \quad (47)$$




FIG. 1. Plot of Orbital Velocities vs.  $r$

the modified orbital velocity tends to the Keplerian expectation for large or small  $r$ , but around  $r_d$  the orbital velocity has a bump (FIG. 1). Observations of orbital velocities show that velocities do not fall off purely exponentially as is the Keplerian expectation, but rather flatten out at some distance near the outer edges of the galactic spiral [44, 45]. The bump in orbital velocity calculated with corrections could align with the cosmological observations for the right choice of  $r_d$ . The exact value of  $r_d$  for a given galaxy is a free parameter to be determined exactly by experiment. The calculated orbital velocities will need to be compared to empirical data to determine if the expected and measured orbital velocities match without the need to add additional sources of mass such as dark matter.

#### IV. THIS MODEL OF GRAVITY AS A GAUGE THEORY

I would like to take this opportunity to present a brief review of the model presented and its comparison to the differential geometry of Standard Model gauge theory, highlighting why it might be better prepared for quantization compared to GR. This model reframes gravity in terms of the affine connection as the fundamental field, with the curvature tensor serving as the field strength. This approach bears striking similarities to the gauge theories of the Standard Model. In this model, the gravitational potential is represented by the affine

connection  $\Gamma$ , analogous to the gauge potential  $A$  in Yang-Mills theories [15, 17–19]. The curvature tensor  $R$  serves as the gravitational field strength, similar to the field strength tensor  $F$  in gauge theories. The equations based on the squared norm of the field strength  $(\mathcal{R}_{\mu\beta\nu}^\alpha \mathcal{R}_\alpha^{\mu\beta\nu})$ , mirrors the Yang-Mills action  $(F_{\mu\nu}^a F_a^{\mu\nu})$  [15].

This formulation offers several advantages for quantization. The model naturally incorporates the geometric nature of gravity while maintaining a structure similar to other quantum field theories. The squared field strength in the action is a promising starting point for renormalization, as it is of the same form as renormalizable gauge theories. The model respects general coordinate invariance, analogous to gauge invariance in the Standard Model [17–19]. The background field method used in this model ( $g_{ab} = \eta_{ab} + h_{ab}$ ) is well-suited for perturbative quantum calculations [27]. This approach potentially allows for a more unified treatment of gravity with other fundamental forces.

In contrast, GR’s action (based on the Ricci scalar  $R$ ) and its direct quantization face several challenges. GR is notoriously difficult to renormalize due to its non-polynomial dependence on the metric [46]. Traditional approaches to quantizing GR often struggle with background independence [47–50]. Perturbative expansions in GR can lead to divergences that are challenging to handle [51]. My model’s closer alignment with the structure of gauge theories potentially offers a more promising path towards a quantum theory of gravity, addressing some of the key obstacles faced in quantizing GR directly.

## Appendix A: About the Invariant Element

In standard GR the invariant element is commonly written as  $\sqrt{-g}d\lambda$ . Writing  $D\lambda$  is more than merely a suppression of the invariance into a different symbolic notation. I write this to explicitly display it as similar to the covariant derivative,

$$D\lambda = d\lambda + \Gamma_{\mu\nu}^\mu \frac{dx^\nu}{d\lambda} d\lambda$$

notice the similarity to the commonly used version through the following,

$$\frac{d}{dx^\nu} \sqrt{-g} = \Gamma_{\mu\nu}^\mu$$

so, this version is equivalent in the case used in this paper. But this version is more versatile and ready to be used in a unified field theory where the invariant element is more complicated, and the advantage comes in the way of using the covariant derivative to expand one’s

understanding,

$$D_\mu V^\nu = \frac{dV^\nu}{dx^\mu} + \Gamma_{\mu\alpha}^\nu V^\alpha + A_\mu V^\nu$$

$$D\lambda = d\lambda + \Gamma_{\mu\nu}^\mu \frac{dx^\nu}{d\lambda} d\lambda + A_\nu \frac{dx^\nu}{d\lambda} d\lambda$$

where  $A$  is another connection such as that of a gauge group in the Standard Model.

## Appendix B: Derivation of the Metric Field Equations of the Squared Norm

$$\mathcal{L} = \mathcal{R}_{abcd} R^{abcd}$$

$$\delta\mathcal{L} = 2\delta\mathcal{R}_{abcd} \cdot R^{abcd}$$

$$\delta\mathcal{R}_{abcd} = D_c\delta\Gamma_{abd} - D_d\delta\Gamma_{abc}$$

$$\delta\Gamma_{abc} = \frac{1}{2}\delta(\partial_a g_{bc} + \partial_b g_{ca} - 2\partial_c g_{ab})$$

$$\delta(\partial_a g_{bc}) = D_a\delta g_{bc}$$

substitute this into  $\delta\mathcal{L}$ ,

$$\begin{aligned} \delta\mathcal{L} &= 2 \left( \frac{1}{2} D_c (D_a \delta g_{bd} - D_b \delta g_{da} - 2D_d \delta g_{ab}) - \frac{1}{2} D_d (D_a \delta g_{bc} - D_b \delta g_{da} - 2D_c \delta g_{ab}) \right) R^{abcd} \\ &= (D_c (D_a \delta g_{bd} - D_b \delta g_{da} - 2D_d \delta g_{ab}) - D_d (D_a \delta g_{bc} - D_b \delta g_{da} - 2D_c \delta g_{ab})) R^{abcd} \end{aligned}$$

to get similar indices and pull out a variation of the metric tensor let us make the substitution

$$(note that  $\mathbf{1}$  is an identity matrix):  $D_c D_a \delta g_{bd} \rightarrow (\mathbf{1}_b^n \mathbf{1}_d^m D_c D_a) \delta g_{nm}$ ,$$

$$\delta\mathcal{L} = (D_c (\mathbf{1}_b^n \mathbf{1}_d^m D_a + \mathbf{1}_d^n \mathbf{1}_a^m D_b - 2 \cdot \mathbf{1}_a^n \mathbf{1}_b^m D_d) - D_d (\mathbf{1}_b^n \mathbf{1}_c^m D_a + \mathbf{1}_c^n \mathbf{1}_a^m D_b - 2 \cdot \mathbf{1}_a^n \mathbf{1}_b^m D_c)) \delta g_{nm} R^{abcd}$$

remembering this is all taking place under an integral we can use integration by parts to

change from a derivative of  $\delta g_{nm}$  to a derivative of  $R^{abcd}$ ,

$$\delta\mathcal{L} = ((D_c (\mathbf{1}_b^n \mathbf{1}_d^m D_a + \mathbf{1}_d^n \mathbf{1}_a^m D_b - 2 \cdot \mathbf{1}_a^n \mathbf{1}_b^m D_d) - D_d (\mathbf{1}_b^n \mathbf{1}_c^m D_a + \mathbf{1}_c^n \mathbf{1}_a^m D_b - 2 \cdot \mathbf{1}_a^n \mathbf{1}_b^m D_c)) R^{abcd}) \delta g_{nm}$$

$$\frac{\delta\mathcal{L}}{\delta g_{nm}} = ((D_c (\mathbf{1}_b^n \mathbf{1}_d^m D_a + \mathbf{1}_d^n \mathbf{1}_a^m D_b - 2 \cdot \mathbf{1}_a^n \mathbf{1}_b^m D_d) - D_d (\mathbf{1}_b^n \mathbf{1}_c^m D_a + \mathbf{1}_c^n \mathbf{1}_a^m D_b - 2 \cdot \mathbf{1}_a^n \mathbf{1}_b^m D_c)) R^{abcd})$$

distributing  $R^{abcd}$ ,

$$\begin{aligned} \frac{\delta\mathcal{L}}{\delta g_{nm}} &= D_c \mathbf{1}_b^n \mathbf{1}_d^m D_a R^{abcd} + D_c \mathbf{1}_d^n \mathbf{1}_a^m D_b R^{abcd} - 2D_c \mathbf{1}_a^n \mathbf{1}_b^m D_d R^{abcd} \\ &\quad - D_d \mathbf{1}_b^n \mathbf{1}_c^m D_a R^{abcd} - D_d \mathbf{1}_c^n \mathbf{1}_a^m D_b R^{abcd} + 2D_d \mathbf{1}_a^n \mathbf{1}_b^m D_c R^{abcd} \\ &= D_c D_a R^{ancm} + D_c D_b R^{mbcn} - 2D_c D_d R^{nmdc} \\ &\quad - D_d D_a R^{anmd} - D_d D_b R^{mbnd} + 2D_d D_c R^{nmdc} \end{aligned}$$

reordering some indices and observing symmetries of the Riemann tensor,

$$\begin{aligned}\frac{\delta\mathcal{L}}{\delta g_{nm}} &= D_b D_a R^{namb} - D_b D_a R^{manb} - 2D_a D_b R^{nmab} \\ &\quad + D_b D_a R^{namb} - D_a D_b R^{mbna} + 2D_b D_a R^{nmab} \\ \frac{\delta\mathcal{L}}{\delta g_{nm}} &= (D_b D_a R^{namb} + D_b D_a R^{namb}) - (D_b D_a R^{manb} - D_a D_b R^{mbna}) - (2D_a D_b R^{nmab} + 2D_b D_a R^{nmab}) \\ &= 2D_b D_a R^{namb} - 2D_b D_a R^{manb} - 2D_{[a} D_{b]} R^{nmab} \\ &= -2D_{[b} D_{a]} R^{namb} - 2D_{[a} D_{b]} R^{nmab} \\ &= -2D_{[a} D_{b]} (R^{namb} + R^{nmab}) \\ D_{[a} D_{b]} R^{namb} &= -\frac{1}{2} \mathcal{R}_{ab} R^{namb}\end{aligned}$$

$$\boxed{\frac{\delta\mathcal{L}}{\delta g_{nm}} = \mathcal{R}_{ab} R^{namb} + 2D_{[a} D_{b]} R^{nmab}}$$

### Appendix C: Plot Orbital Velocity vs. Radius for Characteristic Radii

The following is the code used to create the plot in figure 1

```

1 #!/usr/bin/env python
2 # coding: utf-8
3
4 # In [1]:
5
6
7 import numpy as np
8 import matplotlib.pyplot as plt
9 from matplotlib.lines import Line2D
10
11
12 # In [2]:
13
14
15 G = 6.67 * (10**(-11)) #Newtons const [m^3][kg^-1][s^-2]
```

```

16 M = 2 * (10**(41)) #average mass of spiral galaxy [kg]
17 #4.63 * (10**(20)) #average radius of spiral galaxy [m]
18 rho_0 = 10**(-21) #average central mass density of a spiral galaxy
    [kg][m^-3]
19
20
21 # In[3]:
22
23
24 # Define the parameters
25 char_raddii = [10, 1000, 1000000, 1] # Example values for r_d
26
27
28 # In[ ]:
29
30
31 # Define the function
32 def v_vs_r(r, r_d):
33     return np.sqrt((G*M)/(r) + (G * M * r * (np.log(r_d)/r_d)) *
34                 np.exp(-np.log(r_d)/r_d * r))
35
36 # Generate r values
37 # np.linspace(min, max, steps)
38 r_values = np.linspace(0.1, 10000000, 100000000)
39
40 # Plot the function for each r_d
41 for r_d in char_raddii:
42     v_values = v_vs_r(r_values, r_d)
43     plt.plot(r_values, v_values, label=f'r_d={r_d}' + ('-(
44         Keplarian)' if r_d==1 else '' ), linestyle=(0, (4, 5)) if
45         r_d==1 else '-', color='black' if r_d==1 else None)

```

```

43
44 # # plot newtonian for comparision
45 # v_values = v_vs_r(r_values, 0, 1)
46 # plt.plot(r_values, v_values, label=f'r_d = 1 (Newtonian)',
    linestyle=(0, (4, 5)), color='black')
47
48 plt.yscale('log') # Set the y-axis to logarithmic scale
49 plt.xscale('log') # Set the x-axis to logarithmic scale
50
51 plt.xlabel('r (m)')
52 plt.ylabel('Orbital Velocities (m/s)')
53 plt.title('Orbital Velocites vs. r')
54 plt.legend(title="Characteristic Radii (m)", loc="lower-left")
55 plt.grid(True)
56
57 plt.savefig('figures/graph_orbvelVr.png', dpi=300, bbox_inches='
    tight')
58
59 plt.show()
60
61
62 # In[ ]:
63
64
65 import subprocess
66 subprocess.call("jupyter nbconvert --to-script velocity_curves.
    ipynb")

```

---

[1] A. Einstein, *Annalen der Physik* **49**, 769 (1916).

[2] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Princeton University Press,

- 2017).
- [3] J. Brandes, in *The Physics of Reality: Space, Time, Matter, Cosmos* (World Scientific, 2013) pp. 120–122.
  - [4] S. Weinberg, *Reviews of Modern Physics* **61**, 1 (1989).
  - [5] M. P. Hobson, G. P. Efstathiou, and A. N. Lasenby, *General Relativity: An Introduction for Physicists* (Cambridge University Press, 2006).
  - [6] G. Bertone and D. Hooper, *Reviews of Modern Physics* **90**, 045002 (2018).
  - [7] J. A. Frieman, M. S. Turner, and D. Huterer, *Annual Review of Astronomy and Astrophysics* **46**, 385 (2008).
  - [8] R. M. Wald, *General Relativity* (University of Chicago Press, 1984).
  - [9] C. Rovelli, *Quantum Gravity* (Cambridge University Press, 2004).
  - [10] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, 1973).
  - [11] A. Einstein, *Sitzungsberichte der Preussischen Akademie der Wissenschaften zu Berlin* , 844 (1915).
  - [12] D. Hilbert, *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse* **1915**, 395 (1915).
  - [13] T. Watanabe and M. J. Hayashi, arXiv preprint gr-qc/0409029 (2004).
  - [14] E. Cartan, *Comptes Rendus de l'Academie des Sciences* **174**, 593 (1922).
  - [15] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Westview Press, 1995).
  - [16] S. Weinberg, *The Quantum Theory of Fields, Volume 1: Foundations* (Cambridge University Press, 1995).
  - [17] R. Hauser, Introduction to the standard model of particle physics (part 1) (2019), colorado School of Mines, Alex Flournoy, PHGN 423 Particle Physics, 2018, Lecture Notes.
  - [18] R. Hauser, Introduction to the standard model of particle physics (part 2) (2019), colorado School of Mines, Alex Flournoy, PHGN 423 Particle Physics, 2018, Lecture Notes.
  - [19] R. Hauser, Introduction to general relativity (2019), colorado School of Mines, Alex Flournoy, PHGN 418 General Relativity, 2018, Lecture Notes.
  - [20] D. Philipp, V. Perlick, C. Lämmerzahl, and K. Deshpande, in *2015 IEEE Metrology for Aerospace (MetroAeroSpace)* (IEEE, 2015) pp. 198–203.

- [21] M. Ganiou, I. G. Salako, M. Houndjo, and J. Tossa, *International Journal of Theoretical Physics* **55**, 3954 (2016).
- [22] K. L. Duggal and R. Sharma, *Symmetries of Spacetimes and Riemannian Manifolds* (Springer, 2013).
- [23] R. Emparan and H. S. Reall, *Physical Review D* **66**, 104025 (2002).
- [24] P. Jizba, L. Rachwał, and J. Kňap, *Physical Review D* **102**, 044033 (2020).
- [25] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt, *Exact solutions of Einstein's field equations* (Cambridge university press, 2009).
- [26] S. Hollands and W. Ruan, in *Annales Henri Poincare*, Vol. 3 (Springer, 2002) pp. 635–657.
- [27] Z. Bern, *Living Reviews in Relativity* **5**, 1 (2002).
- [28] A. Einstein and N. Rosen, *Journal of the Franklin Institute* **223**, 43 (1937).
- [29] J. T. Wheeler, *General Relativity and Gravitation* **50**, 1 (2018).
- [30] S. Hofmann, F. Niedermann, and R. Schneider, *Physical Review D—Particles, Fields, Gravitation, and Cosmology* **88**, 064047 (2013).
- [31] J. T. Wheeler, *Nuclear Physics B* **273**, 732 (1986).
- [32] J. Panei, L. Althaus, and O. Benvenuto, arXiv preprint astro-ph/9909499 (1999).
- [33] J. D. Barrow and D. J. Shaw, *International Journal of Modern Physics D* **20**, 2875 (2011).
- [34] C. Caprini and D. G. Figueroa, *Classical and Quantum Gravity* **35**, 163001 (2018).
- [35] M. D. Kruskal, *Physical review* **119**, 1743 (1960).
- [36] C. M. Will, *Theory and Experiment in Gravitational Physics* (Cambridge University Press, 1993).
- [37] D. Lai, L. Bildsten, and V. Kaspi, arXiv preprint astro-ph/9505042 (1995).
- [38] N. Wex, S. Johnston, R. Manchester, A. Lyne, B. Stappers, and M. Bailes, *Monthly Notices of the Royal Astronomical Society* **298**, 997 (1998).
- [39] L. S. Collaboration and V. Collaboration, *The Astrophysical Journal* **892**, L3 (2020).
- [40] L. S. Collaboration and V. Collaboration, *Classical and Quantum Gravity* **38**, 225009 (2021).
- [41] L. S. Collaboration and V. Collaboration, *The Astrophysical Journal* **913**, 1 (2021).
- [42] D. Burstein and V. C. Rubin, *Astrophysical Journal, Part 1 (ISSN 0004-637X)*, vol. 297, Oct. 15, 1985, p. 423-435. **297**, 423 (1985).
- [43] K. C. Freeman, *Astrophysical Journal*, vol. 160, p. 811 **160**, 811 (1970).
- [44] J. F. Navarro, C. S. Frenk, and S. D. White, *The Astrophysical Journal* **490**, 493 (1997).

- [45] J. R. Brownstein and J. W. Moffat, *The Astrophysical Journal* **636**, 721 (2006).
- [46] S. Deser, arXiv preprint arXiv:0910.2975 (2009).
- [47] M. Reuter and H. Weyer, *General Relativity and Gravitation* **41**, 983 (2009).
- [48] A. Ashtekar, *General Relativity and Gravitation* **41**, 1927 (2009).
- [49] A. Ashtekar and J. Lewandowski, *Classical and Quantum Gravity* **21**, R53 (2004).
- [50] J. Read, *Background independence in classical and quantum gravity* (Oxford University Press, 2023).
- [51] S. Deser, *Reviews of Modern Physics* **29**, 417 (1957).